

International
Online Conference



**Algebraic
and Geometric
Methods of Analysis**

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Drozd Yu.

(Kyiv, Ukraine)

Maksymenko S.

(Kyiv, Ukraine)

Plaksa S.

(Kyiv, Ukraine)

Prishlyak A.

(Kyiv, Ukraine)

Bakhtin O.

(Kyiv, Ukraine)

Balan V.

(Bucharest, Romania)

Banakh T.

(Lviv, Ukraine)

Borysenko O.

(Kharkiv, Ukraine)

Cherevko Ye.

(Odesa, Ukraine)

Fedchenko Yu.

(Odesa, Ukraine)

Karlova O.

(Chernivtsi, Ukraine)

Kiosak V.

(Odessa, Ukraine)

Konovenko N.

(Odessa, Ukraine)

Lyubashenko V.

(Kyiv, Ukraine)

Matsumoto K.

(Yamagata, Japan)

Mormul P.

(Warsaw, Poland)

Mykhailyuk V.

(Chernivtsi, Ukraine)

Plachta L.

(Krakov, Poland)

Pokas S.

(Odessa, Ukraine)

Sabitov I.

(Moscow, Russia)

Savchenko O.

(Kherson, Ukraine)

Sergeeva A.

(Odessa, Ukraine)

Shelekhov A.

(Tver, Russia)

Zarichnyi M.

(Lviv, Ukraine)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

ORGANIZING COMMITTEE

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohlyk O.

Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.

Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

Reducing singularities of smooth functions to normal forms

Orevkova Alexandra

(Moscow State University, Moscow, Russia)

E-mail: s15b3_orevkova@179.ru

The talk is devoted to the “uniform” reduction of C^∞ smooth functions on 2-dimensional manifolds to a canonical form at the singular points of these functions.

Definition 1. A smooth function $f = f(u_1, u_2)$ has a singularity E_6 at its critical point $P \in \mathbb{R}^2$ if

- (i) the first and second differentials $df(P) = 0$, $d^2f(P) = 0$, and the third differential $d^3f(P) \neq 0$ and is a perfect cube;
- (ii) there exists a vector $v \in \text{Ker } d^3f(P)$ such that $v^4f \neq 0$ (by v^4f we mean the fourth derivative of f along the tangent vector v at P).

Theorem 2 (Reducing E_6 to normal form). *Let the function $f(u_1, u_2)$ have a singularity E_6 at the critical point P . Then, in some neighborhood of P , there is a local coordinate system in which the point P is the origin, and the function has the normal form*

$$f = f(P) + \tilde{x}^3 \pm \tilde{y}^4.$$

Moreover, this coordinate system can be chosen in such a way that the coordinate change $(u_1, u_2) \rightarrow (\tilde{x}, \tilde{y})$ can be expressed in terms of the original function and its partial derivatives of order ≤ 7 using algebraic operations and the operation of taking a proper integral.

Remark 3. In [1], the first part of Theorem 2, i.e. the existence of a coordinate change, was proved using Tougeron’s theorem [2]; in view of this, obtaining a formula for the corresponding coordinate change requires solving the Cauchy problem for a system of ordinary differential equations. We construct our coordinate change explicitly, without using Tougeron’s theorem.

Our proof of Theorem 2 consists of three steps. At first, we consider a smooth function in two variables of the form $f(u_1, u_2) = u_1^3 + u_2^4 + R(u_1, u_2)$, where the Taylor series of the function $R(u_1, u_2)$ at the origin has zero coefficients at all monomials of the form $u_1^k u_2^l$, $4k + 3l \leq 12$. At second, by the sequence $(u_1, u_2) \rightarrow (x, y)$ of substitutions described in [1], we reduce the function f to such a form that the origin leaves fixed and $f_{x^k y^l}^{(k+l)}(0, 0) = 0$, where $0 \leq k < 3, 0 \leq l < 4$. At third, using the Taylor series expansion of the function with an integral remainder, we reduce the function to the required normal form by the coordinate change

$$\phi : \mathbb{R}_{x,y}^2 \rightarrow \mathbb{R}_{\tilde{x},\tilde{y}}^2 \quad \text{with} \quad \tilde{x} = x \sqrt[3]{g(x,y)}, \quad \tilde{y} = y \sqrt[4]{h(x,y)},$$

where $g(x, y) := \sum_{k=0}^3 \frac{y^k}{k!} f_{y^k}^{(k)}(x, 0)/x^3$ and $h(x, y) := \frac{1}{6} \int_0^1 f_{y^4}^{(4)}(x, sy)(1-s)^3 ds$.

Our next goal is to describe a neighborhood in which the above coordinate change ϕ exists and is regular. To simplify our computations, we will assume that the following is true:

Assumption 4. $f_{y^4}^{(4)}(0, 0) = 24$, $f_{x^3}'''(0, 0) = 6$.

Theorem 5 (Estimating the radius of a neighborhood where the coordinate change is regular). *Under the hypotheses of Theorem 2 and Assumption 4, let $U_0 = \{(x, y) \mid \max(|x|, |y|) < R_0\}$ be a neighborhood of the origin such that $C_{\alpha\beta} = \sup_{U_0} |f_{x^\alpha y^\beta}^{\alpha+\beta}(x, y)| \leq M$ for $(\alpha, \beta) \in \{(0, 5), (1, 4), (3, 1), (3, 2), (3, 3), (4, 0), (4, 1), (4, 2), (4, 3)\}$, where $R_0 > 0$, $M \geq 0$. Let’s consider the neighborhood $U = \{(x, y) \mid \max(|x|, |y|) < R\}$, where the positive constant R is defined by the formula: $R = \min\{R_0, \frac{2}{M+2}\}$. Let also ϕ be the coordinate change from the proof of Theorem 2 that reduces f to the normal form E_6 . Then:*

- (a) the functions $h(x, y)$ and $g(x, y)$ do not change sign in U , that is, the change $\phi|_U$ is well defined and is smooth;
- (b) at every point $\mathbf{x} \in U$, one has $\|\phi'(\mathbf{x}) - I\| < C < 1$, where $C = \frac{2}{5}$, i.e., the coordinate change $\phi|_U$ is close to the identity;
- (c) the coordinate change $\phi|_U$ is injective and regular, i.e., it is an embedding and $\det |\phi'(\mathbf{x})| \neq 0$ at every point $\mathbf{x} \in U$, moreover the image of this embedding contains the disk of radius $(1 - C)R$ centred at the origin.

Remark 6. Our coordinate change $\phi|_U$ from Theorem 2 and Theorem 5 provides a “uniform” reduction of the function f at a singular point of type E_6 to the canonical form $\tilde{x}^3 \pm \tilde{y}^4$ in the sense that the neighbourhood radius and the coordinate change we constructed in this neighbourhood (as well as all partial derivatives of the coordinate change) continuously depend on the function f and its partial derivatives. A uniform reduction of smooth functions near critical points to a canonical form was known earlier for the case of smoothly stable singularities [3]. A uniform reduction of smooth functions to a canonical form by C^k -smooth changes (for finite $k < \infty$) is known for finite type singularities [3] and for topologically stable singularities [5].

To prove Theorem 5, we apply the following lemma to the coordinate transformation ϕ from the proof of Theorem 2. We estimate the norm of the matrix in terms of its elements: $\|A\| \leq \sqrt{\sum a_{ij}^2}$, $A = \{a_{ij}\}_{i,j=1}^n$.

Lemma 7. Let $\phi : U \rightarrow \mathbb{R}^n$ be a smooth mapping, where U is a convex open subset of \mathbb{R}^n . Let the differential of ϕ have the form $\phi'(\mathbf{x}) = I + A(\mathbf{x})$, where I is the unit matrix of dimension n , $\|A\| < C$, $0 < C < 1$. Then ϕ is injective and $\det |\phi'(\mathbf{x})| \neq 0$ at every point $\mathbf{x} \in U$, i.e., ϕ is a diffeomorphism to its image $\phi(U)$. Moreover, $\langle \phi'(\mathbf{x}), \mathbf{x} \rangle \geq (1 - C)|\mathbf{x}|^2$ for every point $\mathbf{x} \in U$.

From the last assertion of Lemma 7 and [6, Corollary 8.3, Step 1], we conclude that $\phi(U)$ contains the disk of radius $(1 - C)R$ centred at the origin. This completes our proof of Theorem 5.

REFERENCES

- [1] Vladimir I Arnol'd. Normal forms for functions near degenerate critical points, the Weyl groups of A_k, D_k, E_k and Lagrangian singularities. *Funct. Anal. Appl.*, 6(4) : 254–272, 1972.
- [2] Jean-Claude Tougeron. Ideaux de fonctions différentiables. I. *Ann. Inst. Fourier*, 18(1) : 177–240, 1968.
- [3] John N Mather. Infinitesimal stability implies stability. *Ann. of Math.*, 89 : 254–291, 1969.
- [4] Anatoly M Samoilenko. The equivalence of a smooth function to a Taylor polynomial in the neighborhood of a finite-type critical point. *Funct. Anal. Appl.*, 2(4) : 318–323, 1968.
- [5] Hans Brodersen. M-t topologically stable mappings are uniformly stable. *Math. Scand.*, 52 : 61–68, 1983.
- [6] Elena A Kudryavtseva, Dmitry A Permyakov. Framed Morse functions on surfaces. *Sbornik Math.*, 201(4) : 501–567, 2010.

Kh. F. Kholturayev <i>Perfect metrizable of the functor of idempotent measures</i>	75
Y. Khomych <i>Quasiareal deformation of surfaces of positive Gauss curvature</i>	77
V. Kiosak, O. Lesechko <i>Canonical infinitesimal deformations of metrics of pseudo-Riemannian spaces</i>	78
R. Salimov, B. Klishchuk <i>On the behavior at infinity of ring Q-homeomorphisms</i>	79
T. Kolomiets, A. Pogorui <i>Elements of probability theory and measures with values in hypercomplex algebras</i>	81
N. Konovenko <i>The invariants of planar 3-webs with respect to group of symplectic diffeomorphisms, for the case of the conformal group</i>	84
E. Kudryavtseva <i>Topology of spaces of smooth functions and gradient-like flows with prescribed singularities on surfaces</i>	85
G. Kuduk <i>Nonlocal problem with integral conditions for homogeneous system of partial differential equations of second order</i>	87
I. Kuznietsova, Yu. Soroka <i>Realization of groups as fundamental groups of orbits of smooth maps</i>	88
K. Gürlebeck, D. Legatiuk <i>Modified quaternionic operator calculus and its application to micropolar elasticity</i>	90
S. Maksymenko, E. Polulyakh <i>On non-Hausdorff manifolds of dimension 1</i>	92
S. Maksymenko <i>Symplectomorphisms preserving smooth functions on surfaces</i>	93
M. Maloid-Hliebova <i>Second classical Zariski topology of multiplicative module</i>	94
I. Marko <i>Incomplete spaces of idempotent measures</i>	95
N. Mazurenko, M. Zarichnyi <i>Hyperspaces of convex sets related to idempotent mathematics</i>	96
A. Mednykh <i>Volumes of knots and links in spaces of constant curvature</i>	98
R. Mohseni, R. A. Wolak <i>Twistor spaces on foliated manifolds</i>	99
P. Mormul <i>Two problems in nonholonomic geometry (in quest of a co-worker)</i>	100
F. Mukhamadiev <i>The local τ-density of a linearly ordered spaces</i>	101
T. Obikhod <i>Entropy and phase transitions in Calabi-Yau space</i>	102
A. Orevkova <i>Reducing singularities of smooth functions to normal forms</i>	104
T. Osipchuk <i>On m-convexity and m-semiconvexity of sets in Euclidean spaces</i>	106
V. Ostrovskiy, O. Ostrovska, D. Proskurin, Yu. Samoilenko <i>On representations of q_{ij}-commuting isometries</i>	108
J.F. Peters <i>Homotopic Nerve Complexes with Free Group Presentations</i>	110
P. Laurain, M. Petrace <i>Uniform measures in Euclidean space</i>	112