

International
Scientific Conference



Algebraic
and Geometric
Methods
of Analysis

27-30 May 2024
Odesa, Ukraine

The purpose of this conference is to bring together researchers in geometry, topology, algebra, analysis and dynamical systems and to provide for them a forum to present their recent work to colleagues from different nationalities. This way we aim to stimulate discussion about the latest findings in geometrical and topological methods in analysis and to increase international collaboration.

The conference continues the traditional annual conference «Geometry in Odesa» holding from 2004, and hosted by Odesa National University of Technology (Odesa National Academy of Food Technologies till 2021). From 2017 the conference was renamed to «Algebraic and geometric methods of analysis» (AGMA).

The Conference languages: Ukrainian and English.

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology, Ukraine
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

- | | |
|---|--|
| • Vladimir Balan (<i>Bucharest, Romania</i>) | • Volodymyr Lyubashenko (<i>Kyiv, Ukraine</i>) |
| • Taras Banakh (<i>Lviv, Ukraine</i>) | • Sergiy Maksymenko (<i>Kyiv, Ukraine</i>) |
| • Dmytro Bolotov (<i>Kharkiv, Ukraine</i>) | • Koji Matsumoto (<i>Yamagata, Japan</i>) |
| • Vyacheslav Boyko (<i>Kyiv, Ukraine</i>) | • Piotr Mormul (<i>Warsaw, Poland</i>) |
| • Yulia Fedchenko (<i>Odesa, Ukraine</i>) | • Maryna Nesterenko (<i>Kyiv, Ukraine</i>) |
| • Oleg Gutik (<i>Lviv, Ukraine</i>) | • Roman Popovych (<i>Kyiv, Ukraine</i>) |
| • Olena Karlova (<i>Chernivtsi, Ukraine</i>) | • Alexandr Prishlyak (<i>Kyiv, Ukraine</i>) |
| • Volodymyr Kiosak (<i>Odesa, Ukraine</i>) | • Aleksandr Savchenko (<i>Kherson, Ukraine</i>) |
| • Nadiia Konovenko (<i>Odesa, Ukraine</i>) | |

ORGANIZING COMMITTEE

- | | |
|---|--|
| • Nadiia Konovenko (<i>Odesa, Ukraine</i>) | • Bohdan Mazhar (<i>Kyiv, Ukraine</i>) |
| • Yuliya Fedchenko (<i>Odesa, Ukraine</i>) | • Sergiy Maksymenko (<i>Kyiv, Ukraine</i>) |
| • Mykola Lysynskiy (<i>Kyiv, Ukraine</i>) | • Alexandr Prishlyak (<i>Kyiv, Ukraine</i>) |

Solutions of N -body harmonic oscillators and Calogero-Moser model using Φ^4 matrix model

Akifumi Sako

(Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo, 162-8601, Japan)

E-mail: sako@rs.tus.ac.jp

The N -body harmonic oscillator system and its generalized system, the Calogero-Moser model, are known as quantum integrable systems, i.e. their Schrödinger equations are solvable and eigenstates of the Hamiltonians can be constructed. It has long been known that there is a connection between these quantum integrable systems and certain kind of generalized matrix models. This talk is concerned with the new correspondence of the quantum solvable systems with some matrix models, which is discovered last year. These matrix models are given as the Grosse-Wulkenhaar models, known as renormalizable scalar Φ^4 -theories on Moyal spaces, which are non-commutative spaces. The Moyal space has Fock representation, so field theories can be expressed by using matrix representation. A scalar Φ^4 -theory on Moyal space corresponds to a Hermitian Φ^4 -matrix model or a real symmetric Φ^4 -matrix model. In particular, Φ^4 -matrix model known as the Grosse-Wulkenhaar models have kinetic terms $\text{Tr}(E\Phi^2)$, where E is a positive diagonal matrix without degenerate eigenvalues. (These matrix models are also obtained by changing the potential of the Kontsevich model from Φ^3 to Φ^4 .) We show that their partition functions of these matrix models correspond to zero-energy solutions of a Schrödinger type equation with the N -body harmonic oscillator Hamiltonian and the Calogero-Moser Hamiltonian, respectively.

Let Φ be a Hermitian or real symmetric $N \times N$ matrix. Let $Z(E, \eta)$ be the partition function defined by

$$Z(E, \eta) = \int d\Phi e^{-S_E[\Phi]}, \quad (1)$$

where $S_E = N \text{Tr}\{E\Phi^2 + \frac{\eta}{4}\Phi^4\}$ and η is a real number. The domains of integrations are the space of Hermitian $N \times N$ -matrices and the space of real symmetric $N \times N$ -matrices, respectively. Let $\Delta(E)$ be the Vandermonde determinant $\Delta(E) := \prod_{k < l} (E_l - E_k)$. Using these, the theorem obtained can be described as follows.

Theorem 1. *Let $\Psi(E, \eta)$ be a function defined by*

$$\Psi(E, \eta) := e^{-\frac{N}{\beta\eta} \sum_{i=1}^N E_i^2} \Delta(E)^{\frac{\beta}{2}} Z(E, \eta).$$

Then $\Psi(E, \eta)$ is a zero-energy solution of the Schrödinger type equation

$$\mathcal{H}\Psi(E, \eta) = 0.$$

Here \mathcal{H} is the Hamiltonian \mathcal{H}_{HO} for the N -body harmonic oscillator system when we consider the Hermitian matrix model with $\beta = 2$:

$$\mathcal{H}_{HO} := -\frac{\eta}{N} \sum_{i=1}^N \left(\frac{\partial}{\partial E_i} \right)^2 + \frac{N}{\eta} \sum_{i=1}^N (E_i)^2.$$

When we consider the real symmetric matrix model with $\beta = 1$, then \mathcal{H} is the Hamiltonian \mathcal{H}_{CM} for Calogero-Moser model:

$$\mathcal{H}_{CM} := \frac{-\eta}{2N} \left(\sum_{i=1}^N \frac{\partial^2}{\partial E_i^2} + \frac{1}{4} \sum_{i \neq j} \frac{1}{(E_i - E_j)^2} \right) + 2 \frac{N}{\eta} \sum_{i=1}^N E_i^2. \quad (2)$$

It is known that the N -body harmonic oscillator system or the Calogero-Moser model is associated with a Virasoro algebra structure. Using this fact, families of differential equations satisfied by the partition functions are also obtained from the Virasoro(Witt) algebra representations:

$$[\tilde{L}_n, \tilde{L}_m] = (n - m) \tilde{L}_{n+m}. \quad (3)$$

The definitions of symbols and terms are left to the references [1, 2], but the following theorem is obtained

Theorem 2. *The partition function defined by (1) satisfies*

$$\mathcal{L}_{SD}(\tilde{L}_{-m}Z(E, \eta)) = -2m(\tilde{L}_{-m}Z(E, \eta)). \quad (4)$$

Here \mathcal{L}_{SD} is a differential operator such that some Schwinger-Dyson equation for the partition function given by

$$\mathcal{L}_{SD}Z(E, \eta) = 0. \quad (5)$$

This means that $\tilde{L}_{-m}Z(E, \eta)$ is an eigenfunction of \mathcal{L}_{SD} with the eigenvalue $-2m$.

This talk is based on [1], in collaboration with H. Grosse, and [2], in collaboration with H. Grosse, N. Kanomata, and R. Wulkenhaar.

REFERENCES

- [1] H. Grosse and A. Sako, "Integrability of Φ^4 Matrix Model as N -body Harmonic Oscillator System," Lett Math Phys 114, 48 (2024). <https://doi.org/10.1007/s11005-024-01783-2> [arXiv:2308.11523 [math-ph]].
- [2] H. Grosse, N. Kanomata, A. Sako and R. Wulkenhaar, "Real symmetric Φ^4 -matrix model as Calogero-Moser model," Lett Math Phys 114, 25 (2024). <https://doi.org/10.1007/s11005-024-01772-5> [arXiv:2311.10974[hep-th]].

Partitioning problem and defensive alliances in the context of zero-divisor graphs of rings

Brahim El Alaoui

(Mohammed V University-Faculty of Sciences in Rabat, Rabat, Morocco)

E-mail: brahim_elalaoui2@um5.ac.ma

This is joint work with **Driss Bennis**

The partitioning of the vertex set of a graph is a well-studied problem in graph theory. It involves dividing the set of vertices of a graph into disjoint subsets or partitions, based on specific criteria or constraints. In this talk, we are interested in partitioning the zero-divisor graph of a commutative ring into global defensive alliances. This problem has been well investigated in graph theory. Here, we connected it with the ring theoretical context. We characterize various finite commutative rings for which the zero-divisor graph is partitionable into global defensive alliances. We also present several examples to illustrate and delimit the scope of the established results.

Table of contents

A. Sako <i>Solutions of N-body harmonic oscillators and Calogero-Moser model using Φ^4 matrix model</i>	2
B. El Alaoui <i>Partitioning problem and defensive alliances in the context of zero-divisor graphs of rings</i>	3
M. Amram <i>On the connection between algebraic, geometric, and topological methods in the classification of algebraic surfaces and curves</i>	4
N. Ando <i>Nilpotent structures of oriented neutral vector bundles</i>	6
M.J. Atteya <i>Multiplicative b-homogeneralized Derivations of Associative Rings</i>	8
T. Banakh <i>Algebra in fields extended by infinity</i>	9
M. Bisci <i>Variational problems in Nonsmooth Analysis</i>	10
D. Bolotov <i>On 2-convex embeddings of non-orientable surfaces in four-dimensional Euclidean space</i>	11
E. Bonacci <i>Mixing optimization in the batch crystallization of CAM</i>	11
V. Bondarenko, M. Styopochkina <i>On representation type of incident algebras of extensions of positive posets</i>	12
F. Bulnes <i>A theorem on hypercohomology groups and singular homology in field theory</i>	13
D. Carfi <i>Geometric and algebraic-topological structures in Schwartz distribution spaces for relativistic Quantum Mechanics</i>	13
Y. Chapovskyi, S. Koval, O. Zhur <i>Lie subalgebras of real order-three special linear Lie algebra revisited</i>	16
Y. Cherevko, O. Chepurna, Y. Kuleshova <i>Conformal mappings and a non-holonomic frame</i>	18
A. Chornenka, O. Gutik <i>On topologization of the bicyclic monoid</i>	19
J. Cuadros, J. Lope <i>An Application to Sasaki Extremal metrics via the Berglund-Hübsch rule</i>	20
E. Sevost'yanov, V. Desyatka <i>On singularities of mappings with a Lebesgue integrable majorant</i>	22
I. Diamantis, S. Lambropoulou, S. Mahmoudi <i>New Combinatorial Invariants of Doubly Periodic Tangles</i>	24
K. v. Dichter <i>Inequalities involving means in high-dimensional spaces</i>	25
P. Petrenko, A. Andreev <i>Two problems in the theory of metric preserving functions</i>	27
Y. Drozd <i>Group action on noncommutative curves</i>	29