

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

Inverse three spectra problem for a Stieltjes string with the Neumann boundary conditions

A. Dudko

(South Ukrainian National Pedagogical University named after K.D. Ushynsky)

E-mail: nastysha00301@gmail.com

V. Pivovarchik

(South Ukrainian National Pedagogical University named after K.D. Ushynsky)

E-mail: vpivovarchik@gmail.com

The notion of Stieltjes string was introduced in [1, Supplement II]. Like in [1] we suppose the string to be a thread, i. e. a string of zero density, bearing a finite number of point masses. Assume that the string consists of two parts, which are joined at one end and free to move in the direction orthogonal to the equilibrium position of the string at the other end. Starting indexing from the free ends, n_j masses $m_k^{(j)} > 0$, $k = 1, \dots, n_j$, are positioned on the j -th part, $j = 1, 2$, which divide the j -th part into $n_j + 1$ substrings, denoted by $l_k^{(j)} > 0$, $k = 0, \dots, n_j$, again starting indexing from the free end. In particular, $l_0^{(j)}$ is the distance on the j -th part between the free endpoint and $m_1^{(j)}$, $l_1^{(j)}$ for $k = 1, \dots, n_j - 1$ is the distance between $m_k^{(j)}$ and $m_{k+1}^{(j)}$, and $l_{n_j}^{(j)}$ is the distance on the j -th thread between the joined end point and $m_{n_j}^{(j)}$. The tension of the thread is assumed to be equal to 1.

Considering small transverse vibrations of such string like in [2, p.55] we obtain the following spectral problem

$$\frac{u_k^{(j)} - u_{k+1}^{(j)}}{l_k^{(j)}} + \frac{u_k^{(j)} - u_{k-1}^{(j)}}{l_{k-1}^{(j)}} - m_k^{(j)} z u_k^{(j)} = 0 \quad (k = 1, 2, \dots, n_j, \quad j = 1, 2), \quad (1)$$

$$u_{n_1+1}^{(1)} = u_{n_2+1}^{(2)}, \quad (2)$$

$$\sum_{j=1}^2 \frac{u_{n_j}^{(j)} - u_{n_j+1}^{(j)}}{l_{n_j}^{(j)}} = 0, \quad (3)$$

$$u_0^{(j)} = u_1^{(j)}, \quad j = 1, 2. \quad (4)$$

where $u_k^{(j)}$ is the amplitude of vibrations of the mass $m_k^{(j)}$, z is the spectral parameter. Also we consider following two problems for the parts of the string:

$$\frac{u_k^{(j)} - u_{k+1}^{(j)}}{l_k^{(j)}} + \frac{u_k^{(j)} - u_{k-1}^{(j)}}{l_{k-1}^{(j)}} - m_k^{(j)} z u_k^{(j)} = 0 \quad (k = 1, 2, \dots, n_j), \quad (5)$$

$$u_0^{(j)} = u_1^{(j)}, \quad (6)$$

$$u_{n_j+1}^{(j)} = 0. \quad (7)$$

We denote by $\{\mu_k\}_{k=1}^n$ where $n = n_1 + n_2$ the spectrum of problem (1)–(4), and by $\{\nu_k^{(j)}\}_{k=1}^{n_j}$, $j = 1, 2$ the spectra of problems (5)–(7). Let $\{\xi_k\}_{k=1}^n = \{\nu_k^{(1)}\}_{k=1}^{n_1} \cup \{\nu_k^{(2)}\}_{k=1}^{n_2}$ be indexed such that $\xi_k \leq \xi_{k+1}$.

Location of the spectra is described by

Theorem 1. *The sequences $\{\mu_k\}_{k=1}^n$ and $\{\xi_k\}_{k=1}^n$ interlace as follows:*

- 1) $0 = \mu_1 < \xi_1 \leq \mu_2 \leq \dots \leq \xi_n$;
- 2) $\mu_k = \xi_{k-1}$ if and only if $\mu_k = \xi_k$;
- 3) the multiplicity of ξ_k does not exceed 2.

The corresponding inverse problem lies in finding the masses $\{m_k^{(j)}\}_{k=1}^{n_j}$ and subintervals $\{l_k^{(j)}\}_{k=1}^{n_j}$ ($j = 1, 2$) using the three spectra, i.e. spectra of problems (1)–(4), (5)–(7) with $j = 1$ and (5)–(7) with $j = 2$.

Theorem 2. *Let $m_j > 0$ ($j = 1, 2$) be given together with the numbers*

$\{\mu_k\}_{k=1}^n$, $\{\nu_k^{(1)}\}_{k=1}^{n_1}$ and $\{\nu_k^{(2)}\}_{k=1}^{n_2-1}$ ($\mu_k < \mu_{k'}, \nu_k^{(j)} < \nu_{k'}^{(j)}$ for $k < k'$) $\{\xi_k\}_{k=1}^{n-1} = \{\nu_k^{(1)}\}_{k=1}^{n_1} \cup \{\nu_k^{(2)}\}_{k=1}^{n_2-1}$ which satisfy the following conditions:

$$0 = \mu_1 < \xi_1 < \mu_2 < \xi_2 < \dots < \xi_{n-1} < \mu_n.$$

Then there exists a unique collection of sequences of positive numbers $\{\{m_k^{(j)}\}_{k=1}^{n_j},$

$(j = 1, 2), \{l_k^{(j)}\}_{k=1}^{n_j} \quad (j = 1, 2)\}$ such that $\sum_{k=1}^{n_j} m_k^{(j)} = m_j$, $\{\mu_k\}_{k=1}^n$ is the spectrum of problem

(1)–(4), $\{\nu_k^{(1)}\}_{k=1}^{n_1}$ is the spectrum of problem (5)–(7) with $j = 1$, $\{\nu_k^{(2)}\}_{k=1}^{n_2-1}$ is a sequence of eigenvalues (a part of the spectrum) of problem (5)–(7) with $j = 2$.

The method of recovering $\{\{m_k^{(j)}\}_{k=1}^{n_j}, \quad (j = 1, 2), \quad \{l_k^{(j)}\}_{k=1}^{n_j} \quad (j = 1, 2)\}$ is similar to that in [3] (see also [2], p. 222).

REFERENCES

- [1] F.R. Gantmakher, M.G. Krein. Oscillation matrices and kernels and small vibrations of mechanical systems. Revised edition. *AMS Chelsea Publishing, Providence, RI*, 2002.
- [2] M. Möller, V. Pivovarchik. Spectral Theory of Operator Pencils, Hermite-Biehler Functions, and their Applications *Birkhäuser*, 2015.
- [3] O. Boyko, V. Pivovarchik. The inverse three-spectra problem for a Stieltjes string and the inverse problem with one-dimensional damping. *Inverse problems*, volume 24, no.1, 015019 13 pp, 2008.

Зміст

N. Aygor, H. Burhanzade <i>Secondary school students' misconceptions about linear algebra</i>	3
S. Bardyla, H. Kvasnytsia <i>Semitopological graph inverse semigroups</i>	4
B. A. Bhayo <i>On inequalities of generalized elliptic integrals</i>	5
Bodzioch M., Choiński M., Foryś U. <i>A criss-cross model of tuberculosis for heterogenous population</i>	6
Bolotov D. V. <i>Foliations with leaves of non-positive curvature and bounded total curvature on closed 3-manifolds</i>	7
E. Bonacci <i>Algebraic and geometric questions about a 6D physics</i>	9
F. Bulnes <i>Mukai-Fourier Transform in Derived Categories to Solutions of the Field Equations: Gravitational Waves as Oscillations in the Space-Time Curvature/Spin IV</i>	10
H. Burhanzade, N. Aygor <i>A study on the teaching methods in determinants</i>	12
Damla Yaman <i>Order continuity properties of lattice ordered algebras</i>	13
Denega I. <i>Problem on non-overlapping polycylindrical domains with poles on the boundary of a polydisk</i>	14
A. Dudko, V. Pivovarchik <i>Inverse three spectra problem for a Stieltjes string with the Neumann boundary conditions</i>	16
Eftekhari Nasab K. <i>On the existence of a global diffeomorphism between Fréchet spaces</i>	18
Glazunov N. <i>Class groups of rings with divisor theory, L-functions and moduli spaces</i>	19
O. Gök <i>b-bimorphisms</i>	21
Gün E. <i>On the second regularized trace formula for a differential operator with unbounded coefficients</i>	22
Hentosh O. Ye., Prykaratsky Ya. A. <i>The Lie-algebraic structure of the Lax-Sato integrable superanalogs for the Liouville heavenly type equations</i>	24
V. Herasymov <i>In a natural topological sense a typical linear nonhomogeneous differential equation in the ring $Z[[x]]$ has no solutions from $Z[[x]]$.</i>	26
Juraev D. A. <i>On the Cauchy problem for matrix factorizations of the Helmholtz equation</i>	27
M. E. Kansu <i>Macroscopic electromagnetism via complex quaternions</i>	29
Vladimir V. Kisil <i>An extension of Möbius–Lie geometry with conformal ensembles of cycles</i>	30
Konovenko N., Lychagin V. <i>Rational differential invariants for oriented primary visual cortex</i>	32