

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

An extension of Möbius–Lie geometry with conformal ensembles of cycles

Vladimir V. Kisil

(School of Mathematics, University of Leeds, Leeds LS2 9JT, England)

E-mail: kisilv@maths.leeds.ac.uk

Lie sphere geometry in the simplest planar setup unifies circles, lines and points—all together called *cycles* in this setup [8]. Symmetries of Lie spheres geometry include (but are not limited to) fractional linear transformations (FLT) of the form:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} : x \mapsto \frac{ax + b}{cx + d}, \quad \text{where } \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} \neq 0. \quad (1)$$

There is a natural set of FLT-invariant geometric relations between cycles (to be orthogonal, to be tangent, etc.) and the restriction of Lie sphere geometry to invariants of FLT is called *Möbius–Lie geometry*. Thus, an ensemble of cycles, structured by a set of such relations, will be mapped by FLT to another ensemble with the same structure.

It is convenient to deal with cycles through Fillmore–Springer–Cnops construction (FSCc) which also has a long history, see [8, § 4.2]. Compared to a plain analytical treatment [3, Ch. 3], FSCc is much more efficient and conceptually coherent in dealing with FLT-invariant properties of cycles. For example, the inner product $\langle C_1, C_2 \rangle$ of cycles C_1 and C_2 is equal to zero if and only if cycles are orthogonal. Other properties, e.g. tangency, has a natural presentation as well, see [8, Ch. 4–5] and [9].

It was shown recently that ensembles of cycles with certain FLT-invariant relations provide helpful parametrisations of new objects as follows.

Example 1. (1) The Poincaré extension of Möbius transformations from the real line to the upper half-plane of complex numbers is described by a triple of cycles $\{C_1, C_2, C_3\}$ such that:

- (a) C_1 and C_2 are orthogonal to the real line;
- (b) $\langle C_1, C_2 \rangle^2 \leq \langle C_1, C_1 \rangle \langle C_2, C_2 \rangle$;
- (c) C_3 is orthogonal to any cycle in the triple including itself.

A modification [10] with ensembles of four cycles describes an extension from the real line to the upper half-plane of complex, dual or double numbers. The construction can be generalised to arbitrary dimensions.

(2) A parametrisation of loxodromes is provided by a triple of cycles $\{C_1, C_2, C_3\}$ such that [12]:

- (a) C_1 is orthogonal to C_2 and C_3 ;
- (b) $\langle C_2, C_3 \rangle^2 \geq \langle C_2, C_2 \rangle \langle C_3, C_3 \rangle$.

Then, main invariant properties of Möbius–Lie geometry, e.g. tangency of loxodromes, can be expressed in terms of this parametrisation [12].

(3) A continued fraction is described by an infinite ensemble of cycles (C_k) such that [2]:

- (a) All C_k are touching the real line (i.e. are *horocycles*);
- (b) (C_1) is a horizontal line passing through $(0, 1)$;
- (c) C_{k+1} is tangent to C_k for all $k > 1$.

This setup was extended in [10] to several similar ensembles. The key analytic properties of continued fractions—their convergence—can be linked to asymptotic behaviour of such an infinite ensemble [2].

(4) An important example of an infinite ensemble is provided by the representation of an arbitrary wave as the envelope of a continuous family of spherical waves. A finite subset of spheres can be used as an approximation to the infinite family. Then, discrete snapshots of time evolution

of sphere wave packets represent a FLT-covariant ensemble of cycles. This and further physical applications of FLT-invariant ensembles may be looked at [5].

Definition 2. The extend Möbius–Lie geometry considers ensembles of cycles interconnected through FLT-invariant relations.

Naturally, “old” objects—cycles—are represented by simplest one-element ensembles without any relation. The paper [9] provides conceptual foundations of such extension and demonstrates its practical implementation as a CPP library **figure**¹. Interestingly, the development of this library shaped the general approach, which leads to specific realisations in [11, 10, 12].

More specifically, the library **figure** manipulates ensembles of cycles (quadrics) interrelated by certain FLT-invariant geometric conditions. The code is build on top of the previous library **cycle** [7, 8, 6], which manipulates individual cycles within the GiNaC [1] computer algebra system. Thinking an ensemble as a graph, one can say that the library **cycle** deals with individual vertices (cycles), while **figure** considers edges (relations between pairs of cycles) and the whole graph. Intuitively, an interaction with the library **figure** reminds compass-and-straightedge constructions, where new lines or circles are added to a drawing one-by-one through relations to already presented objects (the line through two points, the intersection point or the circle with given centre and a point).

It is important that both libraries are capable to work in spaces of any dimensionality and metrics with an arbitrary signatures: Euclidean, Minkowski and even degenerate. Parameters of objects can be symbolic or numeric, the latter admit calculations with exact or approximate arithmetic. Drawing routines work with any (elliptic, parabolic or hyperbolic) metric in two dimensions and the euclidean metric in three dimensions.

REFERENCES

- [1] Christian Bauer, Alexander Frink, and Richard Kreckel. Introduction to the GiNaC framework for symbolic computation within the C++ programming language. *J. Symbolic Computation*, 33(1):1–12, 2002. Web: <http://www.ginac.de/>. [arXiv:cs/0004015](https://arxiv.org/abs/cs/0004015).
- [2] Alan F. Beardon and Ian Short. A geometric representation of continued fractions. *Amer. Math. Monthly*, 121(5):391–402, 2014.
- [3] Walter Benz. *Classical Geometries in Modern Contexts. Geometry of Real Inner Product Spaces*. Birkhäuser Verlag, Basel, second edition edition, 2007.
- [4] GNU. *General Public License (GPL)*. Free Software Foundation, Inc., Boston, USA, version 3 edition, 2007. URL: <http://www.gnu.org/licenses/gpl.html>.
- [5] H.A. Kastrup. On the advancements of conformal transformations and their associated symmetries in geometry and theoretical physics. *Annalen der Physik*, 17(9–10):631–690, 2008. [arXiv:0808.2730](https://arxiv.org/abs/0808.2730).
- [6] Vladimir V. Kisil. Erlangen program at large-0: Starting with the group $SL_2(\mathbf{R})$. *Notices Amer. Math. Soc.*, 54(11):1458–1465, 2007. [arXiv:math/0607387](https://arxiv.org/abs/math/0607387), [On-line. Zbl1137.22006](https://zbmath.org/journal/Zbl1137.22006).
- [7] Vladimir V. Kisil. Fillmore-Springer-Cnops construction implemented in GiNaC. *Adv. Appl. Clifford Algebr.*, 17(1):59–70, 2007. [On-line. A more recent version: arXiv:cs/0512073](https://arxiv.org/abs/cs/0512073). The latest documentation, source files, and live ISO image are at the project page: <http://moebinv.sourceforge.net/>. [Zbl05134765](https://zbmath.org/journal/Zbl05134765).
- [8] Vladimir V. Kisil. *Geometry of Möbius Transformations: Elliptic, Parabolic and Hyperbolic Actions of $SL_2(\mathbf{R})$* . Imperial College Press, London, 2012. Includes a live DVD. [Zbl1254.30001](https://zbmath.org/journal/Zbl1254.30001).
- [9] Vladimir V. Kisil. An extension of Lie spheres geometry with conformal ensembles of cycles and its implementation in a GiNaC library. 2014–2018. [arXiv:1512.02960](https://arxiv.org/abs/1512.02960). Project page: <http://moebinv.sourceforge.net/>.
- [10] Vladimir V. Kisil. Remark on continued fractions, Möbius transformations and cycles. *Izvestiya Komi nauchnogo centra UrO RAN [Izvestiya Komi nauchnogo centra UrO RAN]*, 25(1):11–17, 2016. [arXiv:1412.1457](https://arxiv.org/abs/1412.1457), [on-line](https://arxiv.org/abs/1412.1457).
- [11] Vladimir V. Kisil. Poincaré extension of Möbius transformations. *Complex Variables and Elliptic Equations*, 62(9):1221–1236, 2017. [arXiv:1507.02257](https://arxiv.org/abs/1507.02257).
- [12] Vladimir V. Kisil and James Reid. Conformal parametrisation of loxodromes by triples of circles. 2018. [arXiv:1802.01864](https://arxiv.org/abs/1802.01864).

¹All described software is licensed under GNU GPLv3 [4].

Зміст

| | |
|--|-----------|
| N. Aygor, H. Burhanzade <i>Secondary school students' misconceptions about linear algebra</i> | 3 |
| S. Bardyla, H. Kvasnytsia <i>Semitopological graph inverse semigroups</i> | 4 |
| B. A. Bhayo <i>On inequalities of generalized elliptic integrals</i> | 5 |
| Bodzioch M., Choiński M., Foryś U. <i>A criss-cross model of tuberculosis for heterogenous population</i> | 6 |
| Bolotov D. V. <i>Foliations with leaves of non-positive curvature and bounded total curvature on closed 3-manifolds</i> | 7 |
| E. Bonacci <i>Algebraic and geometric questions about a 6D physics</i> | 9 |
| F. Bulnes <i>Mukai-Fourier Transform in Derived Categories to Solutions of the Field Equations: Gravitational Waves as Oscillations in the Space-Time Curvature/Spin IV</i> | 10 |
| H. Burhanzade, N. Aygor <i>A study on the teaching methods in determinants</i> | 12 |
| Damla Yaman <i>Order continuity properties of lattice ordered algebras</i> | 13 |
| Denega I. <i>Problem on non-overlapping polycylindrical domains with poles on the boundary of a polydisk</i> | 14 |
| A. Dudko, V. Pivovarchik <i>Inverse three spectra problem for a Stieltjes string with the Neumann boundary conditions</i> | 16 |
| Eftekharinasab K. <i>On the existence of a global diffeomorphism between Fréchet spaces</i> | 18 |
| Glazunov N. <i>Class groups of rings with divisor theory, L-functions and moduli spaces</i> | 19 |
| O. Gok <i>b-bimorphisms</i> | 21 |
| Gül E. <i>On the second regularized trace formula for a differential operator with unbounded coefficients</i> | 22 |
| Hentosh O. Ye., Prykaratsky Ya. A. <i>The Lie-algebraic structure of the Lax-Sato integrable superanalogs for the Liouville heavenly type equations</i> | 24 |
| V. Herasymov <i>In a natural topological sense a typical linear nonhomogeneous differential equation in the ring $Z[[x]]$ has no solutions from $Z[[x]]$.</i> | 26 |
| Juraev D. A. <i>On the Cauchy problem for matrix factorizations of the Helmholtz equation</i> | 27 |
| M. E. Kansu <i>Macroscopic electromagnetism via complex quaternions</i> | 29 |
| Vladimir V. Kisil <i>An extension of Möbius–Lie geometry with conformal ensembles of cycles</i> | 30 |
| Konovenko N., Lychagin V. <i>Rational differential invariants for oriented primary visual cortex</i> | 32 |