

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

On the relationship between the Smith normal forms of matrices and of their least common multiple

Romaniv A. M.

(Pidstryhach Institute for Applied Problems of Mechanics and Mathematics NAS of Ukraine)

E-mail: romaniv_a@ukr.net

Let R be a commutative principal ideal domain with $1 \neq 0$, $M_n(R)$ be a ring $n \times n$ matrices over R and let $A, B \in M_n(R)$.

If $A = BC$, then we will say that B is a left divisor of matrix A and A is a right multiple of B . Moreover, if $M = AA_1 = BB_1$ then the matrix M is called a common right multiple of matrices A and B . If in addition the matrix M is a left divisor of any other common right multiple of matrices A and B then we say that M is a **least common right multiple** of A and B . ($[A, B]_r$ in notation).

In the 80's of the last century, M. Newman formulated the problem to establish of the relationship between the Smith normal forms of matrices and of their least common multiple and greatest common divisor over commutative principal ideal domain. The most important results in the study of M. Newman's problem are obtained in the papers [1], [2, 3] and [4].

Let A be non-singular $n \times n$ matrix over R . Then there exist invertible matrices P_A, Q_A , such that

$$P_A A Q_A = E = \text{diag}(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n), \text{ where } \varepsilon_i | \varepsilon_{i+1}, i = 1, \dots, n-1.$$

The matrix E is called the Smith normal form, P_A and Q_A are called left and right transforming matrices for matrix A . Denote by \mathbf{P}_A the set of all left transforming matrices for matrix A . Let the Smith normal form of the matrix B is

$$B \sim \Delta = \text{diag}(\delta_1, \delta_2, \dots, \delta_n), \text{ where } \delta_i | \delta_{i+1}, i = 1, \dots, n-1.$$

Consider the set of matrices

$$\mathbf{L}(E, \Delta) = \{L \in \text{GL}_n(R) \mid \exists L_1 \in M_n(R) : LE = \Delta L_1\}.$$

Theorem 1. *Let R be a commutative principal ideal domain and let*

$$A \sim \text{diag}(1, \varepsilon, \dots, \varepsilon), B \sim \text{diag}(1, \dots, 1, \delta),$$

$P_B P_A^{-1} = \|s_{ij}\|_1^n$, $P_B \in \mathbf{P}_B$, $P_A \in \mathbf{P}_A$. Then

$$[A, B]_r = (L_A P_A)^{-1} \Omega = (L_B P_B)^{-1} \Omega,$$

where

$$\Omega = \text{diag}\left(\frac{(\varepsilon, \delta)}{((\varepsilon, \delta), s_{n1})}, \varepsilon, \dots, \varepsilon, [\varepsilon, \delta]\right),$$

L_A, L_B belong to sets $\mathbf{L}(\Omega, E)$, $\mathbf{L}(\Omega, \Delta)$ respectively and satisfy the equality:

$$(P_B P_A^{-1}) L_A = L_B.$$

REFERENCES

- [1] N. Erawaty, M. Bahri, L. Haraynto, A. Amir Greatest common divisors of Euclidean domain matrices. *Far East Journal of Math. Sci.*, 101(4) : 855–872, 2017.
- [2] A. M. Romaniv, V. P. Shchedryk Greatest common divisor of matrices one of which is a disappear matrix. *Ukrainian Mathematical Journal*, 66(3) : 425–430, 2014.
- [3] A. M. Romaniv, V. P. Shchedryk The greatest common left divisor and the least common right multiple of second order matrices. *Math. Bull. Shevchenko Sci. Soc.*, (9) : 269–284, 2012.
- [4] R. C. Thompson Left multiples and right divisors of integral matrices. *Linear and Multilinear Algebra*, 19(3) : 287–295, 1986.

V. Lomadze <i>A purely algebraic construction of Schwartz distributions</i>	34
S. Maksymenko <i>Deformation of functions on orientable surfaces by symplectic diffeomorphisms</i>	36
Martseniuk V., Bondarenko A., Gefter S. <i>Application the p-adic topology on Z for study determinants of infinite order with integer coefficients</i>	37
K. Matsumoto <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds, II</i>	38
Mormul P. <i>What points in Goursat Monster Tower are strongly nilpotent</i>	39
Mukhamadiev F. G. <i>The local density and the local weak density of N_7^φ-kernel of a topological spaces</i>	40
Tetiana V. Obikhod <i>Topological invariants and microscopic quantum description</i>	41
A. Ozkan <i>On the misconceptions of 10th grade students about analytical geometry</i>	42
Parasyuk I. O. <i>Hyperbolic quasiperiodic motion of charged particle on 2-sphere</i>	43
E. Petrov <i>Representing trees of finite ultrametric spaces and weak similarities</i>	45
Prishlyak A., Hladysh B. <i>Functions with three critical points on closed non-oriented 3-manifolds</i>	47
V.M. Prokip <i>A note on similarity of matrices</i>	49
Romakina L.N. <i>Inversion with respect to an elliptic cycle of a hyperbolic plane of positive curvature</i>	51
Romaniv A. M. <i>On the relationship between the Smith normal forms of matrices and of their least common multiple</i>	53
Ryazanova O. <i>Estimates for the surface with given average curvature</i>	54
Samokhvalov S. E. <i>Kinematic renormalization of energy in the gravity</i>	55
Savchenko A. <i>Fuzzy metrization of the spaces of idempotent measures</i>	56
Sazonova O. <i>General form of the Maxwellian distribution with arbitrary density</i>	57
Skuratovskii R. <i>The commutator and centralizer of Sylow subgroups of alternating and symmetric groups, its minimal generating set</i>	58
Sokhatsky F., Krainichuk H. <i>Semi-lattice of varieties of quasigroups with linearity</i>	59
Salimov R., Stefanchuk M. <i>Asymptotic behavior of solutions to a nonlinear Beltrami equation</i>	61
Mokritskaya T. P., Tushev A. V. <i>On fractal characteristics of the micro-aggregate structure of loess soils</i>	62
Vasilchenko A. N. <i>Spaces of primitive elements in dual modules over Steenrod algebra</i>	64