

International scientific conference

**“Algebraic and Geometric
Methods of Analysis”**

Book of abstracts



May 28 - June 3, 2019

Odesa, Ukraine

Conference webpage: imath.kiev.ua/~topology/conf/agma2019/

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences
- History and methodology of teaching in mathematics

ORGANIZERS

- The Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- The Institute of Mathematics of the National Academy of Sciences of Ukraine
- Odessa I. I. Mechnikov National University
- Taras Shevchenko National University of Kyiv
- The International Geometry Center

PROGRAM COMMITTEE

Chairman: Prishlyak A. (<i>Kyiv, Ukraine</i>)	Konovenko N. (<i>Odesa, Ukraine</i>)	Pokas S. (<i>Odesa, Ukraine</i>)
Balan V. (<i>Bucharest, Romania</i>)	Lyubashenko V. (<i>Kyiv, Ukraine</i>)	Polulyakh E. (<i>Kyiv, Ukraine</i>)
Banakh T. (<i>Lviv, Ukraine</i>)	Maksymenko S. (<i>Kyiv, Ukraine</i>)	Sabitov I. (<i>Moscow, Russia</i>)
Fedchenko Yu. (<i>Odesa, Ukraine</i>)	Matsumoto K. (<i>Yamagata, Japan</i>)	Savchenko A. (<i>Kherson, Ukraine</i>)
Fomenko A. (<i>Moscow, Russia</i>)	Mikesh J. (<i>Olomouc, Czech Republic</i>)	Sergeeva A. (<i>Odesa, Ukraine</i>)
Fomenko V. (<i>Taganrog, Russia</i>)	Mormul P. (<i>Warsaw, Poland</i>)	Shvets V. (<i>Odesa, Ukraine</i>)
Haddad M. (<i>Wadi al-Nasara, Syria</i>)	Moskaliuk S. (<i>Wien, Austria</i>)	Shelekhov A. (<i>Tver, Russia</i>)
Karlova O. (<i>Chernivtsi, Ukraine</i>)	Mykhailyuk V. (<i>Chernivtsi, Ukraine</i>)	Vlasenko I. (<i>Kyiv, Ukraine</i>)
Kiosak V. (<i>Odessa, Ukraine</i>)	Nykyforchyn O. (<i>Ivano-Frankivsk, Ukraine</i>)	Volkov V. (<i>Odessa, Ukraine</i>)
Kirillov V. (<i>Odesa, Ukraine</i>)	Plachta L. (<i>Krakov, Poland</i>)	Zadorozhnyj V. (<i>Odesa, Ukraine</i>)
		Zarichnyi M. (<i>Lviv, Ukraine</i>)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Svytyy I., Dean of the Faculty of Computer Systems and Automation.

ORGANIZING COMMITTEE

Kirillov V.
Konovenko N.
Fedchenko Yu.

Prus A.
Osadchuk E.

Maksymenko S.
Khudenko N.
Cherevko E.

ФІТБ ОНАФТ

Around the homologous sphere of Poincare and its applications

V.S.Dryuma

(IMI «Vladimir Andrunachievici», Moldova, Kishinev)

E-mail: valdryum@gmail.com

Topology of a 3-dim manifolds defined by the system of equations

$$|z_1|^2 + |z_2|^2 + |z_3|^2 - 1 = 0, \quad z_1^l + z_2^m + z_3^n = 0, \quad (1)$$

where $z_k = x_k + iy_k$ are the complex coordinates, depends from the values of the parameters l, m, n . In the case $l = 2, m = 3, n = 5$ the manifold defined by the conditions (10) is a famous homologous sphere of Poincare, which has a set of homologies same with standard 3D-sphere $|z_1|^2 + |z_2|^2 = 1$, but differs from it by self fundamental group. It has an important applications in various branch of modern algebraic topology (J.Milnor,1968).

In this report will be told how to represent the homologous sphere defined by intersection of the five-dimensional sphere with singular manifold ($l = 2, m = 3, n = 5$)

$$|z_1|^2 + |z_2|^2 + |z_3|^2 = 1, \quad z_1^2 + z_2^3 + z_3^5 = 0. \quad (2)$$

in the form of an explicit expression for one function between of the four variables $H(x, y, u, v) = 0$.

Theorem 1. *In the Eulerian coordinates*

$$\begin{aligned} z_1 &= \cos(\theta) e^{-2/3 i \sqrt{3} \phi}, \quad z_2 = -\sin(\theta) \sin(1/2 \beta) e^{-1/2 i (\alpha - \delta + 4/3 \sqrt{3} \phi)}, \\ z_3 &= \sin(\theta) \cos(1/2 \beta) e^{1/2 i (\alpha + \delta - 4/3 \sqrt{3} \phi)}, \end{aligned} \quad (3)$$

the equation of the unit five-dimensional sphere is identically satisfied and the equation of the orbifold $z_1^2 + z_2^3 + z_3^5 = 0$ takes the form

$$\begin{aligned} &(\cos(\theta))^2 e^{-4/3 i \sqrt{3} \phi} - \sin(\theta) \sin(1/2 \beta) e^{-1/2 i \sqrt{3} (\sqrt{3} \alpha - \sqrt{3} \delta + 4 \phi)} + \\ &+ \sin(\theta) \sin(1/2 \beta) e^{-1/2 i \sqrt{3} (\sqrt{3} \alpha - \sqrt{3} \delta + 4 \phi)} (\cos(1/2 \beta))^2 + \\ &+ \sin(\theta) \sin(1/2 \beta) e^{-1/2 i \sqrt{3} (\sqrt{3} \alpha - \sqrt{3} \delta + 4 \phi)} (\cos(\theta))^2 - \\ &- \sin(\theta) \sin(1/2 \beta) e^{-1/2 i \sqrt{3} (\sqrt{3} \alpha - \sqrt{3} \delta + 4 \phi)} (\cos(\theta))^2 (\cos(1/2 \beta))^2 + \\ &+ \sin(\theta) (\cos(1/2 \beta))^5 e^{5/6 i \sqrt{3} (\sqrt{3} \alpha + \sqrt{3} \delta - 4 \phi)} - \\ &- 2 \sin(\theta) (\cos(1/2 \beta))^5 e^{5/6 i \sqrt{3} (\sqrt{3} \alpha + \sqrt{3} \delta - 4 \phi)} (\cos(\theta))^2 + \\ &+ \sin(\theta) (\cos(1/2 \beta))^5 e^{5/6 i \sqrt{3} (\sqrt{3} \alpha + \sqrt{3} \delta - 4 \phi)} (\cos(\theta))^4 = 0. \end{aligned} \quad (4)$$

Using then the variable χ , defined by the condition $e^{5/2 i \alpha + 5/2 i \delta - 10/3 i \sqrt{3} \phi} - e^{5 \chi} = 0$, we express the variable ϕ as $\phi = -1/4 i (i \alpha + i \delta - 2 \chi) \sqrt{3}$ and after separation of the real and imaginary parts of complex equation ((4)), are obtained two equations into the five variables α, δ, χ and θ, β .

As result of elimination of the variable χ from both equations is derived equation of homologous sphere of Poincare in the form of one function of the four variables. The equation is the summa of the functions $\sin()$ and $\cos()$ with linear arguments. It contains more than 200 items.

By analogy can be considered the case of tetrahedral space which corresponds to the intersection of the five-dimensional sphere with singular manifold ($l = 2, m = 3, n = 4$)

$$|z_1|^2 + |z_2|^2 + |z_3|^2 = 1, \quad z_1^2 + z_2^3 + z_3^4 = 0. \quad (5)$$

and the octahedral space defined by the condition

$$|z_1|^2 + |z_2|^2 + |z_3|^2 = 1, \quad z_1^2 + z_2^3 + z_2 z_3^3 = 0. \quad (6)$$

Proposition 2. *The relation between a four variables $F(x, y, a, b) = 0$ which defines some 3D-variety can be considered as General Integral of the pair of the second order ODE's $f(x, y, y', y'') = 0$ and $g(a, b, b', b'') = 0$.*

$$\sphericalangle F(x, y, a, b) = 0 \searrow$$

$$y'' = f(x, y, y') \quad \longleftrightarrow \quad b'' = h(a, b, b')$$

The Liouville-Tresse invariants of both equations with respect to non degenerate transformations of the variables $x = X(u, v), y = Y(u, v)$ or $a = A(p, q), b = B(p, q)$ can be used for the studies of topological properties of the manifold $F(x, y, a, b) = 0$.

Theorem 3. *Spatial homogeneous the first order system of the equations*

$$\begin{aligned} \frac{d}{ds}x(s) &= 4 a_0 z^2 + (4 a_2 y + (3 a_1 - b_2) x) z + 4 a_{22} y^2 + \\ &\quad + (3 a_{12} - 2 b_{22}) xy + (2 a_{11} - b_{12}) x^2, \\ \frac{d}{ds}y(s) &= 4 b_0 z^2 + ((3 b_2 - a_1) y + 4 b_1 x) z + (2 b_{22} - a_{12}) y^2 + \\ &\quad + (-2 a_{11} + 3 b_{12}) xy + 4 b_{11} x^2, \\ \frac{d}{ds}z(s) &= (-b_2 - a_1) z^2 + ((-2 b_{22} - a_{12}) y - b_{12} x - 2 a_{11} x) z, \end{aligned} \quad (7)$$

is projective extension of planar polynomial differential systems

$$\frac{dx}{ds} = a_0 + a_1 x + a_2 y + a_{11} x^2 + a_{12} xy + a_{22} y^2, \quad \frac{dy}{ds} = b_0 + b_1 x + b_2 y + b_{11} x^2 + b_{12} xy + b_{22} y^2 \quad (8)$$

with the parameters a_i, a_{ij} and b_i, b_{ij} .

After eliminating of the variables $(y(x))$ or $z(x)$ it is reduced to the second-order differential equations $F(x, y, y', y'') = 0$.

As a result of the exclusion of the function y'' from the system

$$F(x, y, y', y'') = 0, \quad \frac{\partial F}{\partial y''} = 0,$$

a first order differential equation is arised $C(x, y, y') = 0$. Through each point M of integral curve $C = Q(x, y)$ passes the integral curve of the equation $F(x, y, y', y'') = 0$, for which the point M is the return point of the second type and this allow to study the limit cycles of the system (6).

REFERENCES

- [1] John Milnor, Singular points of complex hypersurfaces, Annals of Mathematics Studies, No. 61, Princeton University Press, Princeton, N.J., 1968.
- [2] V. Dryuma. On dual equations in theory of the second order ODE's, arXiv:nlin/0701047 v1 p.1-17, 2007.
- [3] V. Dryuma. Homogenous extensions of the first order ODE's, Algebraic Topology and Abelian Functions, Buchstaber'70 Conference, 18-22 June 2013, Steklov's MI RAS, Moscow, ABSTRACTS, p.78-79, 2013.
- [4] V. S. Dryuma. On the equation of homologous sphere of Poincare. International conference "Classical and modern geometry", (Moscow, April 22-25, 2019), ed. A.V. Tsarev MGPU, 20-21, 2019.

Зміст

Absamatov Z.A. <i>Formation of algorithmic culture of students in the classroom of higher mathematics</i>	3
Absamatov Z. A., Khamrayev A. Yu. <i>Behavior of the trajectories of a single cubic operator</i>	4
Banaru G. A. <i>On nonexistence of Kenmotsu structure on Kirichenko–Uskorev-hypersurfaces of Kählerian manifolds</i>	5
Banaru M. B. <i>On almost contact metric hypersurfaces in W_4-manifolds</i>	7
Batkhin A. B. <i>Quantum calculus and singularities of quasi-discriminant sets</i>	9
Bernatska J. <i>Derivative Thomae formula for singular half-periods</i>	11
Bilet V., Dovgoshey O. <i>Kuratowski limits of subsets of real line and their applications to pretangent spaces</i>	13
Bonacci E. <i>Algebraic and geometric questions about a FTL physics</i>	15
Bruno A. D. <i>Algorithms for solving an algebraic equation</i>	16
Dryuma V. S. <i>Around the homologous sphere of Poincare and its applications</i>	17
Eftekharinasab K. <i>On the generalization of the Darboux theorems</i>	19
Favorov S. <i>Discrete sets, discrete measures, quasicrystals Fourier, pure crystals</i>	20
Glazunov N. <i>Algebraic-geometric aspects of function field analogues to abelian varieties</i>	21
Gok O. <i>Extensions of almost orthosymmetric lattice bimorphisms</i>	23
Grechneva M., Stegantseva P. <i>The properties of the surface of Minkowski space, which determine the type of its Grassmann image</i>	24
Gutik O., Melnyk K. <i>The semigroup of star partial homeomorphisms of a finite deminsional Euclidean space</i>	25
Gutik O., Sobol O. <i>Extensions of semigroups by symmetric inverse semigroups of a bounded finite rank</i>	26
Prishlyak A., Hatamian H. <i>Non-Oriented Heegaard Diagrams</i>	28
Herasymov V., Gefter S., Arinenkov A. <i>Some many-dimensional extremal geometric problems</i>	30
Juraev D. A. <i>On a regularized solution of the Cauchy problem for matrix factorizations of the Helmholtz equation in m-dimensional bounded domain</i>	31
Kozerenko S. <i>Neighborhood maps on combinatorial trees and their Markov graphs</i>	33
Kuznietsova I., Soroka Yu. <i>First Betti numbers of orbits of Morse functions on surfaces</i>	34
Maksymenko S., Khohliyk O. <i>Diffeomorphisms preserving Morse-Bott foliations</i>	35
Markitan V. <i>Singular monotonic functions defined by a convergent positive series and a double stochastic matrix</i>	36
Matsumoto K. <i>A Flat $(CHR)_3$-curvature tensor in a Trans-Sasakian Manifold</i>	38