

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

Fuzzy metrization of the spaces of idempotent measures

Aleksandr Savchenko

(Kherson State Agrarian University)

E-mail: savchenko.o.g@ukr.net

Let X be a compact Hausdorff space. As usual, by $C(X)$ we denote the space of continuous functions on X endowed with the sup-norm. For any $c \in \mathbb{R}$, we denote by c_X the constant function on X taking the value c .

Let $\mathbb{R}_{\max} = \mathbb{R} \cup \{-\infty\}$. We consider the natural order on $\mathbb{R}_{\max}$. We also use the following traditional notation from idempotent mathematics: $\oplus$ for max and $\odot$ instead of $+$ (this may concern either numbers or functions).

A functional $\mu: C(X) \rightarrow \mathbb{R}$ is said to be an *idempotent measure* (Maslov measure) if the following holds: (1) $\mu(c_X) = c$; (2) $\mu(\varphi \oplus \psi) = \mu(\varphi) \oplus \mu(\psi)$; (3) $\mu(\lambda \odot \varphi) = (\lambda \odot \mu)\varphi$. Remark that the notion of idempotent measure is a counterpart of the notion of probability measure in the so called idempotent mathematics.

The set $I(X)$ is endowed with the weak* topology (see [1]). Actually, the construction I determines a functor in the category of compact Hausdorff spaces. The space $I(X)$ is known to be metrizable for metrizable X .

The aim of the talk is to provide a fuzzy metrization of the space $I(X)$ for fuzzy metrizable X . We use the notion of fuzzy metric in the sense of George and Veeramani [2]. Recall that a triple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm and M is a fuzzy set on $X \times X \rightarrow (0, \infty)$ satisfying the following conditions for all $x, y, z \in X$ and $s, t \in (0, \infty)$:

- (1) $M(x, y, t) > 0$;
- (2) $M(x, y, t) = 1$ if and only if $x = y$;
- (3) $M(x, y, t) = M(y, x, t)$;
- (4) $M(x, y, t) * M(y, z, s) = M(x, z, t + s)$;
- (5) the function $M(x, y, \cdot): (0, \infty) \rightarrow [0, 1]$ is continuous.

Our approach is based on the notion of density of idempotent measure. Also, we use the fuzzy Hausdorff metric on the hyperspaces of fuzzy metric spaces [3]. We also show that the obtained construction determines a functor in the category of compact fuzzy metric spaces and non-expanding maps. We establish some properties of this functor related to the notion of monad.

REFERENCES

- [1] Mikhail M. Zarichnyi. Spaces and maps of idempotent measures. *Izvestiya: Mathematics*, 74(3) : 45–64, 2010.
- [2] A. George and P. Veeramani. On some result in fuzzy metric space. *Fuzzy Sets and System*, 64 : 395–399, 1994.
- [3] J. Rodríguez-López, S. Romaguera. The Hausdorff fuzzy metric on compact sets. *Fuzzy Sets and Systems*, 147 : 273–283, 2004.

V. Lomadze <i>A purely algebraic construction of Schwartz distributions</i>	34
S. Maksymenko <i>Deformation of functions on orientable surfaces by symplectic diffeomorphisms</i>	36
Martseniuk V., Bondarenko A., Gefter S. <i>Application the p-adic topology on Z for study determinants of infinite order with integer coefficients</i>	37
K. Matsumoto <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds, II</i>	38
Mormul P. <i>What points in Goursat Monster Tower are strongly nilpotent</i>	39
Mukhamadiev F. G. <i>The local density and the local weak density of N_T^φ-kernel of a topological spaces</i>	40
Tetiana V. Obikhod <i>Topological invariants and microscopic quantum description</i>	41
A. Ozkan <i>On the misconceptions of 10th grade students about analytical geometry</i>	42
Parasyuk I. O. <i>Hyperbolic quasiperiodic motion of charged particle on 2-sphere</i>	43
E. Petrov <i>Representing trees of finite ultrametric spaces and weak similarities</i>	45
Prishlyak A., Hladysh B. <i>Functions with three critical points on closed non-oriented 3-manifolds</i>	47
V.M. Prokip <i>A note on similarity of matrices</i>	49
Romakina L.N. <i>Inversion with respect to an elliptic cycle of a hyperbolic plane of positive curvature</i>	51
Romaniv A. M. <i>On the relationship between the Smith normal forms of matrices and of their least common multiple</i>	53
Ryazanova O. <i>Estimates for the surface with given average curvature</i>	54
Samokhvalov S. E. <i>Kinematic renormalization of energy in the gravity</i>	55
Savchenko A. <i>Fuzzy metrization of the spaces of idempotent measures</i>	56
Sazonova O. <i>General form of the Maxwellian distribution with arbitrary density</i>	57
Skuratovskii R. <i>The commutator and centralizer of Sylow subgroups of alternating and symmetric groups, its minimal generating set</i>	58
Sokhatsky F., Krainichuk H. <i>Semi-lattice of varieties of quasigroups with linearity</i>	59
Salimov R., Stefanchuk M. <i>Asymptotic behavior of solutions to a nonlinear Beltrami equation</i>	61
Mokritskaya T. P., Tushev A. V. <i>On fractal characteristics of the micro-aggregate structure of loess soils</i>	62
Vasilchenko A. N. <i>Spaces of primitive elements in dual modules over Steenrod algebra</i>	64