

International  
Scientific Conference



Algebraic  
and Geometric  
Methods  
of Analysis

27-30 May 2024  
Odesa, Ukraine

The purpose of this conference is to bring together researchers in geometry, topology, algebra, analysis and dynamical systems and to provide for them a forum to present their recent work to colleagues from different nationalities. This way we aim to stimulate discussion about the latest findings in geometrical and topological methods in analysis and to increase international collaboration.

The conference continues the traditional annual conference «Geometry in Odesa» holding from 2004, and hosted by Odesa National University of Technology (Odesa National Academy of Food Technologies till 2021). From 2017 the conference was renamed to «Algebraic and geometric methods of analysis» (AGMA).

The Conference languages: Ukrainian and English.

#### LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

#### ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology, Ukraine
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

#### SCIENTIFIC COMMITTEE

- |   |  |
|---|--|
| • <b>Vladimir Balan</b> ( <i>Bucharest, Romania</i> ) | • <b>Volodymyr Lyubashenko</b> ( <i>Kyiv, Ukraine</i> )  |
| • <b>Taras Banakh</b> ( <i>Lviv, Ukraine</i> )        | • <b>Sergiy Maksymenko</b> ( <i>Kyiv, Ukraine</i> )      |
| • <b>Dmytro Bolotov</b> ( <i>Kharkiv, Ukraine</i> )   | • <b>Koji Matsumoto</b> ( <i>Yamagata, Japan</i> )       |
| • <b>Vyacheslav Boyko</b> ( <i>Kyiv, Ukraine</i> )    | • <b>Piotr Mormul</b> ( <i>Warsaw, Poland</i> )          |
| • <b>Yulia Fedchenko</b> ( <i>Odesa, Ukraine</i> )    | • <b>Maryna Nesterenko</b> ( <i>Kyiv, Ukraine</i> )      |
| • <b>Oleg Gutik</b> ( <i>Lviv, Ukraine</i> )          | • <b>Roman Popovych</b> ( <i>Kyiv, Ukraine</i> )         |
| • <b>Olena Karlova</b> ( <i>Chernivtsi, Ukraine</i> ) | • <b>Alexandr Prishlyak</b> ( <i>Kyiv, Ukraine</i> )     |
| • <b>Volodymyr Kiosak</b> ( <i>Odesa, Ukraine</i> )   | • <b>Aleksandr Savchenko</b> ( <i>Kherson, Ukraine</i> ) |
| • <b>Nadiia Konovenko</b> ( <i>Odesa, Ukraine</i> )   |  |

#### ORGANIZING COMMITTEE

- |   |  |
|---|--|
| • <b>Nadiia Konovenko</b> ( <i>Odesa, Ukraine</i> ) | • <b>Bohdan Mazhar</b> ( <i>Kyiv, Ukraine</i> )      |
| • <b>Yuliya Fedchenko</b> ( <i>Odesa, Ukraine</i> ) | • <b>Sergiy Maksymenko</b> ( <i>Kyiv, Ukraine</i> )  |
| • <b>Mykola Lysynskiy</b> ( <i>Kyiv, Ukraine</i> )  | • <b>Alexandr Prishlyak</b> ( <i>Kyiv, Ukraine</i> ) |

# Rational factorization of Lax type flows in the space dual to the centrally extended Lie algebra of matrix super-integro-differential operators

O.Ye. Hentosh

(Pidstryhach Inst. for Applied Problems of Mech. and Math., NASU, Lviv, Ukraine)

*E-mail:* ohen@ukr.net

In papers of A. Prykarpatski, O. Hentosh and their coauthors the Lie-algebraic approach to the rational factorization of Lax type flows in spaces dual to certain operator Lie algebras and central extensions of some of them is developed for the central extension of the Lie algebra [1] of matrix super-integro-differential operators with one anticommuting variable.

Let us consider the Lie algebra  $\mathfrak{g}$  which consists of matrix super-integro-differential operators such as  $\mathcal{A} := \mathbf{1}\partial^q + \sum_{p < 2q} A_p D_\theta^p$ , where  $A_r \in C^\infty(\mathbb{S} \times \Lambda_1; gl(m|n))$ ,  $gl(m|n)$  is a semi-simple Lie superalgebra of square supermatrices,  $A_p = A_p(x, \theta) := A_p^0(x) + \theta A_p^1(x)$ ,  $p \in \mathbb{Z}$ , the supermatrices  $A_p$  are even for every even  $p$  and odd for every odd  $p$ ,  $\mathbf{1} \in gl(m+n)$  is a unit matrix,  $q \in \mathbb{N}$ ,  $\partial = \partial/\partial x$ ,  $x \in \mathbb{S} \simeq \mathbb{R}/2\pi\mathbb{Z}$ ,  $\theta \in \Lambda_1$ ,  $\Lambda := \Lambda_0 \oplus \Lambda_1$  is a Grassmann algebra over the field  $\mathbb{C} \subset \Lambda_0$ ,  $D_\theta := \partial/\partial\theta + \theta\partial/\partial x$  is a superderivative, for which  $D_\theta^2 = \partial/\partial x$ , and  $\partial/\partial\theta$  is a left partial derivative by the anticommuting variable  $\theta$ , with the standard commutator  $[\cdot, \cdot]$ , acting by the rule:

$$[\mathcal{A}, \mathcal{B}] = \mathcal{A} \circ \mathcal{B} - \mathcal{B} \circ \mathcal{A}, \quad \mathcal{A}, \mathcal{B} \in \mathfrak{g}, \quad (1)$$

where symbol "o" denotes the product of two matrix super-integro-differential operators (see [1]). The scalar product  $(\mathcal{A}, \mathcal{B}) = \int_{x \in \mathbb{S}} dx \int d\theta \text{sSp res}_{D_\theta}(\mathcal{A}\mathcal{B})$ , where "res $_{D_\theta}$ " denotes the coefficient at  $D_\theta^{-1}$  in the expansion of a matrix super-integro-differential operator and "sSp" is a supermatrix supertrace, being invariant with respect to the commutator (1), allows us to identify the dual space  $\mathfrak{g}^*$  to  $\mathfrak{g}$ , with the Lie algebra itself. The latter is splitting into the direct sum  $\mathfrak{g} := \mathfrak{g}_+ \oplus \mathfrak{g}_-$  of two its Lie subalgebras, where  $\mathfrak{g}_+$  is a Lie subalgebra of formal polynomials by the superderivative operator with supermatrix-valued coefficients, and  $\mathfrak{g}_+^* \simeq \mathfrak{g}_-$ ,  $\mathfrak{g}_-^* \simeq \mathfrak{g}_+$ .

One constructs the central extension  $\hat{\mathfrak{g}} := \tilde{\mathfrak{g}} \oplus \mathbb{C}$  of parameterized Lie algebra  $\tilde{\mathfrak{g}} := \prod_{y \in \mathbb{S}} \mathfrak{g}$  by the Maurer-Cartan two-cocycle on  $\tilde{\mathfrak{g}}$  such as  $\omega_2(\mathcal{A}, \mathcal{B}) = \int_{y \in \mathbb{S}} dy (\mathcal{A}, \partial\mathcal{B}/\partial y)$ , where  $\mathcal{A}, \mathcal{B} \in \tilde{\mathfrak{g}}$ , with the commutator

$$[(\mathcal{A}, d), (\mathcal{B}, e)] = ([\mathcal{A}, \mathcal{B}], \omega_2(\mathcal{A}, \mathcal{B})), \quad (\mathcal{A}, d), (\mathcal{B}, e) \in \hat{\mathfrak{g}},$$

and introduces another commutator on  $\hat{\mathfrak{g}}$  in the form

$$\begin{aligned} [(\mathcal{A}, d), (\mathcal{B}, e)]_{\mathcal{R}} &= ([\mathcal{A}, \mathcal{B}]_{\mathcal{R}}, \omega_{2,\mathcal{R}}(\mathcal{A}, \mathcal{B})), \\ [\mathcal{A}, \mathcal{B}]_{\mathcal{R}} &= [\mathcal{R}\mathcal{A}, \mathcal{B}] + [\mathcal{A}, \mathcal{R}\mathcal{B}], \quad \omega_{2,\mathcal{R}}(\mathcal{A}, \mathcal{B}) = \omega_2(\mathcal{R}\mathcal{A}, \mathcal{B}) + \omega_2(\mathcal{A}, \mathcal{R}\mathcal{B}), \end{aligned} \quad (2)$$

where  $\mathcal{R} = (P_+ - P_-)/2$  and  $P_\pm$  are projectors on the Lie subalgebras  $\tilde{\mathfrak{g}}_\pm$ . On the space  $\hat{\mathfrak{g}}^*$ , dual to  $\hat{\mathfrak{g}}$  with respect to the scalar product  $((\mathcal{A}, d), (\mathcal{B}, e)) = \int_{y \in \mathbb{S}} dy (\mathcal{A}, \mathcal{B}) + de$ , the commutator (2) determines the Lie-Poisson bracket

$$\{\gamma, \mu\}_{\mathcal{R}}(l) = \int_{y \in \mathbb{S}} dy (l, [\nabla_l \gamma(l), \nabla_r \mu(l)]_{\mathcal{R}}) + c\omega_2(\nabla_l \gamma(l), \nabla_r \mu(l)) = (\nabla_l \gamma(l), \Theta \nabla_r \mu(l)), \quad (3)$$

where  $\gamma, \mu \in \mathcal{D}(\tilde{\mathfrak{g}}^*)$  are arbitrary smooth by Frechet functionals on  $\tilde{\mathfrak{g}}^* \simeq \tilde{\mathfrak{g}}$ , at a point  $(l, c) \in \hat{\mathfrak{g}}^*$ . Here  $l \in \tilde{\mathfrak{g}}^*$  is some matrix super-integro-differential operator of order  $q \in \mathbb{N}$ ,  $c \in \mathbb{C}$ ,  $\nabla_l, \nabla_r$  are left and right gradient operators accordingly,  $\Theta : T^*(\tilde{\mathfrak{g}}^*) \rightarrow T(\tilde{\mathfrak{g}}^*)$  is the Poisson operator generating the Lie-Poisson bracket (3) at a point  $l \in \tilde{\mathfrak{g}}^*$  and acting as  $\Theta : \nabla \gamma(l) \mapsto -[l - c\mathbf{1}\partial/\partial y, (\nabla \gamma(l))_-] + [l - c\mathbf{1}\partial/\partial y, \nabla \gamma(l)]_-$  for any  $\gamma \in \mathcal{D}(\tilde{\mathfrak{g}}^*)$ , the subscript "-" denotes the projection of the corresponding element from  $\tilde{\mathfrak{g}}$  on the Lie subalgebra  $\tilde{\mathfrak{g}}_- := \prod_{y \in \mathbb{S}} \mathfrak{g}_-$ , and  $T(\tilde{\mathfrak{g}}^*), T^*(\tilde{\mathfrak{g}}^*)$  are tangent and cotangent spaces to  $\tilde{\mathfrak{g}}^*$ .

The Casimir functionals  $\gamma_j \in \mathcal{I}(\hat{\mathfrak{g}}^*)$ ,  $j \in \mathbb{N}$ , of the central extension  $\hat{\mathfrak{g}}$ , whose left gradients obey the equalities such that  $[l - c\mathbf{1}\partial/\partial y, \nabla_l \gamma_j(l)] = 0$ , where  $\nabla_l \gamma_j(l) := \mathbf{1}\partial^j + \sum_{p < 2j} A_{j,p} D_\theta^p$ ,  $A_{j,p}$  are supermatrix-valued functions of suitable parity,  $j \in \mathbb{N}$ ,  $p \in \mathbb{Z}$ ,  $p < 2j$ , at a point  $(l, c) \in \hat{\mathfrak{g}}^*$ , and the  $\mathcal{R}$ -deformed Lie-Poisson bracket (3) give us the hierarchy of Lax type Hamiltonian flows on  $\tilde{\mathfrak{g}}^* \simeq \tilde{\mathfrak{g}}$ :

$$dl/dt_j = [(\nabla_l \gamma_j(l))_+, l - c\mathbf{1}\partial/\partial y], \quad j \in \mathbb{N}, \quad t_j \in \mathbb{R},$$

where the subscript "+" denotes the projection of the corresponding element from  $\tilde{\mathfrak{g}}$  on the Lie subalgebra  $\tilde{\mathfrak{g}}_+ := \prod_{y \in \mathbb{S}} \mathfrak{g}_+$ . One considers another hierarchy of Lax type Hamiltonian flows on the dual space  $\tilde{\mathfrak{g}}^*$ :

$$d\tilde{l}/dt_j = [(\nabla_l \gamma_j(\tilde{l}))_+, \tilde{l} - c\mathbf{1}\partial/\partial y], \quad j \in \mathbb{N}, \quad t_j \in \mathbb{R},$$

for some matrix super-integro-differential operator  $\tilde{l} \in \tilde{\mathfrak{g}}^*$  of order  $q \in \mathbb{N}$ , which is related with the operator  $l \in \tilde{\mathfrak{g}}^*$  by the generalized gauge transformation

$$\tilde{l}(0) - c\mathbf{1}\partial/\partial y = \mathcal{B}(0)^{-1}(l(0) - c\mathbf{1}\partial/\partial y)\mathcal{B}(0), \quad (4)$$

where  $\mathcal{B}(0) \in \tilde{\mathfrak{g}}_+$  is a matrix superdifferential operator of order  $s \in \mathbb{N}$  with constant coefficients, at the initial moment of the time  $t_j \in \mathbb{R}$  for every  $j \in \mathbb{N}$ .

**Theorem 1.** *If for every  $j \in \mathbb{N}$  at the initial moment of the time  $t_j \in \mathbb{R}$  matrix super-integro-differential operators  $l, \tilde{l} \in \tilde{\mathfrak{g}}^*$  of order  $q \in \mathbb{N}$  are related by the relationship (4), there exist such matrix superdifferential operators of orders  $q + s$  and  $s$  accordingly, where  $s \in \mathbb{Z}_+$ ,  $s < q$ , that the equalities*

$$l = \mathcal{A}\mathcal{B}^{-1}, \quad \tilde{l} = \mathcal{B}^{-1}(\mathcal{A} - c\partial\mathcal{B}/\partial y) \quad (5)$$

hold. The operators  $\mathcal{A}, \mathcal{B} \in \tilde{\mathfrak{g}}_+$  satisfy the following systems of evolution equations

$$\begin{aligned} d\mathcal{A}/dt_j &= (\nabla_l \gamma_j(l))_+ \mathcal{A} - \mathcal{A}(\nabla_l \gamma_j(\tilde{l}))_+ - c(\partial(\nabla_l \gamma_j(l))_+ / \partial y) \mathcal{B}, \\ d\mathcal{B}/dt_j &= (\nabla_l \gamma_j(l))_+ \mathcal{B} - \mathcal{B}(\nabla_l \gamma_j(\tilde{l}))_+, \quad j \in \mathbb{N}, \end{aligned} \quad (6)$$

which possess an infinite sequence of the conservation laws  $H_j \in \mathcal{D}(\tilde{\mathfrak{g}}_+ \times \tilde{\mathfrak{g}}_+)$ ,  $j \in \mathbb{N}$ , in the form

$$H_j(\mathcal{A}, \mathcal{B}) := \gamma_j(l)|_{l=\mathcal{A}\mathcal{B}^{-1}} = \gamma_j(\tilde{l})|_{\tilde{l}=\mathcal{B}^{-1}(\mathcal{A}-c\partial\mathcal{B}/\partial y)}.$$

The equalities (5) determine the Backlund transformation

$$P : (\mathcal{A}, \mathcal{B}) \in \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \mapsto (l, \tilde{l}) \in \tilde{\mathfrak{g}}^* \oplus \tilde{\mathfrak{g}}^*. \quad (7)$$

**Theorem 2.** *For every  $j \in \mathbb{N}$  the system of evolution equations (6), given on the subspace  $\tilde{\mathfrak{g}}_+ \times \tilde{\mathfrak{g}}_+ \subset \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}$ , is Hamiltonian with respect to the Poisson bracket  $\{.,.\}_{\mathcal{L}}$ , which arises as a reduction of the Poisson bracket  $\{.,.\}_{\tilde{\mathcal{L}}}$  with the corresponding Poisson operator  $\tilde{\mathcal{L}} : T^*(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}) \rightarrow T^*(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}})$  such as  $\tilde{\mathcal{L}} = (P')^{-1}(\Theta \oplus \tilde{\Theta})(P'^*)^{-1}$ , where  $\Theta, \tilde{\Theta} : T^*(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}) \rightarrow T^*(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}})$  are Poisson operators generating the Lie-Poisson bracket  $\{.,.\}_{\mathcal{R}}$  at points  $l, \tilde{l} \in \tilde{\mathfrak{g}}^*$  accordingly,  $P'^* : T^*(\tilde{\mathfrak{g}}^* \oplus \tilde{\mathfrak{g}}^*) \rightarrow T^*(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}})$  is an operator adjoint to the Frechet derivative  $P' : T(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}) \rightarrow T(\tilde{\mathfrak{g}}^* \oplus \tilde{\mathfrak{g}}^*)$  of the Backlund transformation (7) and  $(P'^*)^{-1}$  is an operator inverse to  $P'$ , on  $\tilde{\mathfrak{g}}_+ \times \tilde{\mathfrak{g}}_+$  and the Hamiltonians  $\bar{H}_j \in \mathcal{D}(\tilde{\mathfrak{g}}_+ \times \tilde{\mathfrak{g}}_+)$ ,  $j \in \mathbb{N}$ :*

$$\bar{H}_j(\mathcal{A}, \mathcal{B}) := \gamma_j(l)|_{l=\mathcal{A}\mathcal{B}^{-1}} + \gamma_j(\tilde{l})|_{\tilde{l}=\mathcal{B}^{-1}(\mathcal{A}-c\partial\mathcal{B}/\partial y)}.$$

The reductions of the hierarchy (6) on the coadjoint action orbits for the central extension  $\hat{\mathfrak{g}}$  with taking into account the Backlund transformation (7) lead to new integrable hierarchies of nonlinear dynamical systems on matrix functional supermanifolds of two commuting and one anticommuting independent variables, being Hamiltonian ones and possessing infinite sequences of conservation laws.

- [1] Oksana E. Hentosh. Lie-algebraic structure of the Lax-integrable  $(2|1+1)$ -dimensional supersymmetric matrix dynamical systems. *Ukr. Math. J.*, 69(10) : 1537–1560, 2018.

## The some solution of the Bryan-Pidduck equation

**Oleksii Hukalov**

(B. Verkin ILTPE of NASU, Kharkiv, Ukraine)

*E-mail:* hukalov@ilt.kharkov.ua

**Vyacheslav Gordevskyy**

(V. N. Karazin Kharkiv National University, Kharkiv, Ukraine)

*E-mail:* gordevskyy2006@gmail.com

The nonlinear integrodifferential Boltzmann equation [1] that describes the evolution of rarefied gases is one of the main equations of the kinetic theory of gases. The Boltzmann equation for the model of rough spheres (or the Bryan–Pidduck equation) has the form

$$D(f) = Q(f, f); \quad (1)$$

$$D(f) \equiv \frac{\partial f}{\partial t} + \left( V, \frac{\partial f}{\partial x} \right), \quad (2)$$

$$Q(f, f) \equiv \frac{d^2}{2} \int_{\mathbb{R}^3} dV_1 \int_{\mathbb{R}^3} d\omega_1 \int_{\Sigma} d\alpha B(V - V_1, \alpha) \left[ f(t, V_1^*, x, \omega_1^*) f(t, V^*, x, \omega^*) - f(t, V, x, \omega) f(t, V_1, x, \omega_1) \right]. \quad (3)$$

The problem of determination of the exact and approximate solutions of the Bryan–Pidduck equation in the explicit form is quite urgent. At present, the sole known exact solution of the Boltzmann equation is an expression usually called the Maxwell distribution or simply Maxwellian (after J. C. Maxwell, Scottish physicist). In the case of Maxwellians  $M$ , we get

$$D(f) = 0, \quad Q(f, f) = 0. \quad (4)$$

The solution to this equation (1)–(3) will be look for in the next form

$$f(t, x, V, \omega, u) = \int_{\mathbb{R}^3} \varphi(t, x, u) M(V, \omega, u) du. \quad (5)$$

As a measure of the deviation between the parts of equation (1) we will consider a uniform-integral error of the form:

$$\Delta = \sup_{(t,x) \in \mathbb{R}^4} \int_{\mathbb{R}^3} dV \int_{\mathbb{R}^3} d\omega \left| D(f) - Q(f, f) \right|. \quad (6)$$

In the paper [2], we were obtained sufficient conditions for the coefficient functions and hydrodynamic parameters appearing in the distribution, which enable one to make the analyzed error (6) as small as desired.

<b>V. Dryuma</b> <i>6D-Riemannian metric associated at the Navier-Stokes equations and its applications</i>	<b>29</b>
<b>K. Eftekharinasab</b> <i>Interplay of Global Implicit Functions and Critical Point Theory in Infinite Dimensional Spaces</i>	<b>32</b>
<b>M. Egwe</b> <i>Spherical Analysis on Fuzzy Lie Groups</i>	<b>32</b>
<b>V. Fedorchuk, V. Fedorchuk</b> <i>On the construction and classification of the common invariant solutions for the <math>(1 + 3)</math>-dimensional Euler-Lagrange-Born-Infeld and homogeneous Monge-Ampere equations</i>	<b>33</b>
<b>B. Feshchenko</b> <i>Normal forms of functions with degenerate critical points on surfaces whose stabilizers are homotopically non-trivial</i>	<b>35</b>
<b>M. R. Formica</b> <i>Nonlinear interpolation of <math>\alpha</math>-Holderian mappings with applications to quasilinear PDEs</i>	<b>36</b>
<b>J.F. Gálvez-Rodríguez, C. Martín-Aguado, M.A. Sánchez-Granero</b> <i>Riemann Integration on a space with a fractal structure</i>	<b>37</b>
<b>I. Gkeneralis</b> <i>Topological rigidity of quoric manifolds</i>	<b>39</b>
<b>O. Gok</b> <i>On the multiplicative order convergence on Banach lattice <math>f</math>-algebras</i>	<b>41</b>
<b>M. Golasiński</b> <i>Octonionic Stiefel manifolds and vector bundles</i>	<b>43</b>
<b>T. Goy</b> <i>Some series involving central binomial coefficients</i>	<b>44</b>
<b>O. Gutik</b> <i>On non-topologizable semigroups</i>	<b>46</b>
<b>H. Chi</b> <i>Existence and non-existence of cohomogeneity one Einstein metrics</i>	<b>48</b>
<b>I. Havrylenko, E. Petrov</b> <i>Stability of vertical minimal surfaces in three-dimensional sub-Riemannian manifolds</i>	<b>50</b>
<b>O. Ye. Hentosh</b> <i>Rational factorization of Lax type flows in the space dual to the centrally extended Lie algebra of matrix super-integro-differential operators</i>	<b>52</b>
<b>O. Hukalov, V. Gordevskyy</b> <i>The some solution of the Bryan-Pidduck equation</i>	<b>54</b>
<b>S. Ivković</b> <i>Disjoint dynamical properties of wedge operators</i>	<b>55</b>
<b>R. Ilemobade, T.G. Jaiyeola</b> <i>On semi-symmetric <math>(\alpha, \beta, \gamma)</math>-inverse quasigroup</i>	<b>55</b>
<b>O. Karlova</b> <i>About Rolewicz theorem on inversion of continuous bijection between <math>F</math>-spaces</i>	<b>57</b>
<b>L. Fardigola, K. Khalina</b> <i>On boundary controllability problems for the heat equation with variable coefficients on a half-axis</i>	<b>58</b>
<b>N. Kitazawa</b> <i>On explicit reconstruction of real algebraic maps locally like moment maps</i>	<b>60</b>
<b>S.D. Koval, R.O. Popovych</b> <i>Group classification of Kolmogorov backward equations with power diffusivity</i>	<b>63</b>