

International
Online Conference



**Algebraic
and Geometric
Methods of Analysis**

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Drozd Yu.

(Kyiv, Ukraine)

Maksymenko S.

(Kyiv, Ukraine)

Plaksa S.

(Kyiv, Ukraine)

Prishlyak A.

(Kyiv, Ukraine)

Bakhtin O.

(Kyiv, Ukraine)

Balan V.

(Bucharest, Romania)

Banakh T.

(Lviv, Ukraine)

Borysenko O.

(Kharkiv, Ukraine)

Cherevko Ye.

(Odesa, Ukraine)

Fedchenko Yu.

(Odesa, Ukraine)

Karlova O.

(Chernivtsi, Ukraine)

Kiosak V.

(Odessa, Ukraine)

Konovenko N.

(Odessa, Ukraine)

Lyubashenko V.

(Kyiv, Ukraine)

Matsumoto K.

(Yamagata, Japan)

Mormul P.

(Warsaw, Poland)

Mykhailyuk V.

(Chernivtsi, Ukraine)

Plachta L.

(Krakov, Poland)

Pokas S.

(Odessa, Ukraine)

Sabitov I.

(Moscow, Russia)

Savchenko O.

(Kherson, Ukraine)

Sergeeva A.

(Odessa, Ukraine)

Shelekhov A.

(Tver, Russia)

Zarichnyi M.

(Lviv, Ukraine)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

ORGANIZING COMMITTEE

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohlyk O.

Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.

Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

Modified quaternionic operator calculus and its application to micropolar elasticity

Klaus Gürlebeck

(Chair of Applied Mathematics, Bauhaus-Universität Weimar, 99423 Weimar, Germany)

E-mail: klaus.guerlebeck@uni-weimar.de

Dmitrii Legatiuk

(Chair of Applied Mathematics, Bauhaus-Universität Weimar, 99423 Weimar, Germany)

E-mail: dmitrii.legatiuk@uni-weimar.de

Original ideas for the extension of classical elasticity theory to account microeffects of a continuum go back to the work [1] of Cosserat brothers, where they introduced a new theory called the *Cosserat continuum*. The introduced theory grabbed attention of many scientists. Among others, works of Eringen [2], and Nowacki [3] significantly supported further development of the theory. Eringen introduced micro-inertia in the theory, which has led to renaming of the theory to the *micropolar elasticity*. From practical point of view, the micropolar theory models not only displacements of a continuum, as in the classical theory of elasticity, but also its rotations.

In this talk we introduce representation formulae for the solution of spatial boundary value problems of micropolar elasticity. The representation formulae are constructed in the framework of quaternionic analysis, which is a natural extension of the classical complex analysis to higher dimensions. The main toolbox for constructing representation formulae for boundary value problems of mathematical physics in hypercomplex analysis is the co-called quaternionic operator calculus [4, 5]. The essential ingredient is the T -operator (Teodorescu transform), which is a right inverse to the generalised Cauchy-Riemann operator. Accomplishing the T -operator with the F -operator (Cauchy-Bitsadze operator), the higher-dimensional generalisation of the classical Borel-Pompeiu formula can be obtained.

In this talk, we consider the following boundary value problem:

Problem 1. Let $\Omega \subset \mathbb{R}^3$ be a bounded simply connected domain with a sufficiently smooth boundary $\Gamma = \Gamma_0 \cup \Gamma_1$. A boundary value problem of the micropolar elasticity is formulated as follows

$$(\lambda + 2\mu + \kappa) \nabla \nabla \cdot \mathbf{u} - (\mu + \kappa) \nabla \times \nabla \times \mathbf{u} = -\kappa \nabla \times \boldsymbol{\varphi}, \quad (1)$$

$$(\alpha + \beta + \gamma) \nabla \nabla \cdot \boldsymbol{\varphi} - \gamma \nabla \times \nabla \times \boldsymbol{\varphi} - 2\kappa \boldsymbol{\varphi} = -\kappa \nabla \times \mathbf{u}, \quad (2)$$

with boundary conditions

$$\begin{cases} \mathbf{u} = \mathbf{g}_1 & \text{on } \Gamma_0, \\ \boldsymbol{\varphi} = \mathbf{g}_2 & \text{on } \Gamma_0, \end{cases} \quad \text{and} \quad \begin{cases} t_{lk} n_l = t_{(\mathbf{n})k} & \text{on } \Gamma_1, \\ m_{lk} n_l = m_{(\mathbf{n})k} & \text{on } \Gamma_1, \end{cases} \quad (3)$$

where \mathbf{u} is the displacement vector, $\boldsymbol{\varphi}$ is the vector of micropolar rotation, t_{lk} is the stress tensor, m_{lk} is the couple stress tensor, ρ is the material density, j is a rotational inertia, λ and μ are the Lamé parameters, $\kappa, \alpha, \beta, \gamma$ are material parameters of micropolar theory, n_j are components of the unit outer normal vector, $t_{(\mathbf{n})k}$ are given surface forces, and $m_{(\mathbf{n})k}$ are given surface moments.

After that, a quaternionic formulation of the boundary value problem (1)-(3) is considered [6]:

Proposition 2. Considering the displacement field $\mathbf{u} \in C^2(\Omega)$ and micropolar rotations $\boldsymbol{\varphi} \in C^2(\Omega)$ as pure quaternions, i.e. $\mathbf{u} = u_1 \mathbf{e}_1 + u_2 \mathbf{e}_2 + u_3 \mathbf{e}_3$, $\boldsymbol{\varphi} = \varphi_1 \mathbf{e}_1 + \varphi_2 \mathbf{e}_2 + \varphi_3 \mathbf{e}_3$, equations of micropolar elasticity (1)-(2) can be written as follows

$$\begin{aligned} D M_1 D \mathbf{u} + \kappa \text{Vec } D \boldsymbol{\varphi} &= 0, \\ \left(D - i \sqrt{\frac{2\kappa}{\gamma}} \right) M_2 \left(D + i \sqrt{\frac{2\kappa}{\gamma}} \right) \boldsymbol{\varphi} + \kappa \text{Vec } D \mathbf{u} &= 0, \end{aligned} \quad (4)$$

where the operators M_1 and M_2 are defined by

$$\begin{aligned} M_1 \mathbf{w} &:= -(\lambda + 2\mu + \kappa)w_0 - (\mu + \kappa)w_1 \mathbf{e}_1 - (\mu + \kappa)w_2 \mathbf{e}_2 \\ &\quad - (\mu + \kappa)w_3 \mathbf{e}_3, \\ M_2 \mathbf{w} &:= -(\alpha + \beta + \gamma)w_0 - \gamma w_1 \mathbf{e}_1 - \gamma w_2 \mathbf{e}_2 - \gamma w_3 \mathbf{e}_3, \end{aligned}$$

for a quaternion-valued function $\mathbf{w} = w_0 + w_1 \mathbf{e}_1 + w_2 \mathbf{e}_2 + w_3 \mathbf{e}_3$.

By reformulating the system as a system of operator equations, the questions of existence, regularity, stability and uniqueness can be studied by using the classical and modified versions of quaternionic operator calculus [4, 5]:

Theorem 3. *The system of equations*

$$\begin{cases} D M_1 D \mathbf{u} + \kappa \text{Vec } D \varphi & = 0, \\ \left(D - i\sqrt{\frac{2\kappa}{\gamma}} \right) M_2 \left(D + i\sqrt{\frac{2\kappa}{\gamma}} \right) \varphi + \kappa \text{Vec } D \mathbf{u} & = 0, \end{cases} \quad (5)$$

with Dirichlet boundary conditions

$$\begin{cases} \mathbf{u} = \mathbf{g}_1 & \text{on } \Gamma_0, \\ \varphi = \mathbf{g}_2 & \text{on } \Gamma_0, \end{cases}$$

is equivalent to the system of operator equations

$$\begin{cases} \mathbf{u} = F_\Gamma \tilde{\mathbf{g}}_1 + T M_1^{-1} F_\Gamma (\text{tr } T M_1^{-1} F_\Gamma)^{-1} Q_\Gamma \tilde{\mathbf{g}}_1 \\ \quad - \kappa T M_1^{-1} T \text{Vec } D \varphi, \\ \varphi = F_\alpha \tilde{\mathbf{g}}_2 + T_\alpha M_2^{-1} F_{-\alpha} (\text{tr } T_\alpha M_2^{-1} F_{-\alpha})^{-1} Q_\alpha \tilde{\mathbf{g}}_2 \\ \quad - \kappa T_\alpha M_2^{-1} T_{-\alpha} \text{Vec } D \mathbf{u}, \end{cases} \quad (6)$$

where $\tilde{\mathbf{g}}_1 = \mathbf{g}_1 + \kappa \text{tr } T M_1^{-1} T \text{Vec } D \varphi$ and $\tilde{\mathbf{g}}_2 = \mathbf{g}_2 + \kappa \text{tr } T_\alpha M_2^{-1} T_{-\alpha} \text{Vec } D \mathbf{u}$.

Further results related to the uniqueness of solution, as well as the estimate for a difference between the micropolar model and the classical model of elasticity, will be presented in the talk.

REFERENCES

- [1] E. Cosserat, F. Cosserat. Sur la theorie de l'elasticite, *Ann. de l'Ecole Normale de Toulouse*, 10(1), 1896.
- [2] A.C. Eringen. Linear theory of micropolar elasticity, *Journal of Mathematics and Mechanics*, 15(6), pp. 909-923, 1966.
- [3] W. Nowacki. *Theory of micropolar elasticity*, Course Held at the Department for Mechanics of Deformable Bodies, 1970.
- [4] K. Gürlebeck, W. Sprößig. *Quaternionic and Clifford calculus for physicists and engineers*, Wiley, Chichester, 1997.
- [5] K. Gürlebeck, K. Habetha, W. Sprößig. *Application of holomorphic functions in two and higher dimensions*, Springer, 2016.
- [6] K. Gürlebeck, D. Legatiuk. Quaternionic operator calculus for boundary value problems of micropolar elasticity, *Topics in Clifford Analysis*, Springer series "Trends in Mathematics", pp. 221-234, 2019.

Kh. F. Kholturayev <i>Perfect metrizable of the functor of idempotent measures</i>	75
Y. Khomych <i>Quasiareal deformation of surfaces of positive Gauss curvature</i>	77
V. Kiosak, O. Lesechko <i>Canonical infinitesimal deformations of metrics of pseudo-Riemannian spaces</i>	78
R. Salimov, B. Klishchuk <i>On the behavior at infinity of ring Q-homeomorphisms</i>	79
T. Kolomiets, A. Pogorui <i>Elements of probability theory and measures with values in hypercomplex algebras</i>	81
N. Konovenko <i>The invariants of planar 3-webs with respect to group of symplectic diffeomorphisms, for the case of the conformal group</i>	84
E. Kudryavtseva <i>Topology of spaces of smooth functions and gradient-like flows with prescribed singularities on surfaces</i>	85
G. Kuduk <i>Nonlocal problem with integral conditions for homogeneous system of partial differential equations of second order</i>	87
I. Kuznietsova, Yu. Soroka <i>Realization of groups as fundamental groups of orbits of smooth maps</i>	88
K. Gürlebeck, D. Legatiuk <i>Modified quaternionic operator calculus and its application to micropolar elasticity</i>	90
S. Maksymenko, E. Polulyakh <i>On non-Hausdorff manifolds of dimension 1</i>	92
S. Maksymenko <i>Symplectomorphisms preserving smooth functions on surfaces</i>	93
M. Maloid-Hliebova <i>Second classical Zariski topology of multiplicative module</i>	94
I. Marko <i>Incomplete spaces of idempotent measures</i>	95
N. Mazurenko, M. Zarichnyi <i>Hyperspaces of convex sets related to idempotent mathematics</i>	96
A. Mednykh <i>Volumes of knots and links in spaces of constant curvature</i>	98
R. Mohseni, R. A. Wolak <i>Twistor spaces on foliated manifolds</i>	99
P. Mormul <i>Two problems in nonholonomic geometry (in quest of a co-worker)</i>	100
F. Mukhamadiev <i>The local τ-density of a linearly ordered spaces</i>	101
T. Obikhod <i>Entropy and phase transitions in Calabi-Yau space</i>	102
A. Orevkova <i>Reducing singularities of smooth functions to normal forms</i>	104
T. Osipchuk <i>On m-convexity and m-semiconvexity of sets in Euclidean spaces</i>	106
V. Ostrovskiy, O. Ostrovska, D. Proskurin, Yu. Samoilenko <i>On representations of q_{ij}-commuting isometries</i>	108
J.F. Peters <i>Homotopic Nerve Complexes with Free Group Presentations</i>	110
P. Laurain, M. Petrace <i>Uniform measures in Euclidean space</i>	112