



International

Scientific

Conference

Algebraic and Geometric Methods of Analysis

May 24-27, 2022, Odesa, Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology, Ukraine
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Co-Chairs:

Maksymenko S.
(*Kyiv, Ukraine*)

Prishlyak A.
(*Kyiv, Ukraine*)

Balan V.
(*Bucharest, Romania*)

Fedchenko Yu.
(*Odesa, Ukraine*)

Matsumoto K.
(*Yamagata, Japan*)

Banakh T.
(*Lviv, Ukraine*)

Karlova O.
(*Chernivtsi, Ukraine*)

Mormul P.
(*Warsaw, Poland*)

Bolotov D.
(*Kharkiv, Ukraine*)

Kiosak V.
(*Odesa, Ukraine*)

Plachta L.
(*Krakov, Poland*)

Cherevko Ye.
(*Odesa, Ukraine*)

Konovenko N.
(*Odesa, Ukraine*)

Polulyakh Ye.
(*Kyiv, Ukraine*)

Savchenko O.
(*Kherson, Ukraine*)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONTU;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONTU;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONTU;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";

ORGANIZING COMMITTEE

Konovenko N.
Maksymenko S.

Fedchenko Yu.
Cherevko Ye.

Osadchuk Ye.
Sergeeva O.

Soroka Yu.

The symplectic invariants of 3-webs

Konovenko Nadiia

(ONTU, Odesa, Ukraine)

E-mail: ngkonovenko@gmail.com

The classical web geometry ([1],[2],[4]) studies invariants of foliation families with respect to pseudogroup of diffeomorphisms. Thus for the case of planar 3-webs the basic semi invariant is the Blaschke curvature ([3]). It is also curvature of the Chern connection ([4]) that are naturally associated with a planar 3-web.

Let $\mathbf{D} \subset \mathbb{R}^2$ be a connected and simply connected domain in the plane, equipped with symplectic structure given by differential 2-form $\Omega = dx \wedge dy$ in the standard coordinates on the plane.

Remind that a 3-web in the domain is a family of three foliations being in general position. We'll assume that these foliations are integral curves of differential 1-forms ω_i , $i = 1, 2, 3$, and write

$$W_3 = \langle \omega_1, \omega_2, \omega_3 \rangle,$$

where $\omega_i \in \Omega^1(\mathbf{D})$ are such differential 1-forms that $\omega_i \wedge \omega_j \neq 0$ in \mathbf{D} , if $i \neq j$.

Definition 1. We say that two planar 3-webs W_3 and \tilde{W}_3 given in domains \mathbf{D} and $\tilde{\mathbf{D}}$ respectively are symplectively equivalent if there is a symplectomorphism $\phi : \mathbf{D} \rightarrow \tilde{\mathbf{D}}$, such that $\phi(W_3) = \tilde{W}_3$.

Proposition 2. Let $W_3 = \langle \omega_1, \omega_2, \omega_3 \rangle$ and $\tilde{W}_3 = \langle \tilde{\omega}_1, \tilde{\omega}_2, \tilde{\omega}_3 \rangle$ be two planar 3-webs in domains \mathbf{D} and $\tilde{\mathbf{D}}$ respectively given by normalized

$$\omega_1 + \omega_2 + \omega_3 = 0. \quad (1)$$

differential forms. Then a diffeomorphism $\phi : \mathbf{D} \rightarrow \tilde{\mathbf{D}}$ establishes a symplectic equivalence of 3-webs if and only if

$$\phi^*(\tilde{\omega}_i) = \varepsilon \omega_{\sigma(i)},$$

where $(\sigma, \varepsilon) \in \mathbb{A}_3 \times \mathbb{Z}_2$, and $\mathbb{A}_3 \subset \mathbb{S}_3$ is the subgroup of even permutations and $\mathbb{Z}_2 = \{1, -1\}$.

In our case normalization (1) and the above proposition shows that the Chern form γ is itself symplectic invariant of 3-webs.

Let's write down γ in following form

$$\gamma = x_1 \omega_1 + x_2 \omega_2 + x_3 \omega_3,$$

where functions $x_i \in C^\infty(\mathbf{D})$ are barycentric coordinates of γ , i.e.

$$x_1 + x_2 + x_3 = 1.$$

Then we have

$$\begin{aligned} d\omega_1 &= (x_3 - x_2) \omega_1 \wedge \omega_2, \\ d\omega_2 &= (x_1 - x_3) \omega_1 \wedge \omega_2, \\ d\omega_3 &= (x_2 - x_1) \omega_1 \wedge \omega_2. \end{aligned}$$

Using the second normalization (1) condition we'll rewrite these relations in the following form

$$\begin{aligned} d\omega_i &= \lambda_i \Omega, \quad i = 1, 2, 3, \\ \lambda_1 &= x_3 - x_2, \lambda_2 = x_1 - x_3, \lambda_3 = x_2 - x_1, \end{aligned} \quad (2)$$

and

$$x_1 = \frac{1}{3}(1 + \lambda_2 - \lambda_3), x_2 = \frac{1}{3}(1 + \lambda_3 - \lambda_1), x_3 = \frac{1}{3}(1 + \lambda_1 - \lambda_2).$$

Theorem 3. *Functions*

$$\begin{aligned} J_1 &= \lambda_1^2 + \lambda_2^2 + \lambda_3^2, \\ J_2 &= \lambda_1^2 \lambda_2^2 + \lambda_1^2 \lambda_3^2 + \lambda_2^2 \lambda_3^2, \\ J_w &= (\lambda_2^2 - \lambda_1^2)(\lambda_3^2 - \lambda_1^2)(\lambda_3^2 - \lambda_2^2) \\ J_3 &= \lambda_1^2 \lambda_2^2 \lambda_3^2 \end{aligned}$$

are symplectic invariants of 3-webs.

REFERENCES

- [1] Akivis M. A., "Differential geometry of webs", Itogi Nauki i Tekhniki. Ser. Probl. Geom., 15, VINITI, Moscow, 1983, 187–213; J. Soviet Math., 29:5
- [2] Blaschke W., Bol G.. Geometrie der Gewebe. Vol. 335. No. 8. Berlin, 1938.
- [3] Blaschke W. Einführung in die Geometrie der Waben, Birkhäuser 1955
- [4] Chern S.-S., Web geometry. Bulletin of the American Mathematical Society 6.1 (1982) : 1-8.

Зміст

M. Amram <i>On the algorithm of degenerations and fundamental groups as a tool to understand algebraic surfaces</i>	2
D. Bolotov <i>Foliations on closed three-dimensional Riemannian manifolds with a bounded mean curvature of leaves</i>	3
Enzo Bonacci <i>Topological issues about the 6D ISST in Physics</i>	4
Francisco Bulnes <i>Characterized cycles integration on \mathcal{D}-modules as solutions in \mathbb{L}-holomorphic bundles</i>	5
D. Cheban <i>One-dimensional Monotone Non-autonomous Dynamical Systems and Strange Nonchaotic Attractors</i>	6
Y. Cherevko, V. Berezovski, J. Mikeš, Y. Fedchenko <i>Holomorphically Projective Mappings of Kähler Manifolds Preserving The Generalized Einstein Tensor</i>	8
V. Chernov, R. Sadykov <i>Some questions about virtual Legendrian knots</i>	10
Luca Di Beo <i>Morse-Smale flows in the Boy's surface</i>	11
Yu. Drozd <i>Morita equivalence of non-commutative Noetherian schemes</i>	12
K. Eftekharinasab <i>Some critical point results for Fréchet manifolds</i>	14
V. Fedorchuk, V. Fedorchuk <i>On partial preliminary group classification of some class of $(1 + 3)$-dimensional Monge-Ampere equations. One-dimensional Galilean Lie algebras.</i>	16
N. Glazunov <i>On packing and lattice packing of Minkowski-Chebyshev balls</i>	18
Omer GOK <i>Unbounded order and norm convergence of some operators on Banach lattices</i>	20
J. Ham, J. Lee <i>An explicit formula for the A-polynomial of the knot with Conway's notation $C(2n, 4)$</i>	21
N. Konovenko <i>The symplectic invariants of 3-webs</i>	23
Yaacov Kopeliovich <i>Solutions to Mumford's second problem on theta functions</i>	25
I. Kuznietsova, Yu. Soroka <i>Geometric interpretation of first Betti numbers of orbits of smooth functions</i>	26
J-P. Magnot <i>On diffeological principal bundles of non-formal pseudo-differential operators over formal ones</i>	28
S. Maksymenko <i>Topological actions of wreath products</i>	29
J-L. Mo <i>The geometrical properties of degenerations of curves and surfaces</i>	30
P. Mormul <i>Nilpotent approximations in the Goursat Monster Tower</i>	32