



International
Scientific Conference

Algebraic and Geometric Methods of Analysis

26-30 may 2020
Odesa, Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odessa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Odessa I. I. Mechnikov National University
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

PROGRAM COMMITTEE

Chairman: Prishlyak A. (<i>Kyiv, Ukraine</i>)	Kiosak V. (<i>Odesa, Ukraine</i>)	Pokas S. (<i>Odesa, Ukraine</i>)
Balan V. (<i>Bucharest, Romania</i>)	Kirillov V. (<i>Odesa, Ukraine</i>)	Polulyakh E. (<i>Kyiv, Ukraine</i>)
Banakh T. (<i>Lviv, Ukraine</i>)	Konovenko N. (<i>Odesa, Ukraine</i>)	Sabitov I. (<i>Moscow, Russia</i>)
Bolotov D. (<i>Kharkiv, Ukraine</i>)	Lyubashenko V. (<i>Kyiv, Ukraine</i>)	Savchenko A. (<i>Kherson, Ukraine</i>)
Borysenko O. (<i>Kharkiv, Ukraine</i>)	Maksymenko S. (<i>Kyiv, Ukraine</i>)	Sergeeva A. (<i>Odesa, Ukraine</i>)
Cherevko Ye. (<i>Odesa, Ukraine</i>)	Matsumoto K. (<i>Yamagata, Japan</i>)	Shelekhov A. (<i>Tver, Russia</i>)
Fedchenko Yu. (<i>Odesa, Ukraine</i>)	Mormul P. (<i>Warsaw, Poland</i>)	Volkov V. (<i>Odesa, Ukraine</i>)
Karlova O. (<i>Chernivtsi, Ukraine</i>)	Mykhailyuk V. (<i>Chernivtsi, Ukraine</i>)	Zarichnyi M. (<i>Lviv, Ukraine</i>)
	Plachta L. (<i>Krakov, Poland</i>)	

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Svytyy I., Dean of the Faculty of Computer Systems and Automation.

ORGANIZING COMMITTEE

Kirillov V.
Konovenko N.
Fedchenko Yu.

Maksymenko S.
Cherevko Ye.

Osadchuk E.
Prus A.

ІНТЕРНАЦІОНАЛЬНИЙ ЦЕНТР СПІВРОБІТНИЦТВА

On the geometry of submersions

G. M. Abdishukurova

(National University of Uzbekistan, Tashkent, 100174, Tashkent, Uzbekistan)

E-mail: Abdishukurova93@yandex.ru

A. Ya. Narmanov

(National University of Uzbekistan, Tashkent, 100174, Tashkent, Uzbekistan)

E-mail: narmanov@yandex.ru

Let M be a smooth connected Riemannian manifold of dimension n with Riemannian metric g .

Definition 1. Differentiable mapping $\pi : M \rightarrow B$ of maximal rank, where B is a smooth Riemannian manifold of dimension m , called submersion for $n > m$.

Submersion of $\pi : M \rightarrow B$ generates a foliation F of dimension $k = n - m$ on the manifold M , whose leaves are the submanifolds $L_p = \pi^{-1}(p), p \in B$. For a point $q \in L_p$ we denote by $T_q F$ the tangent space of the leaf L_p at the point q , by $H(q)$ the orthogonal complement of the tangent space $T_q F$ of the leaf L_p , i.e. $T_q M = T_q F \oplus H(q)$. We have two distributions $TF : q \rightarrow T_q F, H : q \rightarrow H(q)$. Each vector field X can be represented as $X = X^v + X^h$, where X^v, X^h are the orthogonal projections of X onto TF, H respectively. Here, for convenience, TF, H are considered as subbundles of the tangent bundle TM . If $X^h = 0$, then X is called a vertical field (it is tangent to the foliation), and if $X^v = 0$, then X is called a horizontal field.

Definition 2. A diffeomorphism $\varphi : M \rightarrow M$ is called a diffeomorphism of the foliated manifold (M, F) , if the image $\varphi(L_\alpha)$ of each leaf L_α is a leaf of the foliation F .

The diffeomorphism $\varphi : M \rightarrow M$ of the foliated manifold (M, F) , is denoted by $\varphi : (M, F) \rightarrow (M, F)$. The set of all diffeomorphisms of a foliated manifold is denoted by $Diff_F(M)$. The set $Diff_F(M)$ is a group with respect to the superposition of mappings and is a subgroup of the group $Diff(M)$ of diffeomorphisms of the manifold M . The group $Diff_F(M)$ was studied in [2], in particular, it was proved that this group is a closed subgroup of the group $Diff(M)$ with respect to a compactly open topology.

Definition 3. A diffeomorphism $\varphi : (M, F) \rightarrow (M, F)$ is called an isometry of the foliated manifold (M, F) , if the restriction of the mapping φ to each leaf of the foliation F is an isometry, that is, for each leaf L_α the map $\varphi : L_\alpha \rightarrow \varphi(L_\alpha)$ is an isometry between the manifolds L_α and $\varphi(L_\alpha)$.

Denote by $G_F(M)$ the set of isometries of the foliated manifold (M, F) . The group $G_F(M)$ is subgroup of $Diff(M)$ and therefore it is topological group in compact open topology.

Let us consider submersion $\pi : R^{n+1} \rightarrow R^1$, where

$$\pi(x_1, x_2, \dots, x_n, x_{n+1}) = x_{n+1} - f(x_1, x_2, \dots, x_n), \quad (1)$$

$f(x_1, x_2, \dots, x_n)$ is a differentiable function.

Theorem 4. Diffeomorphism $\varphi : R^{n+1} \rightarrow R^{n+1}$, defined by formula

$$\varphi_\lambda(x_1, x_2, \dots, x_n, x_{n+1}) = (x_1, x_2, \dots, x_n, x_{n+1} + \lambda\pi) \quad (2)$$

at $\lambda \neq -1$ is an isometry of foliation, generated by submersion (1).

Theorem 5. The set of diffeomorphisms

$$G_\Lambda = \{\varphi_\lambda : \lambda \in R^1, \lambda \neq -1\}, \quad (3)$$

is a subgroup of the group $G_F(M)$.

Using the mapping $\varphi_\lambda \rightarrow \lambda$ we identify the set G_Λ with the set $R^1 \setminus \{-1\}$ of real numbers other than -1 . On the set $R^1 \setminus \{-1\}$ we define the multiplication as follows

$$\lambda_1 \cdot \lambda_2 \rightarrow \lambda_1 + \lambda_2 + \lambda_1 \lambda_2, \quad (4)$$

The inverse element is determined by the formula

$$\lambda \rightarrow -\frac{\lambda}{1 + \lambda} \quad (5)$$

and it is obvious that they are differentiable. Therefore, we have following.

Proposition 6. *The set G_Λ is a one-dimensional Lie group.*

Example 7. Consider the submersion $\pi : R^3 \rightarrow R^1$, where $\pi(x_1, x_2, x_3) = x_3 - f(x_1, x_2)$, $f(x_1, x_2) = x_1^2 + x_2^2$. This submersion generates a two-dimensional foliation F . The following vector fields

$$V_1 = \frac{\partial}{\partial x_1} + 2x_1 \frac{\partial}{\partial x_3}, V_2 = \frac{\partial}{\partial x_2} + 2x_2 \frac{\partial}{\partial x_3}$$

are vertical vector fields. Vector field

$$X = -x_2 \frac{\partial}{\partial x_1} + x_1 \frac{\partial}{\partial x_2} + \frac{\partial}{\partial x_3}$$

is a foliated vector field for the foliation F , as shown by the following equalities $[V_1, X] = V_2$, $[V_2, X] = -V_1$. It is known that the flow of foliated vector field consists of diffeomorphisms of foliated manifold (M, F) [1]. The vector field X is a Killing vector field. Therefore, the flow of a vector field X consists of isometries of a foliated manifold. Indeed, the flow of the vector field X consists of diffeomorphisms

$$x \rightarrow A(t)x + bt$$

where $t \in R$, $b = \{0, 0, 1\}^T$, $x = (x_1, x_2, x_3)^T$,

$$A(t) = \begin{pmatrix} \cos t & -\sin t & 0 \\ \sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which are isometries of the foliated manifold (F, R^3) .

Theorem 8. *Suppose for a vector field*

$$V = \sum_{i=1}^n \xi_i \frac{\partial}{\partial x_i}$$

holds equality $V(f) = 0$. Then the flow of the vector field

$$X = V + \frac{\partial}{\partial x_{n+1}}$$

consists of diffeomorphisms of the foliated manifold (F, R^{n+1}) generated by submersion (1). If the field V is a Killing field, then the flow of the vector field X consists of isometries of the foliated manifold (F, R^{n+1}) .

REFERENCES

- [1] Molino P. Riemannian foliations. Boston–Basel: Burkhauser, 1988.
- [2] Narmanov A.Y., Zoyidov A.N. On the group of diffeomorphisms of foliated manifolds. Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Kompyuternye Nauki, 2020, vol. 30, issue 1, pp. 49–58.

ЗМІСТ

G. M. Abdishukurova, A. Ya. Narmanov <i>On the geometry of submersions</i>	3
B. N. Apanasov <i>Hyperbolic 4-cobordisms, Teichmuller spaces and quasiregular mappings in space</i>	5
Aymaz I., Kansu M. <i>Representation of gravi-electromagnetism using matrix algebra</i>	7
V. Bilet, O. Dovgoshey <i>Uniqueness of pretangent spaces at infinity</i>	9
Bolotov D. <i>Foliations of 3-manifolds with small module of mean curvature</i>	10
Bolsinov A. V. <i>On integrability of geodesic flows on 3-dimensional manifolds</i>	11
E. Bonacci <i>Algebraic and geometric questions about the EM helix</i>	12
Borisenko A. A., Sukhorebska D. D. <i>Geodesics on regular tetrahedra in spherical space</i>	13
F. Bulnes <i>Motivic hypercohomology solutions in field theory II</i>	14
I. Denega <i>Estimate of maximum of the products of inner radii of mutually non-overlapping domains</i>	16
A. Dudko, V. Pivovarchik <i>Inverse problem for tree of Stieltjes strings</i>	18
N. Glazunov <i>Formal groups and algebraic cobordism</i>	20
O. Gok <i>A note on tensor product of Archimedean vector lattices</i>	22
E. Gül. <i>Trace Regularization Problem On a Banach Space</i>	24
O. Ye. Hentosh <i>Centrally extended generalization of the superconformal loop Lie algebra and integrable heavenly type systems on supermanifolds</i>	26
B. Hladysh, A. Prishlyak <i>Structure of functions on an oriented 2-manifold with the boundary</i>	28
D. A. Juraev <i>The Cauchy problem for matrix factorizations of the Helmholtz equation in a multidimensional bounded domain</i>	30
A. Kachurovskii <i>Fejer Sums and the von Neumann Ergodic Theorem</i>	31
B. N. Khabibullin, R. R. Muryasov <i>Mixed volumes/areas and distribution of zeros of holomorphic functions</i>	33
B. Klishchuk, R. Salimov <i>On the behavior at infinity of one class of homeomorphisms</i>	35
A. Kravchenko, S. Maksymenko <i>Automorphisms of cellular divisions of 2-sphere induced by functions with isolated critical points</i>	37
A. Kushner, E. Kushner, R. Matviichuk <i>Dynamics and exact solutions of linear PDEs</i>	39