

International scientific conference

**“Algebraic and Geometric
Methods of Analysis”**

Book of abstracts



May 28 - June 3, 2019

Odesa, Ukraine

Conference webpage: imath.kiev.ua/~topology/conf/agma2019/

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences
- History and methodology of teaching in mathematics

ORGANIZERS

- The Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- The Institute of Mathematics of the National Academy of Sciences of Ukraine
- Odessa I. I. Mechnikov National University
- Taras Shevchenko National University of Kyiv
- The International Geometry Center

PROGRAM COMMITTEE

| | | |
|--|---|---|
| Chairman: Prishlyak A. (Kyiv, Ukraine) | Konovenko N. (Odesa, Ukraine) | Pokas S. (Odesa, Ukraine) |
| Balan V. (Bucharest, Romania) | Lyubashenko V. (Kyiv, Ukraine) | Polulyakh E. (Kyiv, Ukraine) |
| Banakh T. (Lviv, Ukraine) | Maksymenko S. (Kyiv, Ukraine) | Sabitov I. (Moscow, Russia) |
| Fedchenko Yu. (Odesa, Ukraine) | Matsumoto K. (Yamagata, Japan) | Savchenko A. (Kherson, Ukraine) |
| Fomenko A. (Moscow, Russia) | Mikesh J. (Olomouc, Czech Republic) | Sergeeva A. (Odesa, Ukraine) |
| Fomenko V. (Taganrog, Russia) | Mormul P. (Warsaw, Poland) | Shvets V. (Odesa, Ukraine) |
| Haddad M. (Wadi al-Nasara, Syria) | Moskaliuk S. (Wien, Austria) | Shelekhov A. (Tver, Russia) |
| Karlova O. (Chernivtsi, Ukraine) | Mykhailyuk V. (Chernivtsi, Ukraine) | Vlasenko I. (Kyiv, Ukraine) |
| Kiosak V. (Odessa, Ukraine) | Nykyforchyn O. (Ivano-Frankivsk, Ukraine) | Volkov V. (Odessa, Ukraine) |
| Kirillov V. (Odesa, Ukraine) | Plachta L. (Krakov, Poland) | Zadorozhnyj V. (Odesa, Ukraine) |
| | | Zarichnyi M. (Lviv, Ukraine) |

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Svytyy I., Dean of the Faculty of Computer Systems and Automation.

ORGANIZING COMMITTEE

Kirillov V.
Konovenko N.
Fedchenko Yu.

Prus A.
Osadchuk E.

Maksymenko S.
Khudenko N.
Cherevko E.

ФІТБ ОНАФТ

Approximation by Fourier sums and interpolation trigonometric polynomials in classes of differentiable functions with high exponents of smoothness

A. S. Serdyuk, I. V. Sokolenko

(Institute of Mathematics of the National Academy of Sciences of Ukraine, Kyiv, Ukraine)

E-mail: serdyuk@imath.kiev.ua, sokol@imath.kiev.ua

Let C and L_p , $1 \leq p \leq \infty$, be the spaces of 2π -periodic functions with the standart norms $\|\cdot\|_C$ and $\|\cdot\|_p$. Further, let $W_{\beta,p}^r$, $1 \leq p \leq \infty$, be the sets of all 2π -periodic functions f , representable as convolutions of the form

$$f(x) = \frac{a_0}{2} + \frac{1}{\pi} \int_{-\pi}^{\pi} \varphi(x-t) B_{r,\beta}(t) dt, \quad a_0 \in \mathbb{R}, \quad \varphi \perp 1, \quad \|\varphi\|_p \leq 1, \quad (1)$$

where $B_{r,\beta}(\cdot)$ are Weyl-Nagy kernels of the form

$$B_{r,\beta}(t) = \sum_{k=1}^{\infty} k^{-r} \cos\left(kt - \frac{\beta\pi}{2}\right), \quad r > 0, \quad \beta \in \mathbb{R}. \quad (2)$$

The classes $W_{\beta,p}^r$ are called as Weyl-Nagy classes (see, e.g., [1]). If $r \in \mathbb{N}$ and $\beta = r$, then the functions of the form (8) are the well-known Bernoulli kernels and the classes $W_{\beta,p}^r$ coincide with the well-known classes W_p^r , which consist of 2π -periodic functions with absolutely continuous derivatives up to $(r-1)$ -th order inclusive and such that $\|f^{(r)}\|_p \leq 1$ and $f^{(r)}(x) = \varphi(x)$ for almost everywhere $x \in \mathbb{R}$, where φ is the function from (10).

For arbitrary $\mathfrak{N} \subset X$, where $X = C$ or L_p , $1 \leq p \leq \infty$, we consider the quantity

$$\varepsilon_n(\mathfrak{N})_X = \sup_{f \in \mathfrak{N}} \|f(\cdot) - S_{n-1}(f; \cdot)\|_X, \quad (3)$$

where $S_{n-1}(f; x)$ is the partial Fourier sum of order $n-1$ of the function f .

In the case of Weyl-Nagy classes $W_{\beta,\infty}^r$ and $X = C$ for the exact upper bounds (3) the following asymptotic estimate holds

$$\varepsilon_n(W_{\beta,\infty}^r)_C = \frac{4}{\pi^2} \frac{\ln n}{n^r} + O\left(\frac{1}{n^r}\right), \quad r > 0, \quad \beta \in \mathbb{R}. \quad (4)$$

For $r \in \mathbb{N}$ and $\beta = r$ this estimate was obtained by A.N. Kolmogorov (1935), for arbitrary $r > 0$ by V.T. Pinkevich (1940) and S.M. Nikol'skii (1941). In the general case the estimate (4) follows from results, which were obtained in the works of A.V. Efimov (1960) and S.A. Telyakovskii (1961). It should be also noticed, that a similar asymptotic equality holds for the classes $W_{\beta,1}^r$ in the metric of the space L_1 . In these works the parameters r and β of the Weyl-Nagy classes were assumed to be fixed, and the question about the dependence of the remainder term in the estimate (4) on these parameters was not considered.

The character of the dependence on r and β of the remainder term in estimate (4) was investigated by I.G. Sokolov (1955), S.G. Selivanova (1955), G.I. Natanson (1961), S.A. Telyakovskii (1968, 1989) and S.B. Stechkin (1980). In the work of S.B. Stechkin [2] the asymptotic behavior, as $n \rightarrow \infty$ and $r \rightarrow \infty$, of the quantities $\varepsilon_n(W_{\beta,\infty}^r)_C$ was completely investigated. Besides, S.B. Stechkin [2, theorem 4] proved that for rapidly growing r the remainder can be improved. Namely, for arbitrary $r \geq n+1$

and $\beta \in \mathbb{R}$ the following equality holds:

$$\varepsilon_n(W_{\beta,\infty}^r)_C = \frac{1}{n^r} \left(\frac{4}{\pi} + O(1) \left(1 + \frac{1}{n} \right)^{-r} \right), \quad (5)$$

where $O(1)$ is a quantity uniformly bounded with respect to n, r and β . If $r/n \rightarrow \infty$, then the estimate (5) becomes the asymptotic equality. It also follows from [2] that for the quantity $\varepsilon_n(W_{\beta,1}^r)_{L_1}$ the analogous estimate to (5) takes place. S.A. Telyakovskii (1989) showed that the remainder in formulas (5) can be replaced by a smaller one, namely, write $O(1)(1 + 2/n)^{-r}$ instead of $O(1)(1 + 1/n)^{-r}$.

We establish generalized analogs of estimates (5) for quantities $\varepsilon_n(W_{\beta,p}^r)_C$ and $\varepsilon_n(W_{\beta,1}^r)_{L_p}$, respectively, for arbitrary values $1 \leq p \leq \infty$.

Theorem 1. *Let $1 \leq p \leq \infty$, $n \in \mathbb{N}$ and $\beta \in \mathbb{R}$. Then for $r \geq n + 1$ the following estimates hold:*

$$\varepsilon_n(W_{\beta,p}^r)_C = \frac{1}{n^r} \left(\frac{\|\cos t\|_{p'}}{\pi} + O(1) \left(1 + \frac{1}{n} \right)^{-r} \right), \quad (6)$$

$$\varepsilon_n(W_{\beta,1}^r)_{L_p} = \frac{1}{n^r} \left(\frac{\|\cos t\|_p}{\pi} + O(1) \left(1 + \frac{1}{n} \right)^{-r} \right), \quad (7)$$

where $1/p + 1/p' = 1$ and $O(1)$ are quantities uniformly bounded in all analyzed parameters. The estimates (6) and (7) are the asymptotic equalities, as $r/n \rightarrow \infty$.

Let $f \in C$. By $\tilde{S}_{n-1}(f; x)$ we denote a trigonometric polynomial of degree $n - 1$, that interpolates $f(x)$ at the equidistant nodes $x_k^{(n-1)} = \frac{2k\pi}{2n-1}$, $k \in \mathbb{Z}$, i.e., such that $\tilde{S}_{n-1}(f; x_k^{(n-1)}) = f(x_k^{(n-1)})$, $k \in \mathbb{Z}$.

For $\mathfrak{N} \subset C$ and $X = C$ or $X = L_p$, $1 \leq p \leq \infty$, consider the following approximative characteristic

$$\tilde{\varepsilon}_n(\mathfrak{N})_X = \sup_{f \in \mathfrak{N}} \|f(\cdot) - \tilde{S}_{n-1}(f; \cdot)\|_X. \quad (8)$$

The problems of finding of asymptotic behavior for quantity of the form (8) in important functional classes \mathfrak{N} was investigated by S.M. Nikol'skii, V.P. Motornyi, A.I. Stepanets, A.S. Serdyuk, and others.

The following statement is true [3].

Theorem 2. *Let $1 \leq p \leq \infty$, $n \in \mathbb{N}$ and $\beta \in \mathbb{R}$. Then for $r \geq n + 1$ the following estimates hold:*

$$\tilde{\varepsilon}_n(W_{\beta,p}^r)_C = \frac{1}{n^r} \left(\frac{2}{\pi} \|\cos t\|_{p'} + O(1) \left(1 + \frac{1}{n} \right)^{-r} \right), \quad (9)$$

$$\tilde{\varepsilon}_n(W_{\beta,1}^r)_{L_p} = \frac{1}{n^r} \left(\frac{2^{1-\frac{1}{p}}}{\pi^{1+\frac{1}{p}}} \|\cos t\|_p^2 + O(1) \left(\frac{1}{n} + \left(1 + \frac{1}{n} \right)^{-r} \right) \right), \quad (10)$$

where $1/p + 1/p' = 1$ and $O(1)$ are quantities uniformly bounded in all analyzed parameters. The estimates (9) and (10) are the asymptotic equalities, as $r/n \rightarrow \infty$, $n \rightarrow \infty$.

Comparing formulas (6), (7), (9) and (10) we see that

$$\begin{aligned} \tilde{\varepsilon}_n(W_{\beta,p}^r)_C &\sim 2\varepsilon_n(W_{\beta,p}^r)_C, \\ \tilde{\varepsilon}_n(W_{\beta,1}^r)_{L_p} &\sim \frac{2^{1-\frac{1}{p}}}{\pi^{\frac{1}{p}}} \|\cos t\|_p \varepsilon_n(W_{\beta,1}^r)_{L_p}, \end{aligned}$$

as $r/n \rightarrow \infty$, $n \rightarrow \infty$.

REFERENCES

- [1] A.I. Stepanets. *Methods of Approximation Theory*, Utrecht, VSP, 2005.
- [2] S. B. Stechkin. An estimation of the remainders of the Fourier series of differentiable functions. *Tr. Mat. Inst. Akad. Nauk SSSR*, 145, 126–151, 1980.
- [3] A.S. Serdyuk, I.V. Sokolenko. Approximation by interpolation trigonometric polynomials in metrics of the spaces L_p on the classes of periodic integer functions, *Ukr. Mat. Zh.*, Vol. 71, no 2, 283 – 292, 2019.

ІНСТИТУТ МАТЕМАТИКИ НАН України

| | |
|--|-----------|
| Mokritskaya T. P., Tushev A. V. <i>On some fractal-based estimations of subsidence volume for various types of soils</i> | 39 |
| Mukhamadiev F. G. <i>The Shanin number and the predshanin number of N_{τ}^{φ}-kernel of a topological spaces</i> | 41 |
| Najmiddinov J. Sh. <i>The effectiveness of the use of computer programs in the teaching of mathematics in academic lyceums</i> | 42 |
| Obikhod T. <i>Gromov-Witten invariants and identification of the energy levels of solitonic states</i> | 43 |
| Ostrovska O., Yakymiv R. <i>On isometries satisfying deformed commutation relations</i> | 45 |
| Prishlyak A., Prus A. <i>Three-color graph of the Morse flow on a compact surface with boundary</i> | 46 |
| Pulemotov A. <i>The Ricci Iteration on Homogeneous Spheres</i> | 48 |
| Rmuš V. <i>The construction of squaring the circle</i> | 49 |
| Samokhvalov S. <i>Riemann-Klein antagonism and problem of energy in general relativity</i> | 51 |
| Savchenko A. <i>On generalized spaces of persistence diagrams</i> | 52 |
| Sazonova O. <i>Continual approximate solution with acceleration and condensation mode</i> | 53 |
| Serdyuk A. S., Sokolenko I. V. <i>Approximation by Fourier sums and interpolation trigonometric polynomials in classes of differentiable functions with high exponents of smoothness</i> | 54 |
| Serdyuk A., Stepanyuk T. <i>Lebesgue-type inequalities for the Fourier sums</i> | 57 |
| Skuratovskii R. <i>Minimal generating set and structure of wreath product of cyclic groups, comutator of wreath product and the fundamental group of orbit Morse function $\pi_1 O(f)$</i> | 59 |
| Vasilchenko A. <i>Spaces of primitive elements in dual modules over Steenrod algebra 2</i> | 61 |
| Morrison P. J. <i>A Geometrical Version of the Maxwell-Vlasov Hamiltonian Structure</i> | 63 |
| Wojtowicz M. <i>Note on congruent numbers</i> | 64 |
| Кадубовський О. А. <i>Про число топологічно нееквівалентних гладких функцій з однією критичною точкою типу сідла на двовимірному торі</i> | 65 |
| Ладиненко Л. П. <i>Щодо геометричної характеристики спеціальних майже геодезичних перетворень просторів афінного зв'язку зі скрутом</i> | 67 |
| Овчаренко О. О. <i>Життєвий та науковий шлях Марка Григоровича Крейна</i> | 68 |
| Подоусова Т. Ю., Вашпанова Н. В. <i>LGT-лінії та A-деформації мінімальних поверхонь</i> | 69 |
| Прокіп В. М. <i>Алгоритм побудови унітального дільника для многочленної матриці</i> | 70 |
| Синюкова О. <i>Про геодезичні відображення просторів дотичних розшарувань зі спеціальною метрикою</i> | 72 |
| Щеглов М. В. <i>Поточкова оцінка відхилення полінома Крякіна від неперервної на відрізку функції</i> | 73 |