

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

What points in Goursat Monster Tower are strongly nilpotent

Piotr Mormul

(Institute of Mathematics, University of Warsaw, Warsaw, Poland)

E-mail: mormul@mimuw.edu.pl

Control systems linear in controls, with linearly independent vector field' generators, sometimes happen to be locally nilpotentizable. That is, to locally possess bases that generate (over reals, not over functions) nilpotent algebras of vector fields. The existence of a nilpotent basis may be somehow mischievously hidden in the nature of a system. When it exists and is at hand, a number of key control problems related with the system (e. g., motion planning) become much simpler. We call such systems *weakly* nilpotent. When a system Σ is given globally on a manifold M , we call weakly nilpotent those points in M , around which Σ is weakly nilpotent.

In turn, *strongly* nilpotent are those points p in M , around which Σ is equivalent to its *nilpotent approximation* at p . Naturally, 'strongly' implies 'weakly', but not vice versa: 'strongly' appears to be a much more stringent property.

An important class of weakly nilpotentizable distributions are *Goursat* distributions – members of Goursat *flags* which live on so-called Monster Manifolds ([1]). Local nilpotent bases found for Goursat distributions permit much more – to compute the *nilpotency orders* (sometimes also called 'indices', sometimes 'steps') of the generated real Lie algebras, see [2]. Those Lie algebras are sometimes called 'Kumpera-Ruiz' after the names of the discoverers of Goursat's singularities as such. A big problem, with only partial answers known to-date, reads

Problem 1. What points in the Goursat Monster Tower are strongly nilpotent ?

Problem 2. What are the dimensions of the Kumpera-Ruiz algebras, hidden in the Goursat distributions ?

(The nilpotency orders of the Kumpera-Ruiz algebras are tractable, but not their real dimensions.)

REFERENCES

- [1] Richard Montgomery & Michail Zhitomirskii. Points and Curves in the Monster Tower. *Memoirs of the American Mathematical Society* **956** (2010).
- [2] Piotr Mormul. Goursat distributions not strongly nilpotent in dimensions not exceeding seven. Volume **281** (2003) of *Lecture Notes in Control and Information Sciences*, 249 – 261.

V. Lomadze <i>A purely algebraic construction of Schwartz distributions</i>	34
S. Maksymenko <i>Deformation of functions on orientable surfaces by symplectic diffeomorphisms</i>	36
Martseniuk V., Bondarenko A., Gefter S. <i>Application the p-adic topology on Z for study determinants of infinite order with integer coefficients</i>	37
K. Matsumoto <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds, II</i>	38
Mormul P. <i>What points in Goursat Monster Tower are strongly nilpotent</i>	39
Mukhamadiev F. G. <i>The local density and the local weak density of N_7^φ-kernel of a topological spaces</i>	40
Tetiana V. Obikhod <i>Topological invariants and microscopic quantum description</i>	41
A. Ozkan <i>On the misconceptions of 10th grade students about analytical geometry</i>	42
Parasyuk I. O. <i>Hyperbolic quasiperiodic motion of charged particle on 2-sphere</i>	43
E. Petrov <i>Representing trees of finite ultrametric spaces and weak similarities</i>	45
Prishlyak A., Hladysh B. <i>Functions with three critical points on closed non-oriented 3-manifolds</i>	47
V.M. Prokip <i>A note on similarity of matrices</i>	49
Romakina L.N. <i>Inversion with respect to an elliptic cycle of a hyperbolic plane of positive curvature</i>	51
Romaniv A. M. <i>On the relationship between the Smith normal forms of matrices and of their least common multiple</i>	53
Ryazanova O. <i>Estimates for the surface with given average curvature</i>	54
Samokhvalov S. E. <i>Kinematic renormalization of energy in the gravity</i>	55
Savchenko A. <i>Fuzzy metrization of the spaces of idempotent measures</i>	56
Sazonova O. <i>General form of the Maxwellian distribution with arbitrary density</i>	57
Skuratovskii R. <i>The commutator and centralizer of Sylow subgroups of alternating and symmetric groups, its minimal generating set</i>	58
Sokhatsky F., Krainichuk H. <i>Semi-lattice of varieties of quasigroups with linearity</i>	59
Salimov R., Stefanchuk M. <i>Asymptotic behavior of solutions to a nonlinear Beltrami equation</i>	61
Mokritskaya T. P., Tushev A. V. <i>On fractal characteristics of the micro-aggregate structure of loess soils</i>	62
Vasilchenko A. N. <i>Spaces of primitive elements in dual modules over Steenrod algebra</i>	64