

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

Spaces of primitive elements in dual modules over Steenrod algebra

Alexander N. Vasilchenko

(19 Cheluskintsev, apt.32, Samara, phone +78463027287, +79372076904)

E-mail: vass-alexandr@yandex.ru

Presented work studies subspace of all primitives $PB(n)$ in $B(n) = (A(n-1)/A(n))^*$, with respect to the coaction $\phi_n^* : B(n) \rightarrow A^* \otimes B(n)$, where the primitives are those elements $\alpha \in B(n)$ for which $\phi_n^*(\alpha) = 1 \otimes \alpha$, here A^* is dual Steenrod algebra. Modules $B(n)$ were studied in [5, 6, 7]. Works [1, 2, 3, 4, 7] contain all useful notions and results: [3] studies structures of modules over Steenrod algebra A and their duals in dual Steenrod algebra A^* [2, 4]; [1] studies modules $A(n)$ over A generated by annihilators of cohomology classes with degrees no greater then n and $A(n)^*$ is the corresponding dual module of $A(n)$; In [5, 7], $A(n)^+$ is defined as annihilators of cohomology operations with excess greater then n , proved that as vector space over Z_p it is generated by all monomials of A^* with multiplicity no greater then n and $A(n)^+$ can be considered both left and right comodule over A^* . The filtration described in [6] Theorem 1 property 2 and 3 yields $PB(n) = \bigcup_t PB(n)_t$ and $PB(n)_t = \bigoplus_s PB(n)_t^s$, where s is the number of τ operations and where t is the biggest index of such operations. By property 4 Theorem 1 follows: $P(B(n)_t/B(n)_{t-1}) \simeq PB(n-1)_{t-1}$. Found properties of $PB(n)_t$ can be shortly summarized here:

Theorem 1 (Properties of $PB(n)_t$ spaces). (1) $PB(n) = \bigcup_t PB(n)_t$ and the following diagram is commutative with exact rows and columns:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & PB(n)_{t-1} & \xrightarrow{i_{t-1}} & PB(n)_t & \xrightarrow{\pi_t} & PB(n-1)_{t-1} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & B(n)_{t-1} & \xrightarrow{i_{t-1}} & B(n)_t & \xrightarrow{\pi_t} & B(n-1)_{t-1} \longrightarrow 0 \\
 & & \downarrow \phi_n & & \downarrow \phi_n & & \downarrow \phi_{n-1} \\
 0 & \longrightarrow & A^* \otimes B(n)_{t-1} & \xrightarrow{Id \otimes i_{t-1}} & A^* \otimes B(n)_t & \xrightarrow{Id \otimes \pi_t} & A^* \otimes B(n-1)_{t-1} \longrightarrow 0
 \end{array}$$

Here all denoted maps are induced by maps defined in Theorem 1 property 4 [6].

(2) For even n : $PB(n)_{-1} = PB(n)_0 = \langle \xi_1^{n/2} \rangle$; for odd n : $PB(n)_0 = \langle \xi_1^{\frac{n-1}{2}} \tau_0 \rangle$.

(3) $PB(n)_t = \bigoplus_s PB(n)_t^s$ and the following diagram is commutative with exact rows and columns:

$$\begin{array}{ccccccc}
 PB(n)_{t-1}^s & \xrightarrow{i_{t-1}^s} & PB(n)_t^s & \xrightarrow{\pi_t^s} & PB(n-1)_{t-1}^{s-1} \\
 \downarrow & & \downarrow & & \downarrow \\
 B(n)_{t-1}^s & \xrightarrow{i_{t-1}^s} & B(n)_t^s & \xrightarrow{\pi_t^s} & B(n-1)_{t-1}^{s-1} \\
 \downarrow \phi_n & & \downarrow \phi_n & & \downarrow \phi_{n-1} \\
 A^* \otimes B(n)_{t-1}^s & \xrightarrow{Id \otimes i_{t-1}^s} & A^* \otimes B(n)_t^s & \xrightarrow{Id \otimes \pi_t^s} & A^* \otimes B(n-1)_{t-1}^{s-1}
 \end{array}$$

Here all denoted maps are induced ones as in (1).

(4) Composition of maps

$$PB(n)_t^{s,deg} \longrightarrow B(n)^{s,deg} \twoheadrightarrow B(n)^{s,deg} / (I \cap B(n)^{s,deg})$$

is monomorphism, here $I = \langle \xi_2, \xi_3, \xi_4, \dots \rangle$ is ideal in A^* , where deg is degree. The dimension of $B(n)^{s,deg}/(I \cap B(n)^{s,deg})$ is less or equal 1.

- (5) $\dim(PB(n)_t^{s,deg}) \leq 1$.
(6) For map $PB(2m+1)_t \xrightarrow{\pi_t} PB(2m)_{t-1}$, $\xi_1^m \in \text{Im}(\pi_0)$ and for $t > 0$: $\xi_1^m \notin \text{Im}(\pi_t)$; if $s > 1$ and $\alpha \in PB(n)_t^s$ then $\alpha = \alpha' \tau_0$.
(7) Denote $J(n)^{s,deg} = B(n)^{s,deg}/(I \cap B(n)^{s,deg})$ then the following diagram is commutative:

$$\begin{array}{ccc}
PB(n)_t^{s,deg} & \xrightarrow{\pi_t^{s,deg}} & PB(n-1)_{t-1}^{s-1,deg-|\tau_t|} \\
\downarrow & & \downarrow \\
B(n)_t^{s,deg} & \xrightarrow{\pi_t^{s,deg}} & B(n-1)_{t-1}^{s-1,deg-|\tau_t|} \\
\downarrow j_n & & \downarrow j_{n-1} \\
J(n)^{s,deg} & & J(n-1)^{s-1,deg-|\tau_t|}
\end{array}$$

These properties may be useful to find bases of $PB(n)_t^{s,deg}$ spaces and $PB(n)$ basis as result.

REFERENCES

- [1] H. Cartan. Algebres d'Eilenberg-MacLane at Homotopie. *Seminare Cartan ENS*, 7e, 1954–1955.
[2] J. Milnor. The Steenrod algebra and its dual. *Annals of Mathematics*, 67 : 150–171, 1958.
[3] J. Milnor, J. Moore. On the structure of Hopf algebras. *Annals of Mathematics*, 81 : 211–264, 1965.
[4] N. Steenrod, D. B. A. Epstein. *Cohomological Operations*, Princeton University Press, 1962.
[5] А. Н. Васильченко. Свойства дуальных модулей над алгеброй Стиррода. Abstracts of the International Conference “Geometry in Odessa” – 2014, Odessa the 26th of May – the 31st of May 2014: p.26
[6] А. Н. Васильченко. Свойства дуальных модулей над алгеброй Стиррода. Abstracts of the International scientific conference “Algebraic and geometric methods of analysis” 2017, Odessa the 31st of May – the 6 June 2017: p.96
[7] А. Н. Васильченко. Свойство дуальных модулей над алгеброй Стиррода. *Вестник СамГУ* 7(118) : 9–16, 2014.

V. Lomadze <i>A purely algebraic construction of Schwartz distributions</i>	34
S. Maksymenko <i>Deformation of functions on orientable surfaces by symplectic diffeomorphisms</i>	36
Martseniuk V., Bondarenko A., Gefter S. <i>Application the p-adic topology on Z for study determinants of infinite order with integer coefficients</i>	37
K. Matsumoto <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds, II</i>	38
Mormul P. <i>What points in Goursat Monster Tower are strongly nilpotent</i>	39
Mukhamadiev F. G. <i>The local density and the local weak density of N_7^φ-kernel of a topological spaces</i>	40
Tetiana V. Obikhod <i>Topological invariants and microscopic quantum description</i>	41
A. Ozkan <i>On the misconceptions of 10th grade students about analytical geometry</i>	42
Parasyuk I. O. <i>Hyperbolic quasiperiodic motion of charged particle on 2-sphere</i>	43
E. Petrov <i>Representing trees of finite ultrametric spaces and weak similarities</i>	45
Prishlyak A., Hladysh B. <i>Functions with three critical points on closed non-oriented 3-manifolds</i>	47
V.M. Prokip <i>A note on similarity of matrices</i>	49
Romakina L.N. <i>Inversion with respect to an elliptic cycle of a hyperbolic plane of positive curvature</i>	51
Romaniv A. M. <i>On the relationship between the Smith normal forms of matrices and of their least common multiple</i>	53
Ryazanova O. <i>Estimates for the surface with given average curvature</i>	54
Samokhvalov S. E. <i>Kinematic renormalization of energy in the gravity</i>	55
Savchenko A. <i>Fuzzy metrization of the spaces of idempotent measures</i>	56
Sazonova O. <i>General form of the Maxwellian distribution with arbitrary density</i>	57
Skuratovskii R. <i>The commutator and centralizer of Sylow subgroups of alternating and symmetric groups, its minimal generating set</i>	58
Sokhatsky F., Krainichuk H. <i>Semi-lattice of varieties of quasigroups with linearity</i>	59
Salimov R., Stefanchuk M. <i>Asymptotic behavior of solutions to a nonlinear Beltrami equation</i>	61
Mokritskaya T. P., Tushev A. V. <i>On fractal characteristics of the micro-aggregate structure of loess soils</i>	62
Vasilchenko A. N. <i>Spaces of primitive elements in dual modules over Steenrod algebra</i>	64