

International
Scientific Conference



Algebraic
and Geometric
Methods
of Analysis

27-30 May 2024
Odesa, Ukraine

The purpose of this conference is to bring together researchers in geometry, topology, algebra, analysis and dynamical systems and to provide for them a forum to present their recent work to colleagues from different nationalities. This way we aim to stimulate discussion about the latest findings in geometrical and topological methods in analysis and to increase international collaboration.

The conference continues the traditional annual conference «Geometry in Odesa» holding from 2004, and hosted by Odesa National University of Technology (Odesa National Academy of Food Technologies till 2021). From 2017 the conference was renamed to «Algebraic and geometric methods of analysis» (AGMA).

The Conference languages: Ukrainian and English.

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology, Ukraine
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

- | | |
|---|--|
| • Vladimir Balan (<i>Bucharest, Romania</i>) | • Volodymyr Lyubashenko (<i>Kyiv, Ukraine</i>) |
| • Taras Banakh (<i>Lviv, Ukraine</i>) | • Sergiy Maksymenko (<i>Kyiv, Ukraine</i>) |
| • Dmytro Bolotov (<i>Kharkiv, Ukraine</i>) | • Koji Matsumoto (<i>Yamagata, Japan</i>) |
| • Vyacheslav Boyko (<i>Kyiv, Ukraine</i>) | • Piotr Mormul (<i>Warsaw, Poland</i>) |
| • Yulia Fedchenko (<i>Odesa, Ukraine</i>) | • Maryna Nesterenko (<i>Kyiv, Ukraine</i>) |
| • Oleg Gutik (<i>Lviv, Ukraine</i>) | • Roman Popovych (<i>Kyiv, Ukraine</i>) |
| • Olena Karlova (<i>Chernivtsi, Ukraine</i>) | • Alexandr Prishlyak (<i>Kyiv, Ukraine</i>) |
| • Volodymyr Kiosak (<i>Odesa, Ukraine</i>) | • Aleksandr Savchenko (<i>Kherson, Ukraine</i>) |
| • Nadiia Konovenko (<i>Odesa, Ukraine</i>) | |

ORGANIZING COMMITTEE

- | | |
|---|--|
| • Nadiia Konovenko (<i>Odesa, Ukraine</i>) | • Bohdan Mazhar (<i>Kyiv, Ukraine</i>) |
| • Yuliya Fedchenko (<i>Odesa, Ukraine</i>) | • Sergiy Maksymenko (<i>Kyiv, Ukraine</i>) |
| • Mykola Lysynskiy (<i>Kyiv, Ukraine</i>) | • Alexandr Prishlyak (<i>Kyiv, Ukraine</i>) |

On Beloch's curve that appears when solving real cubic with origami

Manami Nijima

(JVCKENWOOD Corporation, Japan)

E-mail: sakura5.2ijima@gmail.com

Origami construction is generally defined by 7 Axioms (re-)founded by Justin–Huzita–Hatori in 1980's (cf. [Jus86, p.40–45], [Lan10],[Ned22]). Among other things, Axiom 6 allows so-called “neusis”, which is not allowed in the straightedge and compass construction.

1 (Justin–Huzita–Hatori Axioms 6). *Given two points A and B and two lines l_1 and l_2 on a plane, there exists a fold that places A onto l_1 and B onto l_2 at the same time. In other words, it is possible to construct a certain fold line l such that points symmetrical to points A and B with respect to l are placed on the straight lines l_1 and l_2 , respectively.*

This Axiom 6 enables to construct all solutions of any given real cubic equation by using a perfect piece of origami. The following method was initially shown by Beloch in 1936 [Bel36], based on Lill's enjoyable idea [Lil67] (see also [Kat99], [Hul11], [NO15]).


Theorem 2. *Given segments with length $a, b, c \in \mathbb{R}$, every solution of the cubic formula*

$$x^3 - ax^2 - bx + c = 0 \cdots (*)$$

may be constructed. More precisely,

(1) *If we make a fold that places (I) $A(-1, 0)$ onto $x = 1$ and (II) $P(b, a + c)$ onto $y = a - c$, then the y -intercept r of the fold line l is a solution of $(*)$*

(2) *For any solution $x = r$ of $(*)$, a fold that places $A(-1, 0)$ onto $A'(1, 2r)$ satisfies the conditions (I) and (II) in (1).*



In other words, there is a bijective correspondence between all fold lines satisfying the conditions (I) and (II) and all real solutions of $(*)$. In addition, by considering all fold lines satisfying (I) and parametrizing them by r , we may find all solutions.

We consider all fold lines satisfying the condition (I), and investigate the orbit of the points $P'(r)$ ($r \in \mathbb{R}$), each of which is symmetric to P with respect to a fold line. We set $(p, q) = (b, a + c)$. Our main results may be summarized into the following theorem.


Theorem 3. *The union \mathcal{F} of the orbit of the points P' and the point $P(p, q)$ is a real cubic curve*

$$F(x, y) := 2(q - y)^2 - (q + y)(q - y)(p - x) - (p - x)^2(p + x) = 0,$$

which we call Beloch's curve.

The point P is its uniquely existing singular point, and the Hessian at P is given by $\mathcal{H}_{\mathcal{F}} = -4(4p + q^2)$. We have the following equivalence on the shape of \mathcal{F} and the parabola $\mathcal{G} : 4x + y^2 = 0$.

- P is on the left side of $\mathcal{G} \iff$ The orbit of P' 's does not pass through $P \iff P$ is an isolated point of \mathcal{F} .
- P is on $\mathcal{G} \iff$ The orbit of P' 's passes through P just once $\iff P$ is a cusp of \mathcal{F} .
- P is on the right side of $\mathcal{G} \iff$ The orbit of P' 's passes through P twice $\iff P$ is a self-intersection point of \mathcal{F} .



We may also classify the shapes of real cubic curves $a_0y^2 - a_1xy^2 - a_3x^2 - a_4x^3 = 0$ in a similar manner. In addition, we show that the rotation number of \mathcal{F} around the point A is determined by the relationship between P and $x = 1$.

In the proof of 3, we use the fact that the fold line l is the tangent line of the parabola \mathcal{G} at $(-r^2, 2r)$. In general, given a point A , Q , and a line m , then Axiom 5 allows a fold l such that $Q \in l$ and l maps A onto m . 3 would shed new light on the relationship between Axioms 5 and 6.

REFERENCES

- [Bel36] Margherita Pianolla Beloch. Sul metodo del ripiegamento della carta per la risoluzione dei problemi geometrici. *Periodico di Mat.* IV, XVI, (2):104–108, 1936.
- [Hul11] Thomas C. Hull. Solving cubics with creases: the work of Beloch and Lill. *Am. Math. Mon.*, 118(4):307–315, 2011.
- [Jus86] Jacques Justin. Résolution par le pliage de l'équation du troisième degré et applications géométriques. *L'Ouvret - Journal de l'APMEP d'Alsace et de l'IREM de Strasbourg*, 42:9–19, 1986.
- [Kat99] Koichi Kato. Origami and formulas. 1999.
- [Lan10] Robert J. Lang. Origami and geometric constructions. 2010.
- [Lil67] E. Lill. Résolution graphique des équations numériques de tous les degrés à une seule inconnue, et description d'un instrument inventé dans ce but. *Nouvelles annales de mathématiques : journal des candidats aux écoles polytechnique et normale*, 6:359–362, 1867.
- [Ned22] Dmitri Nedrenco. Axiomatisieren lernen mit papierfalten. *JMU Würzburg, 2022. PhD Thesis. Available online (urn:nbn:de:bvb:20-opus-279383)*., 2022.
- [NO15] Isao Nakai and Eri Ono. Finding roots by a simple origami. *Nat. Sci. Rep. Ochanomizu Univ.*, 66(1):1–17, 2015.

G. Kuduk <i>Integral problem for system of partial differential equations of third order</i>	65
A. Kuramoto <i>The density of Borromean primes</i>	66
I. Kurbatova, N. Konovenko, M. Pistruil <i>Invariant transformation of generalized-recurrent-parabolic spaces that are in a quasi-geodesic mapping</i>	68
J. Lang <i>Notes on the Quality of Non-compactness for Non-compact Sobolev Embeddings</i>	70
D. Lehmann <i>Ordinary linear differential operators and connections. Application to curvilinear webs</i>	70
Jian Liu, Dong Chen, and Guo-Wei Wei <i>Persistent interaction topology in data analysis</i>	73
L. Lotarets <i>Reeb vector field as isometric embedding</i>	73
S. Maksymenko, M. Lysynskiy <i>Classification of smooth structures on line with two origins</i>	75
D. Maingi <i>Vector bundle construction via monads on multiprojective spaces</i>	77
O. Makarchuk <i>About one problem of the Gauss-Kuzmin type</i>	78
S. Maksymenko <i>Homotopy types of stabilizers of Morse-Bott functions on 3-manifolds</i>	79
M. Jinzenji, K. Kuwata <i>Elliptic Virtual Structure Constants and Generalizations of BCOV-Zinger Formula to Projective Fano Hypersurfaces</i>	80
B. Mazhar, S. Maksymenko <i>Deformation properties of smooth functions on Klein bottle</i>	80
Ł. Michalak <i>Algebraic periods of surface homeomorphisms</i>	82
H. Monaim <i>Wigner-Ville distribution associated with quadratic Clifford-Fourier transform</i>	82
P. Mormul <i>Non-simple strongly nilpotent distribution germs</i>	82
V. Mykhaylyuk <i>Extending of partial metrics</i>	83
C. L. Nehaniv <i>Axiomatic Development of Complexity Theory for Finite Groups</i>	84
M. Nesterenko <i>Construction and application of quasicrystals</i>	85
M. Nijjima <i>On Beloch's curve that appears when solving real cubic with origami</i>	87
Z. Novosad, A. Zagorodnyuk <i>Hypercyclicity of symmetric composition operator</i>	89
M. Nxumalo <i>On (i, j)-Baire Bilocales</i>	89
T. Obikhod <i>Application of the dynamical system theory for counting black hole entropy of microstates</i>	90