



International  
Scientific Conference



# Algebraic and Geometric Methods of Analysis



Devoted to 160 anniversary of  
**Dvytro Grave**  
(25.08.1863 - 19.12.1939)  
Academician of the Ukrainian  
Academy of Sciences, the  
first director of the Institute of  
Mathematics of NAS of Ukraine

May 29 – June 1, 2023  
Odesa, Ukraine

## LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

## ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

## SCIENTIFIC COMMITTEE

- |  |   |
|--|---|
| • <b>Bolotov D.</b> ( <i>Kharkiv, Ukraine</i> )  | • <b>Konovenko N.</b> ( <i>Odesa, Ukraine</i> )   |
| • <b>Bondarenko V.</b> ( <i>Kyiv, Ukraine</i> )  | • <b>Maksymenko S.</b> ( <i>Kyiv, Ukraine</i> )   |
| • <b>Boychuk O.</b> ( <i>Kyiv, Ukraine</i> )     | • <b>Mikhailets V.</b> ( <i>Kyiv, Ukraine</i> )   |
| • <b>Boyko V.</b> ( <i>Kyiv, Ukraine</i> )       | • <b>Ostrovskiy V.</b> ( <i>Kyiv, Ukraine</i> )   |
| • <b>Cherevko Ye.</b> ( <i>Odesa, Ukraine</i> )  | • <b>Petravchuk A.</b> ( <i>Kyiv, Ukraine</i> )   |
| • <b>Dorogovtsev A.</b> ( <i>Kyiv, Ukraine</i> ) | • <b>Plaksa S.</b> ( <i>Kyiv, Ukraine</i> )       |
| • <b>Drozd Yu.</b> ( <i>Kyiv, Ukraine</i> )      | • <b>Portenko M.</b> ( <i>Kyiv, Ukraine</i> )     |
| • <b>Gerasymenko V.</b> ( <i>Kyiv, Ukraine</i> ) | • <b>Pratsiovytyi M.</b> ( <i>Kyiv, Ukraine</i> ) |
| • <b>Fedchenko Yu.</b> ( <i>Odesa, Ukraine</i> ) | • <b>Savchenko O.</b> ( <i>Kherson, Ukraine</i> ) |
| • <b>Kiosak V.</b> ( <i>Odesa, Ukraine</i> )     | • <b>Romanyuk A.</b> ( <i>Kyiv, Ukraine</i> )     |
| • <b>Kochubei A.</b> ( <i>Kyiv, Ukraine</i> )    | • <b>Timokha O.</b> ( <i>Kyiv, Ukraine</i> )      |

## ORGANIZING COMMITTEE

- |  |   |
|--|---|
| • <b>Maksymenko S.</b> ( <i>Kyiv, Ukraine</i> )  | • <b>Cherevko Ye.</b> ( <i>Odesa, Ukraine</i> ) |
| • <b>Konovenko N.</b> ( <i>Odesa, Ukraine</i> )  | • <b>Osadchuk Ye.</b> ( <i>Odesa, Ukraine</i> ) |
| • <b>Fedchenko Yu.</b> ( <i>Odesa, Ukraine</i> ) | • <b>Sergeeva O.</b> ( <i>Odesa, Ukraine</i> )  |

- [11] C. B. Rayner. *The exponential map for the Lagrange problem on differentiable manifolds*, Philosophical Transactions of the Royal Society of London, Series A, Mathematical and Physical Sciences Vol. 262, No. 1127 (1967), pp. 299-344.
- [12] L. Rifford. *Sub-Riemannian geometry and optimal transport*, Springer, (2014).
- [13] R. Strichartz. *Sub-Riemannian geometry*. J. Differ. Geom. **24** (1986), 221-263; correction ibid. **30** (1989), 595-596.

## Geometric properties of interception curves

Yagub Aliyev

(School of IT and Engineering, ADA University, Ahmadbey Aghaoglu str. 61, Baku AZ1008, Azerbaijan)

*E-mail*: y.aliyev@ada.edu.az

In this study, a plane curve, which was named as *Interception Curve*, was discussed. This curve can be defined in the following way. Suppose one point moves with constant velocity along a straight line, and another point, at the beginning one unit apart from the line and the first point on this line, moves with the same constant speed so that it always stays on a line passing through the first point and the initial position of the second point. This plane curve appears in problems related to the interception of high-speed targets by beam rider missiles (hence the name *Interception Curve*) [2, 5]. This curve was also mentioned in [4, 6, 1]. In [3], at Sect. 1.460 and Sect. 1.507, some methods based on polar and Cartesian coordinates were proposed to find an explicit representation for this curve.

**Problem 1.** If two points  $P(x, y)$  and  $Q$ , initially at  $O(0, 0)$  and  $A(1, 0)$ , respectively, move uniformly so that  $Q$  is on the line  $x = 1$ , and  $P$  is on the ray  $OQ$  then what curve does the point  $P$  draw?

**Answer.** Let us use polar coordinates  $r = |OP|$  and  $\angle AOQ = \theta$ . We obtain ordinary differential equation

$$r(\theta)^2 + (r'(\theta))^2 = \frac{1}{\cos^4 \theta}, \quad (1)$$

with initial condition  $r(0) = 0$ . Note that in the cartesian coordinates, (1) can be written as

$$x^2 \sqrt{1 + (y'(x))^2} = y'x - y, \quad (2)$$

with initial condition  $y(0) = 0$ . By solving this equation, we obtain the parametrization (cf. [3], Sect. 1.507, where the roles of  $x$  and  $y$  are interchanged)

$$\begin{cases} x(p) = \frac{1}{\sqrt{p}} \int_1^p \frac{\sqrt{tdt}}{2\sqrt{t^2-1}}, \\ y(p) = \frac{\sqrt{p^2-1}}{\sqrt{p}} \int_1^p \frac{\sqrt{tdt}}{2\sqrt{t^2-1}} - \left( \int_1^p \frac{\sqrt{tdt}}{2\sqrt{t^2-1}} \right)^2 \end{cases} \quad (p \geq 1). \quad (3)$$

Using all these, the following results are obtained:

**Theorem 2.** Suppose that  $U$  is the  $y$  intercept of the tangent line of the curve (3) at the point  $P$ , and this tangent line intersects the line  $x = 1$  at point and  $T$ . Then

- (1)  $x \cdot |UP| = |OU|$ ,
- (2)  $\sin \angle QPT = \frac{x^2}{|OP|} = \frac{x}{|OQ|}$ ,

where  $x$  is the abscissa of the point  $P(x, y)$ .

**Theorem 3.** Consider intersection point  $M$  of mid-perpendicular of  $OP$  and the line perpendicular to  $UT$  at the point  $P$ . Similarly, consider intersection point  $N$  of mid-perpendicular of  $OQ$  and the line perpendicular to  $QT$  at the point  $Q$ . Then the points  $M$  and  $N$  are equidistant from the point  $O$  i.e.  $MO=NO$ .

The following result shows that there is a connection between the interception curve and Gauss's constant  $G$  defined by the arithmetic-geometric mean.

**Theorem 4.**

$$\lim_{x \rightarrow 1^-} |PQ| = \frac{1}{4G^2}.$$

**Problem 5.** Suppose that two points  $P$  and  $Q$ , at the beginning at  $B(0,0,1)$  and  $A(1,0,0)$ , respectively, move uniformly so that  $Q$  is on the equator  $z = 0$ ,  $x^2 + y^2 = 1$  of sphere  $x^2 + y^2 + z^2 = 1$  with center  $O(0,0,0)$ , and  $P$  is on the meridian through  $B$  and  $Q$  of the sphere. What curve does the point  $P$  draw?

**Answer.** We can use spherical coordinates to describe this curve:  $\angle AOQ = \theta$  and  $\angle POB = \phi$ . Since  $\rho = |OP| = 1$ , for the coordinates of point  $P(x, y, z)$ , we can write  $x = \cos \theta \sin \phi$ ,  $y = \sin \theta \sin \phi$ , and  $z = \cos \phi$ , where we think of  $\phi = \phi(\theta)$  as a function of  $\theta$ . For this curve we obtain

$$\phi = \tan^{-1} \sinh \theta. \quad (4)$$

Note that (4), which can also be expressed as  $\sin \phi = \tanh \theta$ , is sometimes called Gudermanian function  $\text{gd}(x)$ . For the curve defined by (4) the following results are obtained.

**Theorem 6.**  $\lim_{\theta \rightarrow \infty} |PQ| = 0$ .

In the following, we will use notation  $\widehat{XY}$  for the spherical distance between points  $X$  and  $Y$  on a sphere. Of course, for a unit sphere with center  $O$ ,  $\widehat{XY} = \angle XOY$ .

**Theorem 7.** If a great circle is tangent to the curve (4) at point  $P$ , intersects the equator at point  $T$ , then

- (1)  $\widehat{PT} = \frac{\pi}{2} - \widehat{TQ}$ ,
- (2)  $\widehat{TQ} < \widehat{PT}$ , and  $\lim_{\theta \rightarrow \infty} \widehat{TQ} = \lim_{\theta \rightarrow \infty} \widehat{PT} = \frac{\pi}{4}$ .
- (3)  $\angle BPT = \pi - \widehat{BP}$ .

**Theorem 8.** If a small spherical circle through point  $B$  is tangent to the curve (4) at point  $P$ , then its spherical radius  $R$  satisfies  $\tan R = \frac{1}{2} \sec^2 \frac{1}{2} \widehat{BP}$ .

We can prove some of these results also using simpler plane and spherical geometry methods, which are interesting on their own. It can be shown that the results agree with the angle-preserving property of Mercator and Stereographic projections. The Mercator and Stereographic projections also reveal the symmetry of this curve with respect to Spherical and Logarithmic Spirals.

#### REFERENCES

- [1] Bailey, H.R. The Hiding Path. *Math. Mag.* **1994**, *67*, 40–44.
- [2] Elnan, O.R.S.; Lo, H. Interception of High-Speed Target by Beam Rider Missile. *Aiaa J.* **1963**, *1*, 1637–1639.
- [3] Kamke, E. *Differentialgleichungen Lösungsmethoden und Lösungen*; Springer, Wiesbaden, Germany, 1977.
- [4] Wilder, C.E. A discussion of a differential equation. *Am. Math. Mon.* **1931**, *38*, 17–25.

- [5] Vinh, N.X. Comment on "Interception of High-Speed Target by Beam Rider Missile". *AIAA J.* **1964**, *2*, 409.  
 [6] Zbornik, J. Akademie der Wissenschaften in Wien Mathematisch-Naturwissenschaftliche Klasse. *Ila Sitzungsberichte* **1957**, *166*, 42.

## Planar and non-planar degenerations with related fundamental groups

Meirav Amram  
(SCE)

*E-mail:* meiravt@sce.ac.il

We study planar and non-planar degenerations that are related to algebraic surfaces. It is interesting to see the differences in results and research methods between both cases. We have studied already planar degenerations with an  $R_k$  singularity, non-planar degenerations of degree 4, 6, and 8. The fundamental groups of the Galois covers of the related surfaces were investigated, because those groups are invariants of classification of algebraic surfaces in the moduli space.

**Theorem 1.** *The fundamental groups of surfaces that degenerate to one  $R_k$  singularity are all trivial, for any  $k$ .*




FIGURE 1.1.  $R_k$  singularity

**Theorem 2.** *The fundamental groups of Galois covers related to non-planar degenerations are trivial (for a degree 4 degeneration),  $\mathbb{Z}_2^4$  (for a degree 6 degeneration), and a metabelian group of order  $2^{23}$  (for a degree 8 degeneration).*




FIGURE 2.2. Degree 4 non-planar degeneration

## Table of contents

<b>L. M. Alabdulsada, L. Kozma</b> <i>Hopf-Rinow theorem of sub-Finslerian geometry</i>	<b>2</b>
<b>Y. Aliyev</b> <i>Geometric properties of interception curves</i>	<b>3</b>
<b>M. Amram</b> <i>Planar and non-planar degenerations with related fundamental groups</i>	<b>5</b>
<b>N. Ando</b> <i>Surfaces with zero mean curvature vector in 4-dimensional spaces</i>	<b>6</b>
<b>B. Apanasov</b> <i>Dynamics in nilpotent groups and deformations of locally symmetric rank one manifolds</i>	<b>8</b>
<b>M. Atteya</b> <i>Characterizing Linear Mappings Through Unital Algebra</i>	<b>9</b>
<b>S. Sharma, V. K. Bhat</b> <i>Edge resolvability and topological characteristics of zero-divisor graphs</i>	<b>10</b>
<b>V. Bilet, O. Dovgoshey</b> <i>From minimality to maximality via metric reflection</i>	<b>12</b>
<b>D. Bolotov</b> <i>Thurston norm and Euler classes of bounded mean curvature foliations on hyperbolic 3-Manifolds</i>	<b>14</b>
<b>A. Bolsinov</b> <i>Nijenhuis geometry and its applications</i>	<b>15</b>
<b>E. Bonacci</b> <i>Shape optimization in the batch crystallization of CAM</i>	<b>17</b>
<b>F. Bulnes</b> <i>Homotopies to Diffeomorphisms in Symplectic Field Theory</i>	<b>18</b>
<b>D. Cheban</b> <i>Global asymptotic stability of generalized homogeneous dynamical systems</i>	<b>19</b>
<b>Y. Cherevko, V. Berezovski, J. Mikeš, Y. Fedchenko</b> <i>Hyper-holomorphically projective mappings of hyper-Kähler manifolds</i>	<b>21</b>
<b>S. Dann</b> <i>On a problem of Fejes Toth</i>	<b>22</b>
<b>M. Golasinski, T. de Melo, R. Bononi</b> <i>Gottlieb groups of some Moore spaces</i>	<b>23</b>
<b>A. Białyżyt, A. Denkowska, M. P. Denkowski</b> <i>Inner semi-continuity of medial axes and conflict sets</i>	<b>24</b>
<b>K. v. Dichter</b> <i>The diameter-width-ratio for complete and pseudo-complete sets</i>	<b>25</b>
<b>O. Dovhopiatyi</b> <i>On the possibility of joining two pairs of points in convex domains using paths</i>	<b>27</b>
<b>Yu. Drozd</b> <i>Backström curves</i>	<b>28</b>