



International
Scientific Conference



Algebraic and Geometric Methods of Analysis



Devoted to 160 anniversary of
Dvytro Grave
(25.08.1863 - 19.12.1939)
Academician of the Ukrainian
Academy of Sciences, the
first director of the Institute of
Mathematics of NAS of Ukraine

May 29 – June 1, 2023
Odesa, Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

- | | |
|--|---|
| • Bolotov D. (<i>Kharkiv, Ukraine</i>) | • Konovenko N. (<i>Odesa, Ukraine</i>) |
| • Bondarenko V. (<i>Kyiv, Ukraine</i>) | • Maksymenko S. (<i>Kyiv, Ukraine</i>) |
| • Boychuk O. (<i>Kyiv, Ukraine</i>) | • Mikhailets V. (<i>Kyiv, Ukraine</i>) |
| • Boyko V. (<i>Kyiv, Ukraine</i>) | • Ostrovskiy V. (<i>Kyiv, Ukraine</i>) |
| • Cherevko Ye. (<i>Odesa, Ukraine</i>) | • Petravchuk A. (<i>Kyiv, Ukraine</i>) |
| • Dorogovtsev A. (<i>Kyiv, Ukraine</i>) | • Plaksa S. (<i>Kyiv, Ukraine</i>) |
| • Drozd Yu. (<i>Kyiv, Ukraine</i>) | • Portenko M. (<i>Kyiv, Ukraine</i>) |
| • Gerasymenko V. (<i>Kyiv, Ukraine</i>) | • Pratsiovytyi M. (<i>Kyiv, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Savchenko O. (<i>Kherson, Ukraine</i>) |
| • Kiosak V. (<i>Odesa, Ukraine</i>) | • Romanyuk A. (<i>Kyiv, Ukraine</i>) |
| • Kochubei A. (<i>Kyiv, Ukraine</i>) | • Timokha O. (<i>Kyiv, Ukraine</i>) |

ORGANIZING COMMITTEE

- | | |
|--|---|
| • Maksymenko S. (<i>Kyiv, Ukraine</i>) | • Cherevko Ye. (<i>Odesa, Ukraine</i>) |
| • Konovenko N. (<i>Odesa, Ukraine</i>) | • Osadchuk Ye. (<i>Odesa, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Sergeeva O. (<i>Odesa, Ukraine</i>) |

where $C_0 = 2 \min\{inj(M), (\coth)^{-1}(H_0)\}$, $Vol(M)$ is the volume of M and $inj(M)$ is the injectivity radius of M .

REFERENCES

- [1] W.P. Thurston, A norm for the homology of 3-manifolds. *Memoirs of the American Mathematical Society*, 59 (339): 99–130, 1986.
- [2] D. Sullivan, A homological characterization of foliations consisting of minimal surfaces, *Comm. Math. Helv.*, 54: 218-223, 1979.

Nijenhuis geometry and its applications

Alexey Bolsinov

(Loughborough University, Loughborough LE11 3TU United Kingdom)

E-mail: a.bolsinov@lboro.ac.uk

This presentation is focused on some results of the long-term research programme *Nijenhuis Geometry* initiated several years ago in cooperation with Vladimir Matveev and Andrey Konyaev.

A *Nijenhuis operator* $L = (L_j^i(x))$ is defined to be a field of endomorphisms on a smooth manifold M such that its Nijenhuis torsion identically vanishes, i.e.,

$$\mathcal{N}_L(\xi, \eta) = L^2[\xi, \eta] + [L\xi, L\eta] - L[L\xi, \eta] - L[\xi, L\eta] = 0, \quad (1)$$

for arbitrary vector fields ξ, η on M . The pair (M, L) is called a *Nijenhuis manifold*.

Relation (1) is the simplest differential-geometric condition on a field of endomorphisms, and that is the reason why Nijenhuis operators appear in many areas of differential geometry and mathematical physics. In the theory of integrable bi-Hamiltonian systems, they serve as recursion operators and their role in this area has been well understood for many years due to pioneering works by F. Magri, Y. Kosmann-Schwarzbach and F. Turiel. A classical fact in complex geometry is that an almost complex structure is integrable if and only if it is Nijenhuis (Newlander–Nireberg theorem). In the context of metric projective geometry, Nijenhuis operators played a crucial role in various classification problems (AB and V. Matveev). They naturally occur in the study of infinite dimensional Poisson brackets of hydrodynamic type (E. Ferapontov *et al*). Even in algebra, Nijenhuis operators turns out to be useful in the theory of integrable systems on Lie algebras and Lie pencils (A. Panasyuk), and also appear as left symmetric algebras.

Besides various applications, our motivation is as follows. Classical geometries are defined by means of a tensor of order 2. For Riemannian, sub-Riemannian, symplectic and Poisson structures, this tensor is a bilinear form (co- or contravariant, symmetric or skew-symmetric). In this list, one type of tensors is still missing: linear operators. Nijenhuis geometry would be a very natural candidate to fill this gap.

Thus, *Nijenhuis Geometry* research programme is aimed at systematic development of the theory of Nijenhuis manifolds. Our vision and first results are presented in [1–8]. More specifically, our goal is to *re-direct the research agenda* in this area from *tensor analysis at generic points* to studying *singularities and global properties*. The ultimate goal of our research programme is to answer three fundamental questions:

- (A) **Local description:** to what form can one bring a Nijenhuis operator near almost every point by a local coordinate change?

- (B) **Singular points:** what does it mean for a point to be generic or singular in the context of Nijenhuis geometry? What singularities are non-degenerate/stable? How do Nijenhuis operators behave near non-degenerate and stable singular points?
- (C) **Global properties:** what restrictions on a Nijenhuis operator are imposed by the topology of the underlying manifold? And conversely, what are topological obstructions to a Nijenhuis manifold carrying a Nijenhuis operator with specific properties?

Below are some of our easy-to-formulate results in the area.

Theorem 1. *Let L be a Nijenhuis operator and $\sigma_1, \dots, \sigma_n$ be the coefficients of its characteristic polynomial $\chi(t) = \det(t \cdot \text{Id} - L) = t^n - \sum_{k=1}^n \sigma_k t^{n-k}$. Then in any local coordinate system x_1, \dots, x_n the following matrix relation hold:*

$$J(x) L(x) = S_\chi(x) J(x), \quad \text{where } S_\chi(x) = \begin{pmatrix} \sigma_1(x) & 1 & & \\ \vdots & 0 & \ddots & \\ \sigma_{n-1}(x) & \vdots & \ddots & 1 \\ \sigma_n(x) & 0 & \dots & 0 \end{pmatrix} \quad (2)$$

and $J(x)$ is the Jacobi matrix of the collection of functions $\sigma_1, \dots, \sigma_n$ w.r.t. the variables x_1, \dots, x_n .

Theorem 2. *Let L be a real-analytic Nijenhuis operator of the form*

$$L(x) = L_{\text{lin}}(x) + R(x), \quad \text{where } L_{\text{lin}}(x) = \text{diag}(x_1, x_2, \dots, x_n)$$

and $R(x)$ denotes a non-linear perturbation (of order ≥ 2). Then $L(x)$ is linearisable, i.e., there exists a real analytic change of variables $x \mapsto y$ such that in the new coordinates $L(y) = \text{diag}(y_1, y_2, \dots, y_n)$.

Theorem 3. *A Nijenhuis operator on a closed connected manifold cannot have non-constant complex eigenvalues.*

Theorem 4. *Consider a real analytic gl-regular Nijenhuis operator L (gl-regularity means that each eigenvalue of L may have arbitrary multiplicity but only one linearly independent eigenvector). Then there exist local coordinate systems $u = (u^1, \dots, u^n)$ and $v = (v^1, \dots, v^n)$ in which L reduces to the first and second companion forms:*

$$L(u) = L_{\text{comp1}} = \begin{pmatrix} \sigma_1 & 1 & & \\ \vdots & 0 & \ddots & \\ \sigma_{n-1} & \vdots & \ddots & 1 \\ \sigma_n & 0 & \dots & 0 \end{pmatrix} \quad \text{and} \quad L(v) = L_{\text{comp2}} = \begin{pmatrix} 0 & 1 & & \\ \vdots & \ddots & \ddots & \\ 0 & \dots & 0 & 1 \\ \sigma_n & \sigma_{n-1} & \dots & \sigma_1 \end{pmatrix},$$

where σ_i are the coefficients of the characteristic polynomial of L in the corresponding coordinate system.

Theorem 5. *Let M^2 be either a sphere or a closed Riemann surface of genus ≥ 2 . Then M^2 cannot carry any gl-regular Nijenhuis operator L except for $L = \alpha \text{Id} + \beta A$, where A is a complex structure on M^2 and $\alpha, \beta \in \mathbb{R}$, $\beta \neq 0$. A non-orientable closed 2-manifold different from a Klein bottle cannot carry any gl-regular Nijenhuis operator.*

REFERENCES

- [1] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Nijenhuis geometry. *Advances in Mathematics*, 394: 108001, (2022).
- [2] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Nijenhuis geometry II: Left-symmetric algebras and linearization problem for Nijenhuis operators. *Differential Geometry and its Applications*, 74: 101706, 2021.
- [3] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Nijenhuis Geometry III: gl-regular Nijenhuis operators. *Revista Matemática Iberoamericana*, 2023, DOI 10.4171/RMI/1416.
- [4] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Nijenhuis Geometry IV: conservation laws, symmetries and integration of certain non-diagonalisable systems of hydrodynamic type in quadratures. *Preprint*, arXiv:2304.10626.
- [5] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Applications of Nijenhuis geometry: nondegenerate singular points of Poisson–Nijenhuis structures, *European Journal of Mathematics*, 8: 1355–1376, 2022.
- [6] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Applications of Nijenhuis geometry II: maximal pencils of multi-Hamiltonian structures of hydrodynamic type. *Nonlinearity*, 34(8): 5136–5162, 2021.
- [7] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Applications of Nijenhuis geometry III: Frobenius pencils and compatible non-homogeneous Poisson structures. *Journal of Geometric Analysis*, 2023, arXiv: 2112.09471v2.
- [8] Alexey Bolsinov, Andrey Konyaev, Vladimir Matveev, Applications of Nijenhuis Geometry IV: multicomponent KdV and Camassa-Holm equations. *Dynamics of PDE*, 20(1): 73–98, 2023.

Shape optimization in the batch crystallization of CAM

Enzo Bonacci

(The Mathematics & Statistics Unit of ATINER, Athens, Greece)

E-mail: enzo.bonacci@physics.org

The citric acid monohydrate (CAM) is an important organic substance but, until 1997, the scientific literature covered mostly the kinetics of nucleation [4] and the crystal growth [5] rather than its production via the crystallization by cooling in a stirred tank reactor (STR). The Department of Chemical Engineering at the University “La Sapienza” of Rome decided to fill that sci-tech gap through a meticulous investigation, with three STRs at the laboratories of San Pietro in Vincoli’s district, on the crystallization in discontinuous (batch) of CAM from aqueous solutions. The author participated in that cutting edge experience, as experimenter and coder under the supervision of Prof. Barbara Mazzarotta, in the years 1997-1998 [1]. Our specific tasks were to spot the main operating conditions, to modify them until an *optimal* crystal size distribution (CSD), i.e., large-sized homogeneous crystals of CAM, and to write a QBasic program predicting the outcomes of any test in batch reactors [2]. Here we focus on the influence of the STRs’ geometry, i.e., the role played by the tanks in crystallizing the CAM thanks to their differently shaped bottoms (flat, hemispherical, conical). All the data, collected and simulated, show that the round-bottomed crystallizer gives the best CSD, performing better than the conical-bottomed STR, and that we should discard the flat-bottomed STR for the poor quality of its crystalline product [3]. The homogenous distribution of large crystals from the round-bottomed STR is due to the *optimal* suspension state that such shape provides for the dispersed phase of CAM particles [6], as confirmed by the computational fluid-dynamics software VisiMix.

Table of contents

L. M. Alabdulsada, L. Kozma <i>Hopf-Rinow theorem of sub-Finslerian geometry</i>	2
Y. Aliyev <i>Geometric properties of interception curves</i>	3
M. Amram <i>Planar and non-planar degenerations with related fundamental groups</i>	5
N. Ando <i>Surfaces with zero mean curvature vector in 4-dimensional spaces</i>	6
B. Apanasov <i>Dynamics in nilpotent groups and deformations of locally symmetric rank one manifolds</i>	8
M. Atteya <i>Characterizing Linear Mappings Through Unital Algebra</i>	9
S. Sharma, V. K. Bhat <i>Edge resolvability and topological characteristics of zero-divisor graphs</i>	10
V. Bilet, O. Dovgoshey <i>From minimality to maximality via metric reflection</i>	12
D. Bolotov <i>Thurston norm and Euler classes of bounded mean curvature foliations on hyperbolic 3-Manifolds</i>	14
A. Bolsinov <i>Nijenhuis geometry and its applications</i>	15
E. Bonacci <i>Shape optimization in the batch crystallization of CAM</i>	17
F. Bulnes <i>Homotopies to Diffeomorphisms in Symplectic Field Theory</i>	18
D. Cheban <i>Global asymptotic stability of generalized homogeneous dynamical systems</i>	19
Y. Cherevko, V. Berezovski, J. Mikeš, Y. Fedchenko <i>Hyper-holomorphically projective mappings of hyper-Kähler manifolds</i>	21
S. Dann <i>On a problem of Fejes Toth</i>	22
M. Golasinski, T. de Melo, R. Bononi <i>Gottlieb groups of some Moore spaces</i>	23
A. Białyzyt, A. Denkowska, M. P. Denkowski <i>Inner semi-continuity of medial axes and conflict sets</i>	24
K. v. Dichter <i>The diameter-width-ratio for complete and pseudo-complete sets</i>	25
O. Dovhopiatyi <i>On the possibility of joining two pairs of points in convex domains using paths</i>	27
Yu. Drozd <i>Backström curves</i>	28