



International
Scientific Conference



Algebraic and Geometric Methods of Analysis



Devoted to 160 anniversary of
Dvytro Grave
(25.08.1863 - 19.12.1939)
Academician of the Ukrainian
Academy of Sciences, the
first director of the Institute of
Mathematics of NAS of Ukraine

May 29 – June 1, 2023
Odesa, Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

- | | |
|--|---|
| • Bolotov D. (<i>Kharkiv, Ukraine</i>) | • Konovenko N. (<i>Odesa, Ukraine</i>) |
| • Bondarenko V. (<i>Kyiv, Ukraine</i>) | • Maksymenko S. (<i>Kyiv, Ukraine</i>) |
| • Boychuk O. (<i>Kyiv, Ukraine</i>) | • Mikhailets V. (<i>Kyiv, Ukraine</i>) |
| • Boyko V. (<i>Kyiv, Ukraine</i>) | • Ostrovskiy V. (<i>Kyiv, Ukraine</i>) |
| • Cherevko Ye. (<i>Odesa, Ukraine</i>) | • Petravchuk A. (<i>Kyiv, Ukraine</i>) |
| • Dorogovtsev A. (<i>Kyiv, Ukraine</i>) | • Plaksa S. (<i>Kyiv, Ukraine</i>) |
| • Drozd Yu. (<i>Kyiv, Ukraine</i>) | • Portenko M. (<i>Kyiv, Ukraine</i>) |
| • Gerasymenko V. (<i>Kyiv, Ukraine</i>) | • Pratsiovytyi M. (<i>Kyiv, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Savchenko O. (<i>Kherson, Ukraine</i>) |
| • Kiosak V. (<i>Odesa, Ukraine</i>) | • Romanyuk A. (<i>Kyiv, Ukraine</i>) |
| • Kochubei A. (<i>Kyiv, Ukraine</i>) | • Timokha O. (<i>Kyiv, Ukraine</i>) |

ORGANIZING COMMITTEE

- | | |
|--|---|
| • Maksymenko S. (<i>Kyiv, Ukraine</i>) | • Cherevko Ye. (<i>Odesa, Ukraine</i>) |
| • Konovenko N. (<i>Odesa, Ukraine</i>) | • Osadchuk Ye. (<i>Odesa, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Sergeeva O. (<i>Odesa, Ukraine</i>) |

Explicit construction of explicit real algebraic functions and real algebraic manifolds via Reeb graphs

Naoki Kitazawa

(Institute of Mathematics for Industry, Kyushu University
Osaka Central Advanced Mathematical Institute)

E-mail: naokikitazawa.formath@gmail.com

In this talk, we present explicit real algebraic functions on explicit real manifolds via *Reeb graphs*. Our study is mainly motivated by real algebraic geometry, pioneered by Nash and Tognoli for example. A smooth closed manifold can be regarded as a non-singular real algebraic manifold. Existence theory of real algebraic manifolds and real algebraic maps has been also well-known. We can easily attain such objects and morphisms whereas it is very difficult to know explicitly. For more precise exposition, see [4] for example.

Of course some specific examples of real algebraic maps are well-known. Canonical embeddings and projections of unit spheres are simplest examples. As functions which are in considerable cases regarded as generalized ones, Lie groups and so-called *symmetric spaces* have nice functions represented as real polynomial functions. See [7] for example. However, it is difficult to know their global structures and properties explicitly in general.

Problem 1. Can we know global structures and properties of the functions and maps. For example, can we know information on preimages?

For this, we consider the following problem, established in [10]. This comes from singularity theory of smooth maps and applications to differential topology of manifolds. The *Reeb graph* of a smooth function is the graph whose underlying space is the natural quotient space of the manifold and consists of all connected components of preimages. Its vertex set is the set of all connected components containing some singular points of the function. As [9] shows, for smooth functions with finitely many singular values, we can have such graphs. [8] is a pioneering paper on this notion. Reeb graphs have some information of the manifolds nicely and fundamental and strong tools in geometry of manifolds.

Problem 2. Can we reconstruct a nice smooth function on some manifold whose Reeb graph is the given graph? We do not fix the manifold beforehand.

[10] constructs desired functions on closed surfaces for some nice graphs. [5] extends this to arbitrary finite graphs. [6] considers such a problem for a certain class of finite graphs and Morse functions such that connected components of preimages having no singular points are spheres. Our study [1] considers the following problem first. It is for functions on 3-dimensional closed manifolds. [9] presents a related general result through our informal discussions on [1].

Problem 3. Can we construct the function in Problem 2 with prescribed preimages?

This talk is on answers to the following problem, pioneered by the speaker first in [2].

Problem 4. Can we construct these functions and the manifolds in finer categories such as the real analytic category and the real algebraic category, for example?

We present our main results with several notions we need. An *algebraic domain* D is a bounded open set in the real affine space \mathbb{R}^k surrounded by finitely many mutually disjoint non-singular connected real algebraic hypersurfaces each S_j of which is the zero set of some real polynomial f_j . The *Poincaré-Reeb graph* of it is a canonically obtained graph whose underlying space is the natural quotient space of the closure \overline{D} of the domain and consists of all connected components of preimages for the restriction of the projection $\pi(x_1, \dots, x_k) := x_1$ to \overline{D} . Its vertex set is defined as the set of all connected components containing some singular points of the function $\pi|_{\overline{D}-D}$.

Theorem 5 ([2]). *Let G be a Poincaré-Reeb graph of $D \subset \mathbb{R}^k$. Let D be an algebraic domain represented as the intersection $\bigcap \{x \in \mathbb{R}^k \mid f_j(x) > 0\}$. Then we can construct a smooth real algebraic function whose Reeb graph is isomorphic to G on some non-singular real algebraic closed manifold.*

Example 6. Any Poincaré-Reeb graph G of any bounded connected open set $D \subset \mathbb{R}^k$ surrounded by finitely many mutually disjoint spheres of fixed radii satisfies the assumption of Theorem 5. See also FIGURE 1 of [2].

In the proof, first we construct a nice smooth real algebraic map into \mathbb{R}^k whose image is \overline{D} . More precisely, we construct one such that the preimage of a point in the boundary is a one-point set and that the preimage of a point in the interior is a sphere. Last we compose the projection.

Theorem 7 ([3]). *Let $l > 3$ and $m > 2$ be integers. Let $\{t_j\}_{j=1}^l$ be an increasing sequence of real numbers. Let $\{F_j\}_{j=1}^{l-1}$ be a family of smooth manifolds satisfying the following conditions.*

- F_1 and F_{l-1} are diffeomorphic to the $(m-1)$ -dimensional unit spheres S^{m-1} .
- The others are diffeomorphic to S^{m-1} or represented as connected sums of finitely many manifolds diffeomorphic to the products $S^j \times S^{m-j-1}$ for some integers $1 \leq j \leq m-2$: the connected sum is taken in the smooth category. For adjacent integers $1 \leq j \leq l-2$ and $j+1$, either F_j or F_{j+1} is not diffeomorphic to the unit sphere.

Then we have an m -dimensional non-singular real algebraic closed and connected manifold M and a smooth real algebraic function $f : M \rightarrow \mathbb{R}$ such that the number of singular points is finite, that $\{t_j\}_{j=1}^l$ is the set of all singular values and that the preimage $f^{-1}(p_j)$ is diffeomorphic to F_j for $p_j \in (t_j, t_{j+1})$.

The speaker was supported by JSPS KAKENHI Grant Number JP17H06128 and JSPS KAKENHI Grant Number JP22K18267 as a member. He is also supported by JSPS KAKENHI Grant Number JP23H05437. Principal investigators are all Osamu Saeki. The speaker is also a Postdoctoral Researcher at Osaka Central Advanced Mathematical Institute where he is not employed.

REFERENCES

- [1] N. Kitazawa, *On Reeb graphs induced from smooth functions on 3-dimensional closed orientable manifolds with finitely many singular values*, Topol. Methods in Nonlinear Anal. Vol. 59 No. 2B, 897–912, arXiv:1902.08841.
- [2] N. Kitazawa, *Real algebraic functions on closed manifolds whose Reeb graphs are given graphs*, a positive report for publication has been announced to have been sent and this will be published in Methods of Functional Analysis and Topology, arXiv:2302.02339v3.

- [3] N. Kitazawa, *Construction of real algebraic functions with prescribed preimages*, submitted to a refereed journal, arXiv:2303.00953.
- [4] J. Kollár, *Nash's work in algebraic geometry*, Bulletin (New Series) of the American Mathematical Society (2) 54, 2017, 307–324.
- [5] Y. Masumoto and O. Saeki, *A smooth function on a manifold with given Reeb graph*, Kyushu J. Math. 65 (2011), 75–84.
- [6] L. P. Michalak, *Realization of a graph as the Reeb graph of a Morse function on a manifold*. Topol. Methods in Nonlinear Anal. 52 (2) (2018), 749–762, arXiv:1805.06727.
- [7] S. Ramanujam, *Morse theory of certain symmetric spaces*, J. Diff. Geom. 3 (1969), 213–229.
- [8] G. Reeb, *Sur les points singuliers d'une forme de Pfaff complètement intégrable ou d'une fonction numérique*, Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences 222 (1946), 847–849.
- [9] O. Saeki, *Reeb spaces of smooth functions on manifolds*, International Mathematics Research Notices, maa301, Volume 2022, Issue 11, June 2022, 8740–8768, <https://doi.org/10.1093/imrn/maa301>, arXiv:2006.01689.
- [10] V. Sharko, *About Kronrod-Reeb graph of a function on a manifold*, Methods of Functional Analysis and Topology 12 (2006), 389–396.

Conformal equivalence of 3-webs

Konovenko N.

(ONTU, Odesa, Ukraine)

E-mail: ngkonovenko@gmail.com

Let 3-web $W_3 \langle \omega_1, \omega_2, \omega_3 \rangle$ defined in a domain D on the conformal plane (\mathbb{R}^2, g) . We say that this 3-web is *regular* in D if in this domain:

- (1) The discriminant

$$\tilde{\Delta} = -4I_2 + I_1^2 + 18I_1I_2 - 4I_1^3 - 27I_2^3$$

differs from zero.

- (2) Invariants

$$I_1 = \frac{J_2}{J_1^2} \quad \text{and} \quad I_2 = \frac{J_3}{J_1^3}$$

are functionally independent in the domain, that is, the differential 2-form $\Omega = dI_1 \wedge dI_2 \neq 0$.

Moreover, invariants I_1, I_2 are coordinates in the domain.

We remark that the elementary symmetric functions

$$\begin{aligned} J_1 &= \lambda_1 + \lambda_2 + \lambda_3, \\ J_2 &= \lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3, \\ J_3 &= \lambda_1\lambda_2\lambda_3 \end{aligned}$$

are S_3 - invariants and $\lambda_1, \lambda_2, \lambda_3$ are positive smooth functions.

Let's number now forms $\omega_1, \omega_2, \omega_3$ in the domain and say that the 3-web is *oriented* in the domain if in this numbering $\omega_1 \wedge \omega_2 = r_{12}\Omega$, where $r_{12} > 0$. In this case we'll scale forms ω_i in such a way, that

$$\omega_1 \wedge \omega_2 = \Omega.$$

In opposite case, we call the 3-web *non-oriented* and scale the 1-forms ω_i in such a way, that

$$\omega_1 \wedge \omega_2 = -\Omega.$$

V. Dryuma <i>On geodesic lines of Riemannian metric for Navier-Stokes equations</i>	29
L. Fardigola, K. Khalina <i>On controllability problems for the heat equation in a half-plane in the case of a pointwise control in the Dirichlet boundary condition</i>	32
V. Fedorchuk, V. Fedorchuk <i>On partial preliminary group classification of some class of $(1 + 3)$-dimensional Monge-Ampere equations. Two-dimensional Abelian Lie algebras</i>	34
B. Feshchenko <i>Homotopy type of stabilizers of functions with non-isolated singularities on surfaces</i>	36
N. Glazunov <i>On direct limits of Minkowski's balls, domains, and their critical lattices</i>	37
O. Gok <i>On KB(Kantorovich-Banach) spaces and KB operators</i>	39
M. Golański <i>On polynomial and regular maps of spheres</i>	40
O. Gutik, O. Prokhorenkova <i>On homomorphisms of bicyclic extensions of archimedean totally ordered groups</i>	41
O. Hukalov, V. Gordevskyy <i>The Interaction of an Infinite Number of Eddy Flows</i>	42
S. Ivković <i>Semi-Fredholm theory in unital C^*-algebras</i>	43
T. Jaiyeola, K. Ilori, O. Oyebola <i>On some non-associative hyper-algebraic structures</i>	45
J.-L. Mo <i>The rank of Mordell-Weil groups of surfaces</i>	47
J. Kąkol <i>On Asplund spaces $C_k(X)$ with the compact-open topology</i>	48
N. Kitazawa <i>Explicit construction of explicit real algebraic functions and real algebraic manifolds via Reeb graphs</i>	49
N. Kononenko <i>Conformal equivalence of 3-webs</i>	51
Y. Kopeliovich <i>The fundamental group of Riemann surface via Riemann's existence theorem</i>	52
G. Kuduk <i>Problem with integral conditions for evolution equations in Banach space</i>	53
I. Kuznietsova, S. Maksymenko <i>Deformational symmetries of functions with isolated singularities on the Mobius band</i>	54
R. L'hamri <i>Codes from zero-divisor super-λ graph</i>	55
L. Lotarets <i>Twisted Sasaki metric on the unit tangent bundle and harmonicity</i>	56