



International
Scientific Conference



Algebraic and Geometric Methods of Analysis



Devoted to 160 anniversary of
Dvytro Grave
(25.08.1863 - 19.12.1939)
Academician of the Ukrainian
Academy of Sciences, the
first director of the Institute of
Mathematics of NAS of Ukraine

May 29 – June 1, 2023
Odesa, Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric and topological methods in natural sciences
- Geometric problems in mathematical analysis

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National University of Technology
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

- | | |
|--|---|
| • Bolotov D. (<i>Kharkiv, Ukraine</i>) | • Konovenko N. (<i>Odesa, Ukraine</i>) |
| • Bondarenko V. (<i>Kyiv, Ukraine</i>) | • Maksymenko S. (<i>Kyiv, Ukraine</i>) |
| • Boychuk O. (<i>Kyiv, Ukraine</i>) | • Mikhailets V. (<i>Kyiv, Ukraine</i>) |
| • Boyko V. (<i>Kyiv, Ukraine</i>) | • Ostrovskiy V. (<i>Kyiv, Ukraine</i>) |
| • Cherevko Ye. (<i>Odesa, Ukraine</i>) | • Petravchuk A. (<i>Kyiv, Ukraine</i>) |
| • Dorogovtsev A. (<i>Kyiv, Ukraine</i>) | • Plaksa S. (<i>Kyiv, Ukraine</i>) |
| • Drozd Yu. (<i>Kyiv, Ukraine</i>) | • Portenko M. (<i>Kyiv, Ukraine</i>) |
| • Gerasymenko V. (<i>Kyiv, Ukraine</i>) | • Pratsiovytyi M. (<i>Kyiv, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Savchenko O. (<i>Kherson, Ukraine</i>) |
| • Kiosak V. (<i>Odesa, Ukraine</i>) | • Romanyuk A. (<i>Kyiv, Ukraine</i>) |
| • Kochubei A. (<i>Kyiv, Ukraine</i>) | • Timokha O. (<i>Kyiv, Ukraine</i>) |

ORGANIZING COMMITTEE

- | | |
|--|---|
| • Maksymenko S. (<i>Kyiv, Ukraine</i>) | • Cherevko Ye. (<i>Odesa, Ukraine</i>) |
| • Konovenko N. (<i>Odesa, Ukraine</i>) | • Osadchuk Ye. (<i>Odesa, Ukraine</i>) |
| • Fedchenko Yu. (<i>Odesa, Ukraine</i>) | • Sergeeva O. (<i>Odesa, Ukraine</i>) |

- 1) **Type** T_{023} contains: $E \wr \tilde{A}_n \wr H$, $\tilde{T}_n \wr H$, where $H \in \{\tilde{A}_n, \tilde{A}_{n^2}, S_n\}$. There are 6 subgroups.
- 2) **The second type of subgroups is subclass in** T_{023} with new base of wreath product subgroup \tilde{A}_{n^2} : $E \wr S_n \wr \tilde{A}_{n^2}$, $E \wr A_n \wr \tilde{A}_{n^2}$, $E \wr N_i(S_n \wr S_n)$. Therefore this class has 12 new subgroups. Thus, the total number of normal subgroups in **Type** T_{023} is 18.
- 3) **Type** T_{003} : $A_{00(n^2)}^{(3)} = E \wr E \wr \tilde{A}_{n^2}$, \tilde{T}_{n^2} , $\tilde{T}_n^{(3)}$. Hence, here are 3 new subgroups.
- 4) **Type** T_{123} : $N_i(S_n \wr S_n) \wr S_n$, $N_i(S_n \wr S_n) \wr \tilde{A}_n$ and $N_i(S_n \wr S_n) \wr \tilde{A}_{n^2}$. Thus, there are 29 new normal subgroups in T_{123} , taking into account repetition [5].

REFERENCES

- [1] Skuratovskii R.V. Invariant structures of wreath product of symmetric groups. Naukovuy Chasopus of Science hour writing of the National Pedagogical University named after M.P. Dragomanova. (in ukrainian) Series 01. Physics and Mathematics. — 2009. Issue 10. — P. 163 – 178.
- [2] Sachkov, V.N., Combinatorial methods in discrete Mathematics. Encyclopedia of mathematics and its applications 55. Cambridge Press. 2008. P. 305.
- [3] Kaloujnine L. A. Sur les p -group de Sylow du groupe symétrique du degré p^m . // C. R. Acad. Sci. Paris. — 1945. — **221**. — P. 222–224.
- [4] Drozd, Y.A., Skuratovskii R.V. Generators and relations for wreath products of groups. *Ukr. Math. J.* (2008), *60*, pp. 1168–1171.
- [5] Ruslan Skuratovskii. Normal subgroups of iterated wreath products of symmetric groups and alternating with symmetric groups. 2022, Source: [https://doi.org/10.48550/arXiv.2108.03752]

Asymptotic behavior of the widths of classes of the generalized Poisson integrals

Anatolii Serdyuk

(Institute of Mathematics of NAS of Ukraine, Kyiv, Ukraine)

E-mail: serdyuk@imath.kiev.ua

Igor Sokolenko

(Institute of Mathematics of NAS of Ukraine, Kyiv, Ukraine)

E-mail: sokol@imath.kiev.ua

Let L_p , $1 \leq p \leq \infty$, and C be the spaces of 2π -periodic functions with standard norms $\|\cdot\|_{L_p}$ and $\|\cdot\|_C$, respectively.

Denote by $C_{\beta, p}^\psi$, $1 \leq p \leq \infty$, the set of all 2π -periodic functions f , representable as convolution

$$f(x) = \frac{a_0}{2} + \frac{1}{\pi} \int_{-\pi}^{\pi} \varphi(x-t) \Psi_{\bar{\beta}}(t) dt, \quad a_0 \in \mathbb{R}, \quad \varphi \in B_p^0 = \{g \in L_p : \|g\|_p \leq 1, g \perp 1\}, \quad (1)$$

with a fixed generated kernel $\Psi_{\bar{\beta}} \in L_{p'}$, $1/p + 1/p' = 1$, the Fourier series of which has the form

$$S[\Psi_{\bar{\beta}}](t) = \sum_{k=1}^{\infty} \psi(k) \cos\left(kt - \frac{\beta_k \pi}{2}\right), \quad \beta_k \in \mathbb{R}, \quad \psi(k) \geq 0.$$

A function f in the representation (1) is called $(\psi, \bar{\beta})$ -integral of the function φ and is denoted by $\mathcal{J}_{\bar{\beta}}^\psi \varphi$ ($f = \mathcal{J}_{\bar{\beta}}^\psi \varphi$). If $\psi(k) \neq 0$, $k \in \mathbb{N}$, then the function φ in the representation (1) is called

$(\psi, \bar{\beta})$ -derivative of the function f and is denoted by $f_{\bar{\beta}}^{\psi}$ ($\varphi = f_{\bar{\beta}}^{\psi}$). The concepts of $(\psi, \bar{\beta})$ -integral and $(\psi, \bar{\beta})$ -derivative was introduced by Stepanets [1]. Since $\varphi \in L_p$ and $\Psi_{\bar{\beta}} \in L_{p'}$, then the function f of the form (1) is a continuous function, i.e. $C_{\bar{\beta}, p}^{\psi} \subset C$ (see [1, Proposition 3.9.2.]).

In the case $\beta_k \equiv \beta$, $\beta \in \mathbb{R}$, the classes $C_{\bar{\beta}, p}^{\psi}$ are denoted by $C_{\beta, p}^{\psi}$.

For $\psi(k) = k^{-r}$, $r > 0$, the classes $C_{\bar{\beta}, p}^{\psi}$ and $C_{\beta, p}^{\psi}$ are denoted by $W_{\bar{\beta}, p}^r$ and $W_{\beta, p}^r$, respectively. The classes $W_{\beta, p}^r$ are the well-known Weyl-Nagy classes (see [1]). In other words, $W_{\beta, p}^r$, $1 \leq p \leq \infty$, are the classes of 2π -periodic functions f , representable as convolutions of the Weyl-Nagy kernels $B_{r, \beta}(t) = \sum_{k=1}^{\infty} k^{-r} \cos\left(kt - \frac{\beta\pi}{2}\right)$, $r > 0$, $\beta \in \mathbb{R}$, with functions $\varphi \in B_p^0$.

If $r \in \mathbb{N}$ and $\beta = r$, then the functions $B_{r, \beta}$ are the well-known Bernoulli kernels and the corresponding classes $W_{\beta, p}^r$ coincide with the well-known classes W_p^r which consist of 2π -periodic functions f with absolutely continuous derivatives $f^{(k)}$ up to $(r-1)$ -th order inclusive and such that $\|f^{(r)}\|_p \leq 1$.

For $\psi(k) = e^{-\alpha k^r}$, $\alpha > 0$, $r > 0$, the classes $C_{\bar{\beta}, p}^{\psi}$ and $C_{\beta, p}^{\psi}$ are denoted by $C_{\bar{\beta}, p}^{\alpha, r}$ and $C_{\beta, p}^{\alpha, r}$, respectively. The sets $C_{\beta, p}^{\alpha, r}$ are well-known classes of the generalized Poisson integrals [1], i.e. classes of convolutions with the generalized Poisson kernels

$$P_{\alpha, r, \beta}(t) = \sum_{k=1}^{\infty} e^{-\alpha k^r} \cos\left(kt - \frac{\beta\pi}{2}\right), \quad \alpha > 0, \quad r > 0, \quad \beta \in \mathbb{R}.$$

Further, let K be a convex centrally symmetric subset of C and let B be a unit ball of the space C . Let also F_N be an arbitrary N -dimensional subspace of space C , $N \in \mathbb{N}$, and $\mathcal{L}(C, F_N)$ be a set of linear operators from C to F_N . By $\mathcal{P}(C, F_N)$ denote the subset of projection operators of the set $\mathcal{L}(C, F_N)$, that is, the set of the operators A of linear projection onto the set F_N such that $Af = f$ when $f \in F_N$. The quantities

$$b_N(K, C) = \sup_{F_{N+1}} \sup\{\varepsilon > 0 : \varepsilon B \cap F_{N+1} \subset K\},$$

$$d_N(K, C) = \inf_{F_N} \sup_{f \in K} \inf_{u \in F_N} \|f - u\|_C,$$

$$\lambda_N(K, C) = \inf_{F_N} \inf_{A \in \mathcal{L}(C, F_N)} \sup_{f \in K} \|f - Af\|_C,$$

$$\pi_N(K, C) = \inf_{F_N} \inf_{A \in \mathcal{P}(C, F_N)} \sup_{f \in K} \|f - Af\|_C,$$

are called Bernstein, Kolmogorov, linear, and projection N -widths of the set K in the space C , respectively.

The results containing order estimates of the widths b_N , d_N , λ_N or π_N in the case of $K = C_{\bar{\beta}, p}^{\psi}$ (and, in particular, $W_{\bar{\beta}, p}^r$ and $C_{\bar{\beta}, p}^{\psi}$) can be found, for example, in the monographs of Tikhomirov, Pinkus, Kornejchuk, Romanyuk, Temlyakov etc.

Theorem 1. Let $\bar{\beta} = \{\beta_k\}_{k=1}^{\infty}$, $\beta_k \in \mathbb{R}$, $\alpha > 0$, $r > 1$, $n \in \mathbb{N}$ and be such that

$$(n-1)^r > \frac{1}{\alpha}, \quad (2)$$

then the following inequalities hold

$$\begin{aligned} & \frac{1}{\sqrt{\pi}} e^{-\alpha n^r} \left(1 - \frac{2\gamma_{\alpha,r,n} e^{-2\alpha r(n-1)^{r-1}}}{1 + 2\gamma_{\alpha,r,n} e^{-2\alpha r(n-1)^{r-1}}} \right)^{\frac{1}{2}} \leq P_{2n}(C_{\bar{\beta},2}^{\alpha,r}, C) \\ & \leq P_{2n-1}(C_{\bar{\beta},2}^{\alpha,r}, C) \leq \frac{1}{\sqrt{\pi}} e^{-\alpha n^r} \left(1 + e^{-2\alpha r n^{r-1}} \left(1 + \frac{1}{2\alpha r n^{r-1}} \right) \right)^{\frac{1}{2}}, \end{aligned} \quad (3)$$

where P_N is any of the widths b_N, d_N, λ_N or π_N and

$$\gamma_{\alpha,r,n} = \left(1 + \frac{1}{\alpha r(n-1)^{r-1}} + e^{-2\alpha(n-1)^r} \max \left\{ e^{4\alpha}, \frac{e^2}{\alpha^{1+1/r}} \right\} \right). \quad (4)$$

Theorem 2. Let $\bar{\beta} = \{\beta_k\}_{k=1}^{\infty}, \beta_k \in \mathbb{R}, \alpha > 0, r > 1, n \in \mathbb{N}$ and the condition (2) is satisfied. Then as $n \rightarrow \infty$ the following asymptotic equalities hold

$$\left. \begin{aligned} & P_{2n}(C_{\bar{\beta},2}^{\alpha,r}, C) \\ & P_{2n-1}(C_{\bar{\beta},2}^{\alpha,r}, C) \end{aligned} \right\} = e^{-\alpha n^r} \left(\frac{1}{\sqrt{\pi}} + \mathcal{O}(1) \gamma_{\alpha,r,n} e^{-\alpha r(n-1)^{r-1}} \right), \quad (5)$$

where P_N is any of the widths b_N, d_N, λ_N or π_N and $\gamma_{\alpha,r,n}$ is defined by (4) and $\mathcal{O}(1)$ are the quantities uniformly bounded in all parameters.

Note that the Theorem 2 complements the results of the works of Shevaldin (1992), Stepanets and Serdyuk (1995), Serdyuk (1999), Serdyuk and Sokolenko (2011), Serdyuk and Bodenchuk (2013), which contain exact estimates for the widths of the classes of convolutions with classical or generalized Poisson kernels.

This work was partially supported by the Grant H2020-MSCA-RISE-2019, project number 873071 (SOMPATY: Spectral Optimization: From Mathematics to Physics and Advanced Technology).

REFERENCES

- [1] A.I. Stepanets. *Methods of Approximation Theory*. Utrecht: VSP, 2005.

Poincaré-Reeb graphs of real algebraic domains

Arnaud Bodin

(Université de Lille, CNRS, Laboratoire Paul Painlevé, Lille, France)

E-mail: arnaud.bodin@univ-lille.fr

Patrick Popescu-Pampu

(Université de Lille, CNRS, Laboratoire Paul Painlevé, Lille, France)

E-mail: patrick.popescu-pampu@univ-lille.fr

Miruna-Stefana Sorea

(SISSA - Scuola Internazionale Superiore di Studi Avanzati, Trieste, Italy and Lucian Blaga University of Sibiu, Romania)

E-mail: mirunastefana.sorea@sisssa.it

An *algebraic domain* is a closed topological subsurface of a real affine plane whose boundary consists of disjoint smooth connected components of real algebraic plane curves. We study the geometric shape of an algebraic domain by collapsing all vertical segments contained in it:

E. Petrov, R. Salimov <i>Fixed point theorem for mappings contracting perimeters of triangles and its generalizations</i>	84
A. Prishlyak <i>Structure of codimensional one flows on the 2-sphere with holes</i>	86
A. Arman, A. Bondarenko, A. Prymak <i>Convex bodies of constant width with exponential illumination number</i>	88
G. Riabov <i>Bifurcation points in random dynamical systems</i>	89
D. Ryabogin <i>On symmetries of sections of convex bodies</i>	90
A. Savchenko <i>Fuzzy metrization of spaces of \star-measures</i>	91
O. Sazonova <i>Continual distribution with acceleration and condensation flows</i>	92
R. Servadei <i>On a flower-shape geometry</i>	93
E. Sevost'yanov, N. Ilkevych <i>On equicontinuity of families of mappings with one normalization condition by the prime ends</i>	93
O. Shugailo <i>Equiaffine immersions of codimension two with flat connection</i>	95
H. Sinyukova <i>Some vanishing theorems of sufficient character about holomorphically projective mappings of Kahlerian spaces on the whole</i>	97
A. Skryabina, P. Stegantseva <i>Investigation of the connection between different models of topologies on a finite set</i>	98
R. Skuratovskii <i>Normal subgroups of iterated wreath products of symmetric groups and alternating with symmetric groups</i>	99
A. Serdyuk, I. Sokolenko <i>Asymptotic behavior of the widths of classes of the generalized Poisson integrals</i>	102
A. Bodin, P. Popescu-Pampu, M.-S. Sorea <i>Poincaré-Reeb graphs of real algebraic domains</i>	104
D. Dmytryshyn, D. Gray, and A. Stokolos <i>On univalent trinomials</i>	105
Kh. Sukhorukova <i>On K-ultrametrics and \ast-measures</i>	106
S. Tateno <i>The Iwasawa invariants of Z_p^d-covers of links</i>	106
A. Teleman <i>The Riemann-Hilbert problem and holomorphic bundles framed along a real hypersurface</i>	107
Y. Teplitskaya <i>About some Steiner trees</i>	109
J. Ueki <i>The multiplicities of non-acyclic SL_2-representations and L-functions of twisted Whitehead links</i>	110
J. F. Peters, T. Vergili <i>Proximal connectedness. Spatially and descriptively connected spaces</i>	111