


International
Online Conference



**Algebraic
and Geometric
Methods of Analysis**

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Drozd Yu.

(Kyiv, Ukraine)

Maksymenko S.

(Kyiv, Ukraine)

Plaksa S.

(Kyiv, Ukraine)

Prishlyak A.

(Kyiv, Ukraine)

Bakhtin O.

(Kyiv, Ukraine)

Balan V.

(Bucharest, Romania)

Banakh T.

(Lviv, Ukraine)

Borysenko O.

(Kharkiv, Ukraine)

Cherevko Ye.

(Odesa, Ukraine)

Fedchenko Yu.

(Odesa, Ukraine)

Karlova O.

(Chernivtsi, Ukraine)

Kiosak V.

(Odessa, Ukraine)

Konovenko N.

(Odessa, Ukraine)

Lyubashenko V.

(Kyiv, Ukraine)

Matsumoto K.

(Yamagata, Japan)

Mormul P.

(Warsaw, Poland)

Mykhailyuk V.

(Chernivtsi, Ukraine)

Plachta L.

(Krakov, Poland)

Pokas S.

(Odessa, Ukraine)

Sabitov I.

(Moscow, Russia)

Savchenko O.

(Kherson, Ukraine)

Sergeeva A.

(Odessa, Ukraine)

Shelekhov A.

(Tver, Russia)

Zarichnyi M.

(Lviv, Ukraine)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

ORGANIZING COMMITTEE

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohlyk O.

Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.

Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

On the metric equations generated by symplectic deformations on $P_2(\mathbb{C})$

Anatolij Prykarpatsky

(Drohobych Pedagogical University, Lviv region, Ukraine, and the Department of Computer Science at the Cracov University of Technology, Kraków 31-155, Poland)

E-mail: pryk.anat@cybergal.com

Ihor Mykytyuk

(Pidstryhach Institute for Applied Problems of Mechanics and Mathematics at the NAS, Lviv, 79060, Ukraine, and the Department of Computer Science at the Cracov University of Technology, Kraków, 31-155, Poland)

E-mail: mykytyuk.i@gmail.com

We analyzed in detail the cohomology structure of the symplectic form deformation and applied recently developed generalized transformations which were suggested in the classical works by Enneper and Weierstrass about one and half century ago and succeeded in reformulating the "symplectic" modification of the Monge-Ampere equation by means of specially constructed coordinates, related with the natural projector expansion on $T(P_2(\mathbb{C}))$ and found its special solutions. Let us consider a compact complex n -dimensional manifold M^n , endowed with the Kähler symplectic form $\omega \in \Lambda^2(M^n)$ and define the related Monge-Ampere equation, describing a deformation of this symplectic structure:

$$(\omega + i\partial\bar{\partial}\varphi)^n = (\exp f)\omega^n \quad (1)$$

under the normalizing conditions

$$\int_{M^n} (\exp f)\omega^n = \int_{M^n} \omega^n, \quad \int_{M^n} \varphi\omega^n = 0, \quad (2)$$

where $\varphi \in C^\infty(M^n; \mathbb{R})$ is a real valued function on M^n and $\bar{\partial}$ is the complex ∂ -bar differential, corresponding to the standard differential splitting $d = \partial \oplus \bar{\partial} : \Lambda(M^n) \rightarrow \Lambda(M^n)$ on the complex manifold M^n . In a general case it was supposed [16] that if the two-form $(\omega + i\partial\bar{\partial}\varphi) \in \Lambda^2(M^n)$ is real valued and the first Chern class $c_1(M^n) = 0$ of a Kähler manifold M^n , then there exists a Riemannian metric $g : T(M^n) \times T(M^n) \rightarrow \mathbb{R}$ of the Calabi-Yau type, whose holonomy group [5, 8] coincides with a subgroup of the Lie group $SU(2)$, generating, in particular, a so called Einsteinian metric. The equation (1) is always [16] solvable, yet its holonomy groups, in general, not classified and its unitarity remains to be open.

We here also remark that there exists a slightly different modified Monge-Ampere type deformation equation

$$(\omega + dJ^*d\varphi)^n = (\exp f)\omega^n, \quad (3)$$

on a real symplectic manifold $\bar{M}^{2n} \simeq M^n$, where $f \in C^\infty(\bar{M}^{2n}; \mathbb{R})$ and $J : T(\bar{M}^{2n}) \rightarrow T(\bar{M}^{2n})$, $J^2 = -I$, is a suitably chosen nonintegrable quasi-complex structure on the manifold \bar{M}^{2n} and $J^* : T^*(\bar{M}^{2n}) \rightarrow T^*(\bar{M}^{2n})$ denotes its conjugate. It was proved [2] that if the structure $J : T(\bar{M}^{2n}) \rightarrow T(\bar{M}^{2n})$ is integrable, then the equation (3) reduces to the Monge-Ampere equation (1) on the related complex manifold $M^n \simeq \bar{M}^{2n}$ owing to the classical Newlander-Nirenberg [10] criterion. Otherwise, if the equation (3) is solvable for its arbitrarily chosen right hand side, then the quasi-complex structure $J : T(\bar{M}^{2n}) \rightarrow T(\bar{M}^{2n})$ proves to be necessary [2, 9, 11] a complex one, once more reducing the equation (3) to the Monge-Ampere equation (1).

In our note we are interested in the following "symplectic" modification

$$(\omega + dd^s\varphi)^2 = (\exp f)\omega^2 \quad (4)$$

of the Monge-Ampere (1) on the complex Kähler manifold $M^2 = P_2(\mathbb{C})$, where $\varphi \in \Lambda^2(M^2)$ is a searched for two-form and $d^s := (-1)^{k+1} \star_s d \star_s$, $dd^s = -d^s d$, denotes the symplectic Hodge type differentiation. It is well known that any compact two-dimensional Kähler manifold M^2 with the Chern class $c_1(M^2) = 0$ is hiper-Kähler, possessing exactly three Kähler fundamental forms ω_I, ω_J and $\omega_K \in \Lambda^2(\bar{M}^4)$, corresponding to three complex structures I, J and $K : T(\bar{M}^4) \rightarrow T(\bar{M}^4)$. As for the compact projective two-dimensional Kähler manifold $M^2 = P_2(\mathbb{C})$ the Chern class $c_1(M^2) \neq 0$, it is not hiper-Kähler, its holomorphic volume two-form is not composed of the symplectic forms ω_J and $\omega_K \in \Lambda^2(\bar{M}^4)$. Notwithstanding this fact, based on the equalities (??) and the well known [1, 14, 15] relationship

$$\star_s \eta = -\eta \quad (5)$$

for an arbitrary "primitive" holomorphic volume two-form $\eta \in \Lambda_{hol}^2(M^2)$, satisfying the additional condition $\eta \wedge \omega = 0$, one easily derives that for any two cohomological "primitive" holomorphic volume two-forms Ω_1 and $\Omega_2 \in \Lambda_{hol}^2(M^2)$ there holds the following interesting relationship:

$$\Omega_1 - \Omega_2 = dd^s \psi \quad (6)$$

for some smooth two-form $\psi \in \Lambda^2(M^2)$, solving the problem (4) for the case when the symplectic structure $\omega \in \Lambda^2(M^2)$ is replaced by a holomorphic volume form $\Omega \in \Lambda_{hol}^2(M^2)$. Having analyzed in detail the cohomology structure of the two-form expression $(\omega + dd^s \varphi) \in \Lambda^2(M^2)$ and applied generalized transformations which were suggested in the classical works by Enneper [4] and Weierstrass [13] about one and half century ago and recently developed in [6], we succeeded in reformulating the "symplectic" modification of the Monge-Ampere (4) by means of specially constructed coordinates, related with the natural projector expansion from in $P_2(\mathbb{C})$ and find its special solutions.

REFERENCES

- [1] Chern S.S. Complex manifolds. The University of Chicago Publ, 1956
- [2] Delanoë Ph. L'analogie presque-complexe de l'équation de Calabi-Yau. Osaka J. Math.33 (1996), 829-846
- [3] Ehresmann C., Libermann P. Sur le problème d'équivalence des formes différentielles extérieures quadratiques. C. R. Acad. Sci. Paris. V. 229 (1949), 697-698.
- [4] Enneper A. Nachr. Konigl. Gesell. Wissensch. Georg - Augustus Univ. Göttingen 12 (1868), 258-277.
- [5] Joyce D. Compact manifolds with special holonomy. Oxford Univ. Press, 2000.
- [6] Konopelchenko B.G., Taimanov I.A. Constant mean curvature surfaces via integrable dynamical system. Consortium EINSTEIN, Dipartimento di Fisica, Università, arXiv:dg-ga/9505006v1 26 May 1995
- [7] Libermann P. Sur les structures presque complexes et autres structures infinitésimales irrégulières. Bull. Soc. Math. France. V. 83 (1955), 195-224.
- [8] Lichnerowicz A. Théorie globale des connexions et des groupes d'holonomie, Cremonese, Roma, 1955
- [9] Moser J. On the volume elements on a manifold, Trans. Amer. Math. Soc. 120 (1965), 286-294
- [10] Nijenhuis N. et Wolf W.B. Some integration problems in almost-complex and complex manifolds, Annals of Math. 77 (1963), 424-489
- [11] Nongue W., Peng Z. On a generalized Calabi-Yau equation. arxiv:0911.0784
- [12] Tseng Li-Sheng, Yau S.-T. Cohomology and Hodge theory on symplectic manifolds. I. arXiv:0909.5418
- [13] Weierstrass K. Fortsetzung der Untersuchung über die Minimalflächen, Mathematische Werke, Vol. 3, Verlagsbuch-handlung, Hillesheim, 1866, 219-248
- [14] Weil A. Introduction à l'Étude des variétés Kähleriennes. Publications de l'Institut de Mathématique de l'Université de Nancago; VI., Paris: Hermann, 1958
- [15] Wells R.O. Differential analysis on complex manifolds. Prentice Hall Inc., NJ, 1973
- [16] Yau S.-T. On the Ricci curvature of compact Kähler manifold and the complex Monge-Ampère equation. I. Comm. Pure and Applied Math. 31 (1978), 339-411

Y. Chapovskyi, D. Efimov, A. Petravchuk <i>Centralizers of elements in Lie algebras of vector fields</i>	113
E. Petrov, R. Salimov <i>Properties of quasisymmetric mappings to preserve the structures of spaces</i>	114
D. R. Popovych <i>IW contractions and their generalizations</i>	116
Prabhjot Singh <i>Weak Separation Condition coincides Finite Type Condition</i>	118
Pratyush Pranav <i>Topological data analysis for cosmology: theory and applications</i>	120
A. Prishlyak, Ch. Hatamian <i>Morse flows with singularities on boundary of 3-manifolds</i>	121
V. Prokip <i>On the matrix equation $AX - YB = C$ over Bezout domains</i>	122
A. Prykarpatsky, I. Mykytyuk <i>On the metric equations generated by symplectic deformations on $P_2(C)$</i>	124
O. Reinov <i>On a result of G. Pisier concerning Sidon sets</i>	126
A. K. Sadullaev, F. G. Mukhamadiev <i>On the τ-placedness of space of the permutation degree</i>	127
T. N Safarov <i>The analogue of Darboux equation in Galilean space</i>	128
V. M. Safonov, I. V. Zamrii, O. V. Safonova <i>On countable multiplicity of mappings</i>	129
A. Savchenko <i>Fuzzy ultrametrization of spaces of non-additive measures on fuzzy ultrametric spaces</i>	130
O. Sazonova <i>About one class of continual approximate solutions with arbitrary density</i>	131
J. Segert <i>Painlevé VI Solutions From Equivariant ADHM Instanton Bundles</i>	133
A. Serdyuk, I. Sokolenko <i>Asymptotic estimates for the widths of classes of periodic functions of high smoothness</i>	135
H. Sinyukova <i>Some generalizations of the known theorems of the type of geodesical unique definability</i>	138
S. Som, A. Bera, L. K. Dey <i>Some remarks on the Metrizable of F-metric spaces</i>	139
P. Stegantseva, M. Grechneva <i>The surfaces with the flat normal connection and the constant curvature of Grassmann image in Minkowski space</i>	140
A. Skryabina, P. Stegantseva <i>The relation between T_0-topologies with the weight $2^{n-2} < k \leq 2^{n-1}$ on n-element set and T_0-topologies close to the discrete on $(n - 1)$-element set</i>	142
Ya. B. Stelmakh <i>The Golomb and Kirch topologies on the set of nonzero integers</i>	143
D. Dmitrishin, A. Stokolos <i>On symmetrization of univalent polynomials</i>	144