

International
Online Conference



**Algebraic
and Geometric
Methods of Analysis**

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Drozd Yu.

(Kyiv, Ukraine)

Maksymenko S.

(Kyiv, Ukraine)

Plaksa S.

(Kyiv, Ukraine)

Prishlyak A.

(Kyiv, Ukraine)

Bakhtin O.

(Kyiv, Ukraine)

Balan V.

(Bucharest, Romania)

Banakh T.

(Lviv, Ukraine)

Borysenko O.

(Kharkiv, Ukraine)

Cherevko Ye.

(Odesa, Ukraine)

Fedchenko Yu.

(Odesa, Ukraine)

Karlova O.

(Chernivtsi, Ukraine)

Kiosak V.

(Odessa, Ukraine)

Konovenko N.

(Odessa, Ukraine)

Lyubashenko V.

(Kyiv, Ukraine)

Matsumoto K.

(Yamagata, Japan)

Mormul P.

(Warsaw, Poland)

Mykhailyuk V.

(Chernivtsi, Ukraine)

Plachta L.

(Krakov, Poland)

Pokas S.

(Odessa, Ukraine)

Sabitov I.

(Moscow, Russia)

Savchenko O.

(Kherson, Ukraine)

Sergeeva A.

(Odessa, Ukraine)

Shelekhov A.

(Tver, Russia)

Zarichnyi M.

(Lviv, Ukraine)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

ORGANIZING COMMITTEE

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohlyk O.

Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.

Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

On the solution of separate differential equations with variational derivatives of the first and second orders

Leonid A. Yanovich

(Institute of Mathematics of the National Academy of Sciences of Belarus, Minsk, Belarus)

E-mail: yanovich@im.bas-net.by

Marina V. Ignatenko

(Belarusian State University, Minsk, Belarus)

E-mail: ignatenkomv@bsu.by

The theory of equations with variational (functional) derivatives is a fairly extensive area of mathematics. This class of equations has numerous applications in statistical physics, quantum field theory, hydromechanics and other fields. The theory of variational derivatives, differential and integro-differential equations with variational derivatives is quite fully stated, for example, in the monographs [1–3] and the works [4–10].

The problems that are formulated and investigated in this area of mathematics are similar to the problems considered in the case of ordinary and partial differential equations. Explicit formulas for solving equations with variational derivatives are known only in a few cases. This applies mainly to the set of linear equations [5–8]. Therefore, the main methods for solving such equations are approximate.

The problem of an approximate solution of equations with variational derivatives is not sufficiently studied. When solving this class of problems, it may be useful to apply methods that take into account the given initial and boundary values. In particular, in the Cauchy problem for an equation of the n -th order, the desired functional $F(x)$ can be approximately found from the values of the functional $F(x_0)$ and its variational derivatives up to the $(n - 1)$ -th order, which are known at the point $x_0(t)$. For this, it is natural to use the operator interpolation apparatus [11, 12]. Consider one of the ways for the approximate solution of equations with variational derivatives, based on interpolation of the functional included in the equation.

We formulate the definition of the variational derivative for functionals defined on sets of functions [9]. Let X be a linear space of real functions defined on a segment $[a, b]$ of the real axis \mathbb{R} , and F be an operator or functional given on the X .

The k -th order Gateaux differential $\delta^k F[x; h_1, h_2, \dots, h_k]$ ($k \in \mathbb{N}$) of the mapping F at the point $x \in X$ in the directions $h_1, h_2, \dots, h_k \in X$ is defined by the equality

$$\begin{aligned} \delta^k F[x; h_1, h_2, \dots, h_k] &= \lim_{\lambda \rightarrow 0} \frac{\delta^{k-1} F[x + \lambda h_k; h_1, h_2, \dots, h_{k-1}] - \delta^{k-1} F[x; h_1, h_2, \dots, h_{k-1}]}{\lambda} = \\ &= \left. \frac{\partial^k F(x + \lambda_1 h_1 + \lambda_2 h_2 + \dots + \lambda_k h_k)}{\partial \lambda_k \dots \partial \lambda_1} \right|_{\lambda_1 = \dots = \lambda_k = 0}, \delta^0 F[x] \equiv F(x). \end{aligned}$$

If there exists the k -th order Gateaux differential $\delta^k F[x; h_1, h_2, \dots, h_k]$ ($x, h_i \in X$; $i = 1, 2, \dots, k$) of the functional $F(x)$ at the point $x \in X$ in the directions $h_1, h_2, \dots, h_k \in X$, that can be represented as

$$\delta^k F[x; h_1, h_2, \dots, h_k] = \int_{[a,b]^k} a(x; t_1, \dots, t_k) h_1(t_1) \dots h_k(t_k) dt_1 \dots dt_k, \quad (1)$$

where $a(x; t_1, \dots, t_k)$ is some function depending on $x = x(s)$ and variables $t_1, \dots, t_k \in \mathbb{R}$, then $a(x; t_1, \dots, t_k)$ is called the variational derivative of the k -th order of the functional $F(x)$ with respect to x at the point $t = (t_1, t_2, \dots, t_k)$ and denoted by the symbol $\frac{\delta^k F(x)}{\delta x(t_1) \dots \delta x(t_k)}$. Variational derivatives can be generalized functions and other types of functionals.

As X , one can choose the space $C[a, b]$ of continuous functions with a uniform norm, the Hilbert space $L_2[a, b]$ or any other space such that the integral on the right-hand side of (1) makes sense.

We give formulas for the exact solution of some of the simplest differential equations with variational derivatives.

For example, for the equation

$$\frac{\delta F(x)}{\delta x(t)} = 2p(t) \cos x(t) \int_0^1 p(t) \sin x(t) dt,$$

the solution is the functional $F(x) = \left(\int_0^1 p(t) \sin x(t) dt \right)^2$, where $p(t)$ and $x(t)$ are elements of space $C[0, 1]$.

The functional $F(x) = \int_a^b p(t) f[x(t)] dt$ is the solution of the equation $\frac{\delta F(x)}{\delta x(t)} = p(t) f'[x(t)]$.

The solution of the equation

$$\frac{\delta F(x)}{\delta x(t)} = p_0(t) e^{x(t)} + p_1(t) \cos x(t) + p_2(t) \sin x(t) + p_3(t) x(t) + p_4(t) \int_0^1 p_4(\tau) x(\tau) d\tau$$

has the form

$$F(x) = \int_0^1 \left[p_0(t) e^{x(t)} + p_1(t) \sin x(t) - p_2(t) \cos x(t) + \frac{1}{2} \left(\int_0^1 p_4(t) x(t) dt \right)^2 \right] p(t) \sin x(t) dt,$$

where $p_0(t)$ ($i = 0, 1, \dots, 4$) are arbitrary functions for which reduced integrals exist.

Next, we consider a differential equation of the hyperbolic type with the second-order variational derivatives:

$$\frac{\delta^2 u(x, y)}{\delta x^2(t)} - a^2(t) \frac{\delta^2 u(x, y)}{\delta y^2(t)} = 0 \quad (x = x(t) \geq 0, y = y(t), a(t) \neq 0; t \in [a, b] \subseteq \mathbb{R}). \quad (2)$$

The solution of this equation is the functional

$$u(x, y) = f_1 \left[\int_a^b (y(t) + a(t)x(t)) dt \right] + f_2 \left[\int_a^b (y(t) - a(t)x(t)) dt \right], \quad (3)$$

where $f_1(\cdot) \ni f_2(\cdot)$ are any functions that are twice differentiable on \mathbb{R} . The representation (3) is an analogue of the classical D'Alembert formula.

We give the Hermite interpolation formula $H(x, y)$ with respect to a single node of the second multiplicity, which is an approximation to the solution $u(x, y)$ of the Cauchy problem for equation (2) with the initial conditions

$$u(x_0, y) = u_0(y), \quad \frac{\delta u(x_0, y)}{\delta x(t)} = u_1(y), \quad (4)$$

where $u_0(y)$ and $u_1(y)$ are some functionals defined on $C[a, b]$.

Theorem 1. An approximate solution of the Cauchy problem (2), (4) can be represented as

$$H(x, y) = u_0(y) + u_1(y) \int_a^b (x(t) - x_0(t)) dt + \frac{1}{2} a^2(t) u_0''(y) [x(t) - x_0(t)]^2. \quad (5)$$

The proof of this theorem is based on a direct verification of the interpolation conditions (4).

Substituting the approximation $H(x, y)$ of the form (5) to the solution $u(x, y)$ of the equation (2) in the left-hand side of equality (2), we obtain

$$\frac{\delta^2 H(x, y)}{\delta x^2(t)} - a^2(t) \frac{\delta^2 H(x, y)}{\delta y^2(t)} =$$

$$= -a^2(t) \left(u_1''(y) \int_a^b (x(t) - x_0(t)) dt + \frac{1}{2} a^2(t) u_0^{(4)}(y) [x(t) - x_0(t)]^2 \right) \delta^2(t-s),$$

where the delta function $\delta(t) = \begin{cases} 0, & t \neq 0; \\ +\infty, & t = 0. \end{cases}$ We note that in the case $t \neq s$, the value $\delta(t-s) = 0$ and the equality (2) takes place for any (x, y) from the domain of definition.

The obtained results can serve as a basis for further research of the theory of differential equations with variational derivatives that is not well developed, and can also be used to construct approximate interpolation methods for solving some linear and nonlinear differential equations with variational derivatives of the first and second order that are found in various applied fields and mathematical physics.

REFERENCES

- [1] Paul Le'vy. *Concrete Problems of Functional Analysis*. Moscow : Nauka Publ., 1967.
- [2] Mordecai M. Vainberg. *Variational Methods for Investigation of Non-linear Operators*. Moscow : Gosudarstvennoye tekhnicheskoye izdatel'stvo Publ., 1956.
- [3] Vito Volterra. *Theory of Functionals and of Integral and Integro-Differential Equations*. New York : Dover Publications, 2005.
- [4] Yuri L. Daletsky. Differential Equations with Functional Derivatives and Stochastic Equations for Generalized Random Processes. *Doklady of the Academy of Sciences of USSR*, 166(5) : 1035–1038, 1966.
- [5] Vladimir G. Zadorozhniy. Second-order Differential Equations with Variational Derivatives. *Differential equations*, 25(10) : 1679–1683, 1989.
- [6] Vera P. Danilovich, Ivan M. Kovalchik. Cauchy Formula for Linear Equations with Functional Derivatives. *Differential equations*, 13(8) : 1509–1511, 1977.
- [7] Ivan M. Kovalchik. Representation of Solutions of Certain Equations with Functional Derivatives Using Wiener Integrals. *Doklady of the Academy of Sciences of Ukraine. Series A. Physical, Mathematical, and Technical Sciences*, 12 : 1079–1083, 1978.
- [8] Ivan M. Kovalchik. Linear Equations with Functional Derivatives. *Doklady of the Academy of Sciences of USSR*, 194(4) : 763–766, 1970.
- [9] Yuri L. Daletsky. Infinite-dimensional Elliptic Operators and Parabolic Equations Connected with them. *Successes of Mathematical Sciences*, 22(4(136)) : 3–54, 1967.
- [10] Vladimir I. Averbukh, Oieg G. Smolyanov. The Theory of Differentiation in Linear Topological Spaces. *Successes of Mathematical Sciences*, 22(6(138)) : 201–260, 1967.
- [11] Vladimir L. Makarov, Vladimir V. Khlobystov, Leonid A. Yanovich. *Methods of Operator Interpolation*, volume 83 of *Proceedings of the Institute of Mathematics of the National Academy of Sciences of Ukraine*. Kiev : Institute of Mathematics of the National Academy of Sciences of Ukraine, 2010.
- [12] Leonid A. Yanovich, Marina V. Ignatenko. *Bases of the theory of interpolation of functions of matrix variables*. Minsk : Belaruskaya Nabuka Publ., 2016.

I. Gelbukh <i>Realization of a graph as the Reeb graph of a Morse, Morse–Bott or round function</i>	38
N. Glazunov <i>Foliations in moduli spaces of abelian varieties and bounded global G-shtukas</i>	40
A. Glebova <i>Geometry of curves in three-dimensional space and invariants of nonlinear differential equations of the second order</i>	41
P. Petrenko, A. Andreev <i>On Orthosymmetric n-morphisms</i>	42
M. Golasinski <i>On homotopy nilpotency of Moore space</i>	43
E. Afanas'eva, A. Golberg <i>Metric viewpoint in mapping theory between Riemannian manifolds</i>	45
S. V. Gryshchuk <i>Monogenic functions with values in commutative complex algebras of the second rank with unity and generalized biharmonic equation with non-zero simple characteristics</i>	46
A. Guale, J. Vielma <i>The Collatz conjecture from an algebraic point of view</i>	47
O. Gutik, P. Khylynskyi <i>On the monoid of cofinite partial isometries of positive integers with a bounded finite noise</i>	48
O. Gutik, M. Mykhalenych <i>On some generalization of the bicyclic monoid</i>	50
Y. Gutman <i>Variational principles for metric mean dimension</i>	51
O. Ye. Hentosh, A. K. Prykarpatski <i>A generalized Lie-algebraic approach to constructing of integrable fractional dynamical systems</i>	52
L. Yanovich, M. Ignatenko <i>On the solution of separate differential equations with variational derivatives of the first and second orders</i>	54
Sh. Sh. Ismoilov <i>Special mean and total curvature of a dual surface in isotropic spaces</i>	57
D. Jumaev, A. Zaitov <i>A topological transformation group of a hyperspace</i>	59
D. A. Juraev <i>On the Carleman function for matrix factorizations of the Helmholtz equation</i>	62
A. Kachurovskii <i>Measuring the rate of convergence in the Birkhoff ergodic theorem</i>	64
O. Kadubov's'kyi <i>Enumeration of topologically non-equivalent functions with one degenerate saddle critical point on triple torus</i>	65
O. Karupu <i>On some properties of moduli of smoothness of conformal mapping of simply connected domains</i>	68
D. Katz <i>Integration over non-rectifiable curves: spirals of high torsion</i>	70
A. A. Katz <i>On real Σ^*-algebras of operators</i>	72
L. Fardigola, K. Khalina <i>On controllability problems for the heat equation with variable coefficients controlled by the Dirichlet boundary condition on a half-axis</i>	73