

International scientific conference
**«Algebraic and geometric
methods of analysis»**

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

<https://www.imath.kiev.ua/~topology/conf/agma2018>

Problem on non-overlapping polycylindrical domains with poles on the boundary of a polydisk

Iryna Denega

(Institute of mathematics of the NAS of Ukraine, Department of complex analysis and potential theory, Tereshchenkivska 3, Kiev-4, 01004, Ukraine)

E-mail: iradenega@gmail.com

The goal of the present work is the study of the problem of a product of powers of the generalized conformal radii of polycylindrical nonoverlapping domains with poles on the boundary of a polydisk. The spatial analogs of a number of known results concerning the nonoverlapping domains on a plane were obtained in [1], where a generalization of the notion of inner radius was given. Namely, the notion of harmonic radius of the spatial domain $B \subset \mathbb{R}^n$ relative to some internal point was introduced. Work [1] was the essential break-through in the consideration of nonoverlapping domains in the spatial case. Then, work [2] advanced an approach that allowed the transfer of some results known in the case of a complex plane onto \mathbb{C}^n . At the same time, the problems of nonoverlapping domains in the case of a complex plane represent a sufficiently well-developed trend of the geometric theory of functions of complex variable (see, e.g., [1–6]).

Let \mathbb{N} , \mathbb{R} and \mathbb{C} be the sets of natural, real, and complex numbers, respectively, and $\mathbb{R}^+ = (0, \infty)$. Let $\overline{\mathbb{C}}$ be a Riemann sphere (extended complex plane). It is well known that $\mathbb{C}^n = \underbrace{(\mathbb{C} \times \mathbb{C} \times \dots \times \mathbb{C})}_{n\text{-times}}$, $n \in \mathbb{N}$. $\overline{\mathbb{C}}^n = \underbrace{(\overline{\mathbb{C}} \times \overline{\mathbb{C}} \times \dots \times \overline{\mathbb{C}})}_{n\text{-times}}$ is a compactification of the space \mathbb{C}^n (see, e.g., [3, 4, 5]), where

the set of infinitely remote points has the complex dimension $n - 1$. Let $[D]^n$ (Cartesian degree of a domain $D \in \overline{\mathbb{C}}$) denote the Cartesian product $\underbrace{D \times D \times \dots \times D}_{n\text{-times}}$, and let $[d]^n$ (Cartesian degree of a point $d \in \overline{\mathbb{C}}$) denote the point with $\overline{\mathbb{C}}^n$, which have the coordinates $\underbrace{(d, \dots, d)}_{n\text{-times}}$. It is clear that $\mathbb{C}^1 = \mathbb{C}$,

$\overline{\mathbb{C}}^1 = \overline{\mathbb{C}}$. The topology in $\overline{\mathbb{C}}^n$ is introduced like in a Cartesian product of topological spaces. In this topology, $\overline{\mathbb{C}}^n$ is compact (see [3, 4, 5]).

The domain $\mathbb{B} = B_1 \times B_2 \times \dots \times B_n \subset \overline{\mathbb{C}}^n$, where each domain $B_k \subset \overline{\mathbb{C}}$, $k = \overline{1, n}$, is called a polycylindrical domain in $\overline{\mathbb{C}}^n$ (see [3]). The domains B_k , $k = \overline{1, n}$, are called coordinate domains of the domain \mathbb{B} .

Let $r(B, a)$ be inner radius of the domain $B \subset \overline{\mathbb{C}}$ with respect to the point $a \in B$. The generalized inner radius of the polycylindrical domain \mathbb{B} relative to the point $\mathbb{A} = (a_1, a_2, \dots, a_n) \in \mathbb{B}$, $a_k \in B_k$, $k = \overline{1, n}$, is

$$R(\mathbb{B}, \mathbb{A}) := \left(\prod_{k=1}^n r(B_k, a_k) \right)^{\frac{1}{n}},$$

where the quantities $r(B_k, a_k)$, $k = \overline{1, n}$, mean the inner radii of the coordinate domains B_k relative to a_k .

Let $\mathbb{U}^n = [U]^n$, where $U = \{z \in \mathbb{C} : |z| < 1\}$ (unit disk in the complex plane \mathbb{C}). By Γ_n we denote the skeleton of the polydisk \mathbb{U}^n i.e., the set of points $\mathbb{A} = (a_1, a_2, \dots, a_n) \in \mathbb{C}^n$, $|a_s| = 1$, $s = \overline{1, n}$.

The system $\{\mathbb{B}_k\}_{k=1}^m$ ($\mathbb{B}_k = B_1^{(k)} \times \dots \times B_n^{(k)}$, $k = \overline{1, m}$) is called a system of nonoverlapping polycylindrical domains, if, for every fixed p_0 , $p_0 = \overline{1, n}$, the system of domains $\{B_{p_0}^{(k)}\}$, $k = \overline{1, m}$, is a system of nonoverlapping domains on $\overline{\mathbb{C}}$.

Let $m \in \mathbb{N}$, $m \geq 2$. The system of points $\Delta_m := \{a_k\}_{k=1}^m$, $a_k \in \mathbb{C}$, is called m -radial, if $|a_k| \in \mathbb{R}^+$ for $k = \overline{1, m}$, $0 = \arg a_1 < \arg a_2 < \dots < \arg a_m < 2\pi$.

The system of points $\{\mathbb{A}_k\}_{k=1}^m$ ($\mathbb{A}_k = (a_1^{(k)}, a_2^{(k)}, \dots, a_n^{(k)}) \in \mathbb{C}^n$, $k = \overline{1, m}$), is called radial in the space \mathbb{C}^n , if, for every fixed p_0 the sequence $\{a_{p_0}^{(k)}\}$, $k = \overline{1, m}$, is an m -radial system of points on the corresponding complex plane \mathbb{C} .

We will consider radial systems of points in the space \mathbb{C}^n of the form

$$\{\mathbb{A}_k\}_{k=1}^m, \quad \mathbb{A}_k = (a_1^{(k)}, a_2^{(k)}, \dots, a_n^{(k)}) \in \mathbb{C}^n, \quad k = \overline{1, m}, \quad a_{p_0}^{(1)} > 0, \quad p_0 = \overline{1, n},$$

$$\arg a_{p_0}^{(k)} < \arg a_{p_0}^{(k+1)}, \quad k = \overline{1, m-1}, \quad \arg a_{p_0}^{(m)} < 2\pi. \quad (1)$$

We introduce the following notations

$$\alpha_{p_0}^{(1)} := \frac{1}{\pi} \left(\arg a_{p_0}^{(2)} - \arg a_{p_0}^{(1)} \right), \quad \alpha_{p_0}^{(2)} := \frac{1}{\pi} \left(\arg a_{p_0}^{(3)} - \arg a_{p_0}^{(2)} \right), \dots, \alpha_{p_0}^{(m)} := \frac{1}{\pi} \left(2\pi - \arg a_{p_0}^{(m)} \right).$$

Let

$$F_\delta(x) = 2^{x^2+6} \cdot x^{x^2+2-2\delta} \cdot (2-x)^{-\frac{1}{2}(2-x)^2} (2+x)^{-\frac{1}{2}(2+x)^2},$$

$$x \in (0, 2], \quad \delta \in [0, 1], \quad F_\delta(x) \subset \overline{\mathbb{C}}.$$

Then the following proposition is valid.

Theorem 1. [6] *Let $m, n \in \mathbb{N}$, $m \geq 7$, $\gamma \in (0, \gamma_0]$, $\gamma_0 = \sqrt[3]{m}$ and $\delta \in [0; 0, 7]$. Then, for any radial system of points of the form (1) $\{\mathbb{A}_k\}_{k=1}^m = \{a_p^{(k)}\}_{k=1}^m \in \overline{\mathbb{C}^n}$, $p = \overline{1, n}$, such that $\mathbb{A}_k \in \Gamma_n$, $k = \overline{1, m}$, and for any collection of nonoverlapping polycylindrical domains \mathbb{B}_k , $\mathbb{A}_k \in \mathbb{B}_k \subset \overline{\mathbb{C}^n}$, $k = \overline{0, m}$, $\mathbb{A}_0 \in \mathbb{B}_0 \subset \overline{\mathbb{C}^n}$, the following inequality holds*

$$R^\gamma(\mathbb{B}_0, \mathbb{A}_0) \prod_{k=1}^m R(\mathbb{B}_k, \mathbb{A}_k) \leq \gamma^{-\frac{\delta \cdot m}{2}} \cdot \left(\prod_{k=1}^m \prod_{p=1}^n \alpha_p^{(k)} \right)^{\frac{\delta}{n}} \cdot \left[F_\delta \left(\frac{2}{m} \sqrt{\gamma} \right) \right]^{\frac{m}{2}}.$$

One of the extreme systems is the system

$$\{\mathbb{B}_k\}_{k=0}^m = \{[B_0^{(0)}]^n, [B_1^{(0)}]^n, [B_2^{(0)}]^n, \dots, [B_m^{(0)}]^n\},$$

$$\{\mathbb{A}_k\}_{k=0}^m = \{[0]^n, [b_1^{(0)}]^n, [b_2^{(0)}]^n, \dots, [b_m^{(0)}]^n\},$$

where the domains $B_k^{(0)}$ and the points $b_k^{(0)}$, $k = \overline{1, m}$, are, respectively, the circular domains and the poles of the quadratic differential

$$Q(w)dw^2 = -\frac{(n^2 - \gamma)w^n + \gamma}{w^2(w^n - 1)^2} dw^2.$$

REFERENCES

- [1] V. N. Dubinin and E. G. Prilepkina. On the extreme partition of spatial domains. *Zap. Nauch. Sem. POMI*, 254 : 95–107, 1998.
- [2] A. K. Bakhtin. A generalization of some results of the theory of one-sheeted functions onto multidimensional complex spaces. *Dop. NAN Ukr.*, (3) : 7–11, 2011.
- [3] B. V. Shabat. *Introduction to Complex Analysis, Part II*, Amer. Math. Soc. Providence, RI, 1992.
- [4] E. M. Chirka. *Complex Analytic Sets*, Springer : Berlin, 1989.
- [5] B. A. Fuks. *Introduction to the Theory of Analytic Functions of Several Complex Variables*, Amer. Math. Soc. Providence, RI, 1963.
- [6] Iryna Denega, Yaroslav Zabolotnyi. Problem of nonoverlapping polycylindrical domains with poles on the boundary of a polydisk. *Journal of Mathematical Sciences*, 227(1) : 26–32, 2017.

Зміст

N. Aygor, H. Burhanzade <i>Secondary school students' misconceptions about linear algebra</i>	3
S. Bardyla, H. Kvasnytsia <i>Semitopological graph inverse semigroups</i>	4
B. A. Bhayo <i>On inequalities of generalized elliptic integrals</i>	5
Bodzioch M., Choiński M., Foryś U. <i>A criss-cross model of tuberculosis for heterogenous population</i>	6
Bolotov D. V. <i>Foliations with leaves of non-positive curvature and bounded total curvature on closed 3-manifolds</i>	7
E. Bonacci <i>Algebraic and geometric questions about a 6D physics</i>	9
F. Bulnes <i>Mukai-Fourier Transform in Derived Categories to Solutions of the Field Equations: Gravitational Waves as Oscillations in the Space-Time Curvature/Spin IV</i>	10
H. Burhanzade, N. Aygor <i>A study on the teaching methods in determinants</i>	12
Damla Yaman <i>Order continuity properties of lattice ordered algebras</i>	13
Denega I. <i>Problem on non-overlapping polycylindrical domains with poles on the boundary of a polydisk</i>	14
A. Dudko, V. Pivovarchik <i>Inverse three spectra problem for a Stieltjes string with the Neumann boundary conditions</i>	16
Eftekharinasab K. <i>On the existence of a global diffeomorphism between Fréchet spaces</i>	18
Glazunov N. <i>Class groups of rings with divisor theory, L-functions and moduli spaces</i>	19
O. Gok <i>b-bimorphisms</i>	21
Gei E. <i>On the second regularized trace formula for a differential operator with unbounded coefficients</i>	22
Hentosh O. Ye., Prykaratsky Ya. A. <i>The Lie-algebraic structure of the Lax-Sato integrable superanalogs for the Liouville heavenly type equations</i>	24
V. Herasymov <i>In a natural topological sense a typical linear nonhomogeneous differential equation in the ring $Z[[x]]$ has no solutions from $Z[[x]]$.</i>	26
Juraev D. A. <i>On the Cauchy problem for matrix factorizations of the Helmholtz equation</i>	27
M. E. Kansu <i>Macroscopic electromagnetism via complex quaternions</i>	29
Vladimir V. Kisil <i>An extension of Möbius–Lie geometry with conformal ensembles of cycles</i>	30
Konovenko N., Lychagin V. <i>Rational differential invariants for oriented primary visual cortex</i>	32