


International
Online Conference



**Algebraic
and Geometric
Methods of Analysis**

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

SCIENTIFIC COMMITTEE

Drozd Yu.

(Kyiv, Ukraine)

Maksymenko S.

(Kyiv, Ukraine)

Plaksa S.

(Kyiv, Ukraine)

Prishlyak A.

(Kyiv, Ukraine)

Bakhtin O.

(Kyiv, Ukraine)

Balan V.

(Bucharest, Romania)

Banakh T.

(Lviv, Ukraine)

Borysenko O.

(Kharkiv, Ukraine)

Cherevko Ye.

(Odesa, Ukraine)

Fedchenko Yu.

(Odesa, Ukraine)

Karlova O.

(Chernivtsi, Ukraine)

Kiosak V.

(Odessa, Ukraine)

Konovenko N.

(Odessa, Ukraine)

Lyubashenko V.

(Kyiv, Ukraine)

Matsumoto K.

(Yamagata, Japan)

Mormul P.

(Warsaw, Poland)

Mykhailyuk V.

(Chernivtsi, Ukraine)

Plachta L.

(Krakov, Poland)

Pokas S.

(Odessa, Ukraine)

Sabitov I.

(Moscow, Russia)

Savchenko O.

(Kherson, Ukraine)

Sergeeva A.

(Odessa, Ukraine)

Shelekhov A.

(Tver, Russia)

Zarichnyi M.

(Lviv, Ukraine)

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

ORGANIZING COMMITTEE

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohlyk O.

Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.

Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

Asymptotic estimates for the widths of classes of periodic functions of high smoothness

Anatolii Serdyuk

(Institute of Mathematics of NAS of Ukraine, Kyiv, Ukraine)

E-mail: serdyuk@imath.kiev.ua

Igor Sokolenko

(Institute of Mathematics of NAS of Ukraine, Kyiv, Ukraine)

E-mail: sokol@imath.kiev.ua

Let L_p , $1 \leq p \leq \infty$, and C be the spaces of 2π -periodic functions with standart norms $\|\cdot\|_p$ and $\|\cdot\|_C$, respectively.

Denote by $C_{\bar{\beta},p}^\psi$, $1 \leq p \leq \infty$, the set of all 2π -periodic functions f , representable as convolution

$$f(x) = \frac{a_0}{2} + \frac{1}{\pi} \int_{-\pi}^{\pi} \varphi(x-t) \Psi_{\bar{\beta}}(t) dt, \quad a_0 \in \mathbb{R}, \quad \varphi \in B_p^0 = \{g \in L_p : \|g\|_p \leq 1, g \perp 1\}, \quad (1)$$

with a fixed generated kernel $\Psi_{\bar{\beta}} \in L_{p'}$, $1/p + 1/p' = 1$, the Fourier series of which has the form

$$S[\Psi_{\bar{\beta}}](t) = \sum_{k=1}^{\infty} \psi(k) \cos\left(kt - \frac{\beta_k \pi}{2}\right), \quad \beta_k \in \mathbb{R}, \quad \psi(k) \geq 0. \quad (2)$$

A function f in the representation (1) is called $(\psi, \bar{\beta})$ -integral of the function φ and is denoted by $\mathcal{J}_{\bar{\beta}}^\psi \varphi$ ($f = \mathcal{J}_{\bar{\beta}}^\psi \varphi$). If $\psi(k) \neq 0$, $k \in \mathbb{N}$, then the function φ in the representation (1) is called $(\psi, \bar{\beta})$ -derivative of the function f and is denoted by $f_{\bar{\beta}}^\psi$ ($\varphi = f_{\bar{\beta}}^\psi$). The concepts of $(\psi, \bar{\beta})$ -integral and $(\psi, \bar{\beta})$ -derivative was introduced by Stepanets (see [4]). Since $\varphi \in L_p$ and $\Psi_{\bar{\beta}} \in L_{p'}$, then the function f of the form (1) is a continuous function, i.e. $C_{\bar{\beta},p}^\psi \subset C$ (see [4, Proposition 3.9.2.]).

In the case $\beta_k \equiv \beta$, $\beta \in \mathbb{R}$, the classes $C_{\bar{\beta},p}^\psi$ are denoted by $C_{\beta,p}^\psi$. For $\psi(k) = k^{-r}$, $r > 0$, the classes $C_{\bar{\beta},p}^\psi$ and $C_{\beta,p}^\psi$ are denoted by $W_{\bar{\beta},p}^r$ and $W_{\beta,p}^r$, respectively. The classes $W_{\beta,p}^r$ are the well-known Weyl-Nagy classes. For $\psi(k) = e^{-\alpha k^r}$, $\alpha > 0$, $r > 0$, the classes $C_{\bar{\beta},p}^\psi$ and $C_{\beta,p}^\psi$ are denoted by $C_{\bar{\beta},p}^{\alpha,r}$ and $C_{\beta,p}^{\alpha,r}$, respectively. The sets $C_{\beta,p}^{\alpha,r}$ are well-known classes of the generalized Poisson integrals.

Let \mathfrak{N} be some functional class from the space C ($\mathfrak{N} \subset C$). The quantity

$$E_n(\mathfrak{N})_C = \sup_{f \in \mathfrak{N}} E_n(f)_C = \sup_{f \in \mathfrak{N}} \inf_{T_{n-1} \in \mathcal{T}_{2n-1}} \|f - T_{n-1}\|_C \quad (3)$$

is called the best uniform approximation of the class \mathfrak{N} by elements of the subspace \mathcal{T}_{2n-1} of trigonometric polynomials T_{n-1} of the order $n - 1$.

The order estimates for the best approximations $E_n(K)_C$ of classes $K = C_{\bar{\beta},p}^\psi$, $1 \leq p \leq \infty$, (and, hence, classes $W_{\bar{\beta},p}^r$, $C_{\bar{\beta},p}^{\alpha,r}$ and $C_{\bar{\beta},p}^\psi$) depending on rate of decreasing to zero of sequences $\psi(k)$ were obtained, in particular, in the works of Temlyakov (1993), Hrabova and Serdyuk (2013), Serdyuk and Stepanyuk (2014) etc.

If the sequences $\psi(k)$ decrease to zero faster than any geometric progression, then asymptotic equations of the best uniform approximations are even known (see [3] and the bibliography available there).

In [3] it was shown that for such classes $C_{\bar{\beta},p}^\psi$ the following asymptotic equations take places

$$E_n(C_{\bar{\beta},p}^\psi)_C \sim \mathcal{E}_n(C_{\bar{\beta},p}^\psi)_C \sim \frac{\|\cos t\|_{p'}}{\pi} \psi(n), \quad 1 \leq p \leq \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1, \quad (4)$$

where $\mathcal{E}_n(C_{\bar{\beta},p}^\psi) = \sup_{f \in C_{\bar{\beta},p}^\psi} \|f - S_{n-1}(f)\|_C$, $S_{n-1}(f)$ is the partial Fourier sum of order $n-1$ of the function f , and $A(n) \sim B(n)$ as $n \rightarrow \infty$ means that $\lim_{n \rightarrow \infty} A(n)/B(n) = 1$.

For $p = \infty$ in the case of $K = W_{\bar{\beta},\infty}^r$, $r > 0$, and in the cases of $K = C_{\bar{\beta},\infty}^{\alpha,r}$, $r \geq 1$, and $K = C_{\bar{\beta},\infty}^\psi$ ($K = C_{\bar{\beta},\infty}^\psi$) for certain restrictions on sequences ψ and $\bar{\beta}$ the exact values of the best uniform approximations are known thanks to the works of Favard (1936, 1937), Akhiezer and Krein (1937), Krein (1938), Nagy (1938), Stechkin (1956), Dziadyk (1959, 1974), Sun (1961), Bushanskij (1978), Pinkus (1985), Serdyuk (1995, 1999, 2002) etc.

For $p = 2$ and for arbitrary $\bar{\beta} = \beta_k \in \mathbb{R}$, $\sum_{k=1}^{\infty} \psi^2(k) < \infty$ the exact values for the quantity $\mathcal{E}_n(C_{\bar{\beta},2}^\psi)_C$ are also known (see [2]).

Let K be a convex centrally symmetric subset of C and let $b_N(K, C)$, $d_N(K, C)$, $\lambda_N(K, C)$, and $\pi_N(K, C)$ be Bernstein, Kolmogorov, linear, and projection N -widths of the set K in the space C [1].

The results containing order estimates of the widths b_N , d_N , λ_N or π_N in the case of $K = C_{\bar{\beta},p}^\psi$ (and, in particular, $W_{\bar{\beta},p}^r$ and $C_{\bar{\beta},p}^\psi$) can be found, for example, in the works of Tikhomirov (1976), Pinkus (1985), Kornejchuk (1987), Kashin (1977), Kushpel' (1989), Temlyakov (1990, 1993) etc.

Theorem 1. *Let $\{\beta_k\}_{k=1}^{\infty}$, $\beta_k \in \mathbb{R}$, and $\psi(k) > 0$ satisfies the condition $\sum_{k=1}^{\infty} \psi^2(k) < \infty$. Then for all $n \in \mathbb{N}$ the following inequalities hold*

$$\frac{1}{\sqrt{\pi}} \left(\frac{1}{\psi^2(n)} + 2 \sum_{k=1}^{n-1} \frac{1}{\psi^2(k)} \right)^{-\frac{1}{2}} \leq P_{2n}(C_{\bar{\beta},2}^\psi, C) \leq P_{2n-1}(C_{\bar{\beta},2}^\psi, C) \leq \frac{1}{\sqrt{\pi}} \left(\sum_{k=n}^{\infty} \psi^2(k) \right)^{\frac{1}{2}}, \quad (5)$$

where P_N is any of the widths b_N , d_N , λ_N or π_N .

If, in addition, $\psi(k)$ satisfies the condition $\lim_{n \rightarrow \infty} \max \left\{ \psi(n) \left(\sum_{k=1}^{n-1} \frac{1}{\psi^2(k)} \right)^{\frac{1}{2}}, \frac{1}{\psi(n)} \left(\sum_{k=n+1}^{\infty} \psi^2(k) \right)^{\frac{1}{2}} \right\} = 0$, then the following asymptotic equalities hold

$$\left. \begin{array}{l} P_{2n}(C_{\bar{\beta},2}^\psi, C) \\ P_{2n-1}(C_{\bar{\beta},2}^\psi, C) \end{array} \right\} = \psi(n) \left(\frac{1}{\sqrt{\pi}} + \mathcal{O}(1) \max \left\{ \psi(n) \left(\sum_{k=1}^{n-1} \frac{1}{\psi^2(k)} \right)^{\frac{1}{2}}, \frac{1}{\psi(n)} \left(\sum_{k=n+1}^{\infty} \psi^2(k) \right)^{\frac{1}{2}} \right\} \right), \quad (6)$$

where $\mathcal{O}(1)$ are the quantities uniformly bounded in all parameters.

The equalities (6) are realized by trigonometric Fourier sums $S_{n-1}(f)$.

REFERENCES

- [1] A. Pinkus, *n*-widths in approximation theory, Springer-Verlag, Berlin, 1985.
- [2] A.S. Serdyuk and I.V. Sokolenko, *Uniform approximation of classes of $(\psi, \bar{\beta})$ -differentiable functions by linear methods*, Approximation Theory of Functions and Related Problems, Zb. prac' Inst. mat. NAN Ukr., Kyiv, **8**, 2011, no. 1, P. 181–189. (Ukrainian)
- [3] A.S. Serdyuk, I.V. Sokolenko, *Asymptotic estimates for the best uniform approximations of classes of convolution of periodic functions of high smoothness* // Journal of Mathematical Sciences, Vol. 252, No. 4, 2021, P. 526–540.

[4] A.I. Stepanets, *Methods of Approximation Theory*, Utrecht, VSP, 2005.

ІНСТИТУТ МАТЕМАТИКИ НАН України

Y. Chapovskyi, D. Efimov, A. Petravchuk <i>Centralizers of elements in Lie algebras of vector fields</i>	113
E. Petrov, R. Salimov <i>Properties of quasisymmetric mappings to preserve the structures of spaces</i>	114
D. R. Popovych <i>IW contractions and their generalizations</i>	116
Prabhjot Singh <i>Weak Separation Condition coincides Finite Type Condition</i>	118
Pratyush Pranav <i>Topological data analysis for cosmology: theory and applications</i>	120
A. Prishlyak, Ch. Hatamian <i>Morse flows with singularities on boundary of 3-manifolds</i>	121
V. Prokip <i>On the matrix equation $AX - YB = C$ over Bezout domains</i>	122
A. Prykarpatsky, I. Mykytyuk <i>On the metric equations generated by symplectic deformations on $P_2(C)$</i>	124
O. Reinov <i>On a result of G. Pisier concerning Sidon sets</i>	126
A. K. Sadullaev, F. G. Mukhamadiev <i>On the τ-placedness of space of the permutation degree</i>	127
T. N Safarov <i>The analogue of Darboux equation in Galilean space</i>	128
V. M. Safonov, I. V. Zamrii, O. V. Safonova <i>On countable multiplicity of mappings</i>	129
A. Savchenko <i>Fuzzy ultrametrization of spaces of non-additive measures on fuzzy ultrametric spaces</i>	130
O. Sazonova <i>About one class of continual approximate solutions with arbitrary density</i>	131
J. Segert <i>Painlevé VI Solutions From Equivariant ADHM Instanton Bundles</i>	133
A. Serdyuk, I. Sokolenko <i>Asymptotic estimates for the widths of classes of periodic functions of high smoothness</i>	135
H. Sinyukova <i>Some generalizations of the known theorems of the type of geodesical unique definability</i>	138
S. Som, A. Bera, L. K. Dey <i>Some remarks on the Metrizable of F-metric spaces</i>	139
P. Stegantseva, M. Grechneva <i>The surfaces with the flat normal connection and the constant curvature of Grassmann image in Minkowski space</i>	140
A. Skryabina, P. Stegantseva <i>The relation between T_0-topologies with the weight $2^{n-2} < k \leq 2^{n-1}$ on n-element set and T_0-topologies close to the discrete on $(n - 1)$-element set</i>	142
Ya. B. Stelmakh <i>The Golomb and Kirch topologies on the set of nonzero integers</i>	143
D. Dmitrishin, A. Stokolos <i>On symmetrization of univalent polynomials</i>	144