

International scientific conference
«Algebraic and geometric methods
of analysis»

Book of abstracts



May 31 - June 5, 2017
Odessa
Ukraine

LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences
- History and methodology of teaching in mathematics

ORGANIZERS

- The Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- The Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- The International Geometry Center

PROGRAM COMMITTEE

Chairman: Prishlyak A. (<i>Kyiv, Ukraine</i>)	Maksymenko S. (<i>Kyiv, Ukraine</i>)	Rahula M. (<i>Tartu, Estonia</i>)
Balan V. (<i>Bucharest, Romania</i>)	Matsumoto K. (<i>Yamagata, Japan</i>)	Sabitov I. (<i>Moscow, Russia</i>)
Banakh T. (<i>Lviv, Ukraine</i>)	Mashkov O. (<i>Kyiv, Ukraine</i>)	Savchenko A. (<i>Kherson, Ukraine</i>)
Fedchenko Yu. (<i>Odesa, Ukraine</i>)	Mykytyuk I. (<i>Lviv, Ukraine</i>)	Sergeeva A. (<i>Odesa, Ukraine</i>)
Fomenko A. (<i>Moscow, Russia</i>)	Milka A. (<i>Kharkiv, Ukraine</i>)	Strikha M. (<i>Kyiv, Ukraine</i>)
Fomenko V. (<i>Taganrog, Russia</i>)	Mikesh J. (<i>Olomouc, Czech Republic</i>)	Shvets V. (<i>Odesa, Ukraine</i>)
Glushkov A. (<i>Odesa, Ukraine</i>)	Mormul P. (<i>Warsaw, Poland</i>)	Shelekhov A. (<i>Tver, Russia</i>)
Haddad M. (<i>Wadi al-Nasara, Syria</i>)	Moskaliuk S. (<i>Wien, Austria</i>)	Shurygin V. (<i>Kazan, Russia</i>)
Herega A. (<i>Odesa, Ukraine</i>)	Panzhenskiy V. (<i>Penza, Russia</i>)	Vlasenko I. (<i>Kyiv, Ukraine</i>)
Khruslov E. (<i>Kharkiv, Ukraine</i>)	Pastur L. (<i>Kharkiv, Ukraine</i>)	Zadorozhnyj V. (<i>Odesa, Ukraine</i>)
Kirichenko V. (<i>Moscow, Russia</i>)	Plachta L. (<i>Krakov, Poland</i>)	Zarichnyi M. (<i>Lviv, Ukraine</i>)
Kirillov V. (<i>Odesa, Ukraine</i>)	Pokas S. (<i>Odesa, Ukraine</i>)	Zelinskiy Y. (<i>Kyiv, Ukraine</i>)
Konovenko N. (<i>Odesa, Ukraine</i>)	Polulyakh E. (<i>Kyiv, Ukraine</i>)	

ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Volkov V., Director of the Educational Research Institute of Mechanics, Automation and Computer Systems named after P. M. Platonov;
- Bukaros A., Dean of the Faculty of automation, mechatronics and robotics

ORGANIZING COMMITTEE

Kirillov V.
Konovenko N.
Fedchenko Yu.

Hladysh B.
Nuzhnaya N.
Osadchuk E.

Maksymenko S.
Khudenko N.
Cherevko E.

НТБ ОНАФТ

Landau-type inequalities for curves on Riemannian manifolds

Igor O. Parasyuk

(National Taras Shevchenko University of Kyiv, Volodymyrs'ka str., 64, Kyiv, Ukraine, 01601)

E-mail: pio@univ.kiev.ua

Let for a natural number n a function $f(\cdot) \in C^n(\mathbb{R} \mapsto \mathbb{R})$ satisfy the inequalities

$$\|f(\cdot)\|_\infty := \sup_{t \in \mathbb{R}} |f(t)| < \infty, \quad \|f^{(n)}(\cdot)\|_\infty < \infty.$$

In the case where $n = 2$, the famous Landau–Hadamard inequality reads

$$\|f'(\cdot)\|_\infty \leq \sqrt{2 \|f(\cdot)\|_\infty \|f''(\cdot)\|_\infty}.$$

In the general case $n \geq 2$, $1 \leq k < n$, Kolmogorov determined the best constants $C_{n,k}$ for inequality

$$\|f^{(k)}(\cdot)\|_\infty \leq C_{n,k} \|f(\cdot)\|_\infty^{1-k/n} \|f^{(n)}(\cdot)\|_\infty^{k/n}.$$

(see, e.g., [1]). The goal of the present report is to discuss how the above inequalities can be generalized for the case of mappings taking values in Riemannian manifolds.

Let $(\mathcal{M}, \mathfrak{g} = \langle \cdot, \cdot \rangle)$ be a smooth complete Riemannian manifold with the metric tensor \mathfrak{g} , and let ∇ be the Levi-Civita connection with respect to \mathfrak{g} . For a given smooth mapping $x(\cdot) : I \mapsto \mathcal{M}$ of an interval $I \subset \mathbb{R}$ and for a smooth vector field $\xi(\cdot) : I \mapsto T\mathcal{M}$ along $x(\cdot)$, denote by $\nabla_{\dot{x}}\xi(t)$ the covariant derivative of $\xi(\cdot)$ along the tangent vector $\dot{x}(t) \in T_{x(t)}\mathcal{M}$ at the point $t \in I$, and by $\nabla_{\dot{x}}^k$ the k -th iterate of $\nabla_{\dot{x}}$. (Here $T\mathcal{M} = \bigsqcup_{x \in \mathcal{M}} T_x\mathcal{M}$ stands for the total space of the tangent bundle with natural projection $\pi(\cdot) : T\mathcal{M} \mapsto \mathcal{M}$, and $T_x\mathcal{M} = \pi^{-1}(x)$ denotes the tangent space to \mathcal{M} at x .)

For a smooth function $U(\cdot) : \mathcal{M} \mapsto \mathbb{R}$ denote by $\nabla U(x) \in T_x\mathcal{M}$ and by $H_U(x) : T_x\mathcal{M} \mapsto T_x\mathcal{M}$, respectively, the gradient vector and the Hesse form of $U(\cdot)$ at point x (by the definition

$$\langle H_U(x)\xi, \eta \rangle = \langle \nabla_\xi \nabla U(x), \eta \rangle$$

for any $x \in \mathcal{M}$ and any $\xi, \eta \in T_x\mathcal{M}$). Define the natural norm for tangent vector ξ as $\|\xi\| := \sqrt{\langle \xi, \xi \rangle}$ and the norm for vector field $\xi(\cdot)$ along $x(\cdot)$ as

$$\|\xi(\cdot)\|_\infty := \sup_{t \in \mathbb{R}} \|\xi(t)\|.$$

We obtain the following Landau-type inequality.

Theorem 1. *Let $x(\cdot) : \mathbb{R} \mapsto \mathcal{M}$ be a smooth mapping such that*

$$\|\nabla_{\dot{x}}\dot{x}(\cdot)\|_\infty < \infty.$$

Suppose that there exists a smooth function $U(\cdot) : \mathcal{M} \mapsto \mathbb{R}$ satisfying the inequalities

$$\sup_{t \in \mathbb{R}} U \circ x(t) < \infty, \quad \|\nabla U \circ x(\cdot)\|_\infty < \infty,$$

and

$$\lambda := \inf_{t \in \mathbb{R}} \min \{ \langle [H_U \circ x(t)] \xi, \xi \rangle : \xi \in T_{x(t)}\mathcal{M}, \|\xi\| = 1 \} > 0.$$

Then

$$\|\dot{x}(\cdot)\|_\infty \leq C \sqrt{\|\nabla U \circ x(\cdot)\|_\infty \|\nabla_{\dot{x}}\dot{x}(\cdot)\|_\infty / \lambda}$$

where the constant C does not exceed the positive root of the equation $\zeta^3 - 3\zeta = 1$. In particular, $C < 1.87939$.

Remark 2. If $\mathcal{M} = \mathbb{R}^d$ and $U(x) := \|x\|^2/2$, then $\lambda = 1$, and Theorem 1 leads to the Landau inequality with the constant C somewhat greater than the best one $C_{2,1} = \sqrt{2}$.

If the closure of $x(\cdot)$ is a compact subset \mathcal{K} of a domain $\mathcal{D} \subset \mathcal{M}$ and there exists a point $x_0 \in \mathcal{D} \setminus \mathcal{K}$ such that the distance function

$$\rho(\cdot, x_0) : \mathcal{D} \setminus \{x_0\} \mapsto (0, \infty)$$

is smooth, then Theorem 1 holds true for $U(x) := \rho^2(x, x_0)$.

It turns out that it is much easier to obtain a counterpart of the Landau – Kolmogorov inequality for vector fields along mappings.

Theorem 3. Let $\xi(\cdot) : \mathbb{R} \mapsto T\mathcal{M}$ be a smooth vector field along a smooth mapping $x(\cdot) : \mathbb{R} \mapsto \mathcal{M}$ and let $n \geq 2$ be a natural number. Suppose that

$$\|\xi(\cdot)\|_\infty < \infty, \quad \|\nabla_x^n \xi(\cdot)\|_\infty < \infty.$$

Then for any natural $k < n$ there holds the inequality

$$\left\| \nabla_x^k \xi(\cdot) \right\|_\infty \leq C_{n,k} \|\xi(\cdot)\|_\infty^{1-k/n} \|\nabla_x^n \xi(\cdot)\|_\infty^{k/n}$$

where $C_{n,k}$ are the Kolmogorov constants.

REREFENCES

- [1] Steven R. Finch. *Mathematical constants*, volume 94 of *Encyclopedia of Mathematics and its Applications*. Cambridge: Cambridge University Press, 2003.

Konovenko N., Lychagin V. <i>On projective classes of rational functions</i>	71
Kozerenko S. <i>Orientations of trees and signed Markov graphs</i>	73
Kuzmenko T. <i>Constructive description of G-monogenic mappings in the algebra of complex quaternions</i>	74
Lyubashenko V. <i>Moyal and Rankin-Cohen deformations of algebras</i>	76
Markitan V. <i>Fractal properties of sets associated with Markov representation of real numbers defined by a double stochastic matrix</i>	78
Matsumoto K. <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds</i>	79
Mormul P. <i>Weak and strong nilpotentizability in the monster towers hosting flag distributions</i>	80
Mukhamadiev F. G. <i>The local density and the local weak density of $N_7^{\mathcal{O}}$-kernel of a topological space X and superextensions</i>	82
Muradoglu Z., Gunduz Aras C. <i>A study for decision making problems by using interval soft sets</i>	84
Muradov R. S. <i>Archimedean copula functions and their some algebraic properties with applications</i>	85
Obikhod T. V. <i>BPS states of Fourfolds as candidates for Kaluza-Klein modes</i>	87
Parasyuk I. O. <i>Landau-type inequalities for curves on Riemannian manifolds</i>	88
Prislyak A., Prus A. <i>Morse-Smale flows on torus with hole</i>	90
Reinov O. <i>On nuclear operators with trace $V = 1$ and $V^2 = 0$</i>	91
Sabitov I. Kh. <i>Multiple roots of the volume polynomials for polyhedra</i>	92
Samokhvalov S. <i>Theory of gravity in the affine frame</i>	93
Shamolin M. V. <i>Integrable systems with dissipation on the tangent bundle of two-dimensional manifold</i>	94
Turhan T., Ayyildiz N. <i>On geometry of spatial kinematics in Lorentzian space</i>	96
Turhan T., Ayyildiz N. <i>A study on the integral invariants of a closed spacelike ruled surface</i>	97
Vasilchenko A. N. <i>Dual modules over Steenrod algebra 2</i>	98
Vlasenko I. <i>Topology of the basin of attraction of surface endomorphisms.</i>	100
Voloshyna V. <i>About some properties of functions determined as transformations from W^n to W^m-representation</i>	101
Vyhivska L. <i>On the problem of product of inner radii symmetric non-overlapping domains</i>	103
Yildirim S., Ayyildiz N. <i>A Study on Rectifying Curves in Semi-Euclidean Spaces</i>	104
Арсеньева О. Е., Кириченко В. Ф., Суровцева Е. В. <i>Эрмитова геометрия почти контактного метрического многообразия</i>	105