

International scientific conference  
«Algebraic and geometric methods  
of analysis»

Book of abstracts



May 31 - June 5, 2017  
Odessa  
Ukraine

## LIST OF TOPICS

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences
- History and methodology of teaching in mathematics

## ORGANIZERS

- The Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- The Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- The International Geometry Center

## PROGRAM COMMITTEE

<b>Chairman: Prishlyak A.</b> ( <i>Kyiv, Ukraine</i> )	<b>Maksymenko S.</b> ( <i>Kyiv, Ukraine</i> )	<b>Rahula M.</b> ( <i>Tartu, Estonia</i> )
<b>Balan V.</b> ( <i>Bucharest, Romania</i> )	<b>Matsumoto K.</b> ( <i>Yamagata, Japan</i> )	<b>Sabitov I.</b> ( <i>Moscow, Russia</i> )
<b>Banakh T.</b> ( <i>Lviv, Ukraine</i> )	<b>Mashkov O.</b> ( <i>Kyiv, Ukraine</i> )	<b>Savchenko A.</b> ( <i>Kherson, Ukraine</i> )
<b>Fedchenko Yu.</b> ( <i>Odesa, Ukraine</i> )	<b>Mykytyuk I.</b> ( <i>Lviv, Ukraine</i> )	<b>Sergeeva A.</b> ( <i>Odesa, Ukraine</i> )
<b>Fomenko A.</b> ( <i>Moscow, Russia</i> )	<b>Milka A.</b> ( <i>Kharkiv, Ukraine</i> )	<b>Strikha M.</b> ( <i>Kyiv, Ukraine</i> )
<b>Fomenko V.</b> ( <i>Taganrog, Russia</i> )	<b>Mikesh J.</b> ( <i>Olomouc, Czech Republic</i> )	<b>Shvets V.</b> ( <i>Odesa, Ukraine</i> )
<b>Glushkov A.</b> ( <i>Odesa, Ukraine</i> )	<b>Mormul P.</b> ( <i>Warsaw, Poland</i> )	<b>Shelekhov A.</b> ( <i>Tver, Russia</i> )
<b>Haddad M.</b> ( <i>Wadi al-Nasara, Syria</i> )	<b>Moskaliuk S.</b> ( <i>Wien, Austria</i> )	<b>Shurygin V.</b> ( <i>Kazan, Russia</i> )
<b>Herega A.</b> ( <i>Odesa, Ukraine</i> )	<b>Panzhenskiy V.</b> ( <i>Penza, Russia</i> )	<b>Vlasenko I.</b> ( <i>Kyiv, Ukraine</i> )
<b>Khruslov E.</b> ( <i>Kharkiv, Ukraine</i> )	<b>Pastur L.</b> ( <i>Kharkiv, Ukraine</i> )	<b>Zadorozhnyj V.</b> ( <i>Odesa, Ukraine</i> )
<b>Kirichenko V.</b> ( <i>Moscow, Russia</i> )	<b>Plachta L.</b> ( <i>Krakov, Poland</i> )	<b>Zarichnyi M.</b> ( <i>Lviv, Ukraine</i> )
<b>Kirillov V.</b> ( <i>Odesa, Ukraine</i> )	<b>Pokas S.</b> ( <i>Odesa, Ukraine</i> )	<b>Zelinskiy Y.</b> ( <i>Kyiv, Ukraine</i> )
<b>Konovenko N.</b> ( <i>Odesa, Ukraine</i> )	<b>Polulyakh E.</b> ( <i>Kyiv, Ukraine</i> )	

## ADMINISTRATIVE COMMITTEE

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Volkov V., Director of the Educational Research Institute of Mechanics, Automation and Computer Systems named after P. M. Platonov;
- Bukaros A., Dean of the Faculty of automation, mechatronics and robotics

## ORGANIZING COMMITTEE

Kirillov V.  
Konovenko N.  
Fedchenko Yu.

Hladysh B.  
Nuzhnaya N.  
Osadchuk E.

Maksymenko S.  
Khudenko N.  
Cherevko E.

НТБ ОНАФТ

# Archimedean copula functions and their some algebraic properties with applications

Muradov Rustamjon Sobitkhonovich

(Tashkent University of Information Technologies named after Muhammad al-Khwarizmi)

*E-mail:* rustamjonmuradov@mail.ru

The word “copula” was first used in a mathematical sense by Sklar (1959, see [3]). But the functions themselves predate the use of the term, appearing in the work of Hoeffding, Fréchet, Dall’Aglia, and many others(see, [2]). Over the past forty years or so, copulas have played an important role in finance, engineering, bio-medical research, hydrology, climate and weather research, social research, econometrics, insurance, statistical research, dynamical systems and many areas. Axiomatically, a copula can be defined as follows.

**Definition 1.** A two-dimensional copula  $C$  is a mapping from  $I^2 = [0, 1] \times [0, 1]$  to  $I = [0, 1]$  which satisfies the following three conditions:

1.  $C(u, 0) = C(0, u) = 0$  for every  $u \in [0, 1]$ ;
2.  $C(u, 1) = C(1, u) = u$  for every  $u \in [0, 1]$ ;
3.  $C(u_2, v_2) - C(u_1, v_2) - C(u_2, v_1) + C(u_1, v_1) \geq 0$  for every  $u_1, v_1, u_2, v_2 \in [0, 1]$  satisfying  $u_1 \leq u_2, v_1 \leq v_2$ .

For example of copulas are the product or independence copula  $C_{\perp}(u, v) = uv$ , minimum copula  $C_{min}(u, v) = \min(u, v)$  and maximal copula  $C_{max}(u, v) = \max(u + v - 1, 0)$ . In figure 1 we present the graphs of the copulas  $C_{\perp}$ ,  $C_{min}$  and  $C_{max}$ .




FIGURE 1.1. Graphs of the copulas  $C_{\perp}$ ,  $C_{min}$  and  $C_{max}$ .

In this work, we discuss an important class of copulas known as Archimedean copulas. These copulas find a wide range of applications for a number of reasons: 1) the ease with which they can be constructed; 2) the great variety of families of copulas which belong to this class; and 3) the many nice properties possessed by the members of this class.

Let  $\varphi$  be a continuous, strictly decreasing function from  $[0, 1]$  to  $[0, \infty]$  such that  $\varphi(1) = 0$ .

**Definition 2.** The pseudo-inverse of  $\varphi$  is the function  $\varphi^{[-1]}$  with  $Dom\varphi^{[-1]} = [0, \infty]$  and  $Ran(f)$  given by

$$\varphi^{[-1]}(t) = \begin{cases} \varphi^{-1}(t), & 0 < t < \varphi(0), \\ 0, & \varphi(0) \leq t \leq \infty. \end{cases} \quad (1)$$

Note that  $\varphi^{[-1]}$  is continuous and non increasing on  $[0, \infty]$ , and strictly decreasing on  $[0, \varphi(0)]$ . Furthermore,

$$\varphi(\varphi^{[-1]}(t)) = \begin{cases} t, & 0 < t < \varphi(0), \\ 0, & \varphi(0) \leq t \leq \infty \end{cases} = \min(t, \varphi(0))$$

If  $\varphi(0) = \infty$ , then  $\varphi^{[-1]} = \varphi^{-1}$ .

**Definition 3.** Copulas of the form

$$C(u, v) = \varphi^{[-1]}(\varphi(u) + \varphi(v)), \quad (2)$$

are called Archimedean copulas, where the function  $\varphi$  is called a generator of the copula  $\varphi(1) = 0$ .

**Theorem 4.** Let  $\varphi$  be a continuous, strictly decreasing function from  $I$  to  $[0, \infty]$  such that  $\varphi(1) = 0$ , and let  $\varphi^{[-1]}$  be the pseudo-inverse of  $\varphi$  defined by (2) (taken from [2]). Then the function  $C$  given by (3) is a copula if and only if  $\varphi$  is convex.

We conclude this work with two theorems concerning some algebraic properties of Archimedean copulas.

**Theorem 5.** Let  $C$  be an Archimedean copula (taken from [2]) with generator  $\varphi$ . Then:

1.  $C$  is symmetric, i.e.,  $C(u, v) = C(v, u)$  for all  $u, v$  in  $I$ ;
2.  $C$  is associative, i.e.,  $C(C(u, v), w) = C(u, C(v, w))$  for all  $u, v, w$  in  $I$ ;
3. If  $c > 0$  is any constant, then  $c\varphi$  is also a generator of  $C$ .

For convenience, let  $\Omega$  denote the set of continuous strictly decreasing convex functions  $\varphi$  from  $I$  to  $[0, \infty]$  with  $\varphi(1) = 0$ . By now the reader is surely wondering about the meaning of the term ‘‘Archimedean’’ for these copulas. Recall the Archimedean axiom for the positive real numbers: If  $a, b$  are positive real numbers, then there exists an integer  $n$  such that  $na > b$ . An Archimedean copula behaves like a binary operation on the interval  $I$ , in that the copula  $C$  assigns to each pair  $u, v$  in  $I$  a number  $C(u, v)$  in  $I$ . From Theorem 5, we see that the ‘‘operation’’  $C$  is commutative and associative, and preserves order, i.e.,  $u_1 \leq u_2$  and  $v_1 \leq v_2$  implies  $C(u_1, v_1) \leq C(u_2, v_2)$ . Algebraists call  $(I, C)$  an ordered Abelian semigroup. For any  $u$  in  $I$ , we can define the  $C$ -powers  $u_C^n$  of  $u$  recursively:  $u_C^1 = u$ , and  $u_C^{n+1} = C(u, u_C^n)$ , note that  $u_C^2$  belongs to the diagonal section  $\delta_C(u)$  of  $C$ . The version of the Archimedean axiom for  $(I, C)$  is, for any two numbers  $u, v$  in  $(0, 1)$ , there exists a positive integer  $n$  such that  $u_C^n < v$ . The next theorem shows that Archimedean copulas satisfy this version of the Archimedean axiom and hence merit their name. The term ‘‘Archimedean’’ for these copulas was introduced in Ling (1965, see [2]).

**Theorem 6.** Let  $C$  be an Archimedean copula (taken from [2]) generated by  $\varphi$  in  $\Omega$ . Then for any  $u, v$  in  $I$ , there exists a positive integer  $n$  such that  $u_C^n < v$ .

In many applications, the random variables (r.v.-s) of interest represent the lifetimes of individuals or objects in some population. In survival analysis our interest focuses on a nonnegative r.v.-s denoting death times of biological organisms or failure times of mechanical systems. A difficulty in the analysis of survival data is the possibility that the survival times can be subjected to random censoring by other nonnegative r.v.-s and therefore we observe incomplete data. In article [1] we consider only right censoring model and problem of estimation of survival function when the survival times and censoring times are dependent and estimates of survival function assuming that the dependence structure is described by a known Archimedean copula function. We demonstrate almost sure asymptotic representation which provides a key tool for obtaining weak convergence result for estimator.

## REFERENCES

- [1] Abdushukurov A.A., Muradov R.S. Some algebraic properties of Archimedean copula functions and their applications in statistical estimation of the survival function, *New Trends in Mathematical Sciences*, 4(3): 88–98, 2016.
- [2] Nelsen R.B. An Introduction to Copulas. Springer, New York, 2006.
- [3] Sklar A. Fonctions de repartition à  $n$  dimensions et leurs marges, Vol.8. of *Publications de l'Institut de Statistique de l'Université de Paris*, 229–231, 1959.

<b>Konovenko N., Lychagin V.</b> <i>On projective classes of rational functions</i>	<b>71</b>
<b>Kozerenko S.</b> <i>Orientations of trees and signed Markov graphs</i>	<b>73</b>
<b>Kuzmenko T.</b> <i>Constructive description of <math>G</math>-monogenic mappings in the algebra of complex quaternions</i>	<b>74</b>
<b>Lyubashenko V.</b> <i>Moyal and Rankin-Cohen deformations of algebras</i>	<b>76</b>
<b>Markitan V.</b> <i>Fractal properties of sets associated with Markov representation of real numbers defined by a double stochastic matrix</i>	<b>78</b>
<b>Matsumoto K.</b> <i>Warped product semi-slant submanifolds in locally conformal Kaehler manifolds</i>	<b>79</b>
<b>Mormul P.</b> <i>Weak and strong nilpotentizability in the monster towers hosting flag distributions</i>	<b>80</b>
<b>Mukhamadiev F. G.</b> <i>The local density and the local weak density of <math>N_7^{\mathcal{O}}</math>-kernel of a topological space <math>X</math> and superextensions</i>	<b>82</b>
<b>Muradoglu Z., Gunduz Aras C.</b> <i>A study for decision making problems by using interval soft sets</i>	<b>84</b>
<b>Muradov R. S.</b> <i>Archimedean copula functions and their some algebraic properties with applications</i>	<b>85</b>
<b>Obikhod T. V.</b> <i>BPS states of Fourfolds as candidates for Kaluza-Klein modes</i>	<b>87</b>
<b>Parasyuk I. O.</b> <i>Landau-type inequalities for curves on Riemannian manifolds</i>	<b>88</b>
<b>Prislyak A., Prus A.</b> <i>Morse-Smale flows on torus with hole</i>	<b>90</b>
<b>Reinov O.</b> <i>On nuclear operators with trace <math>V = 1</math> and <math>V^2 = 0</math></i>	<b>91</b>
<b>Sabitov I. Kh.</b> <i>Multiple roots of the volume polynomials for polyhedra</i>	<b>92</b>
<b>Samokhvalov S.</b> <i>Theory of gravity in the affine frame</i>	<b>93</b>
<b>Shamolin M. V.</b> <i>Integrable systems with dissipation on the tangent bundle of two-dimensional manifold</i>	<b>94</b>
<b>Turhan T., Ayyildiz N.</b> <i>On geometry of spatial kinematics in Lorentzian space</i>	<b>96</b>
<b>Turhan T., Ayyildiz N.</b> <i>A study on the integral invariants of a closed spacelike ruled surface</i>	<b>97</b>
<b>Vasilchenko A. N.</b> <i>Dual modules over Steenrod algebra 2</i>	<b>98</b>
<b>Vlasenko I.</b> <i>Topology of the basin of attraction of surface endomorphisms.</i>	<b>100</b>
<b>Voloshyna V.</b> <i>About some properties of functions determined as transformations from <math>W^n</math> to <math>W^m</math>-representation</i>	<b>101</b>
<b>Vyhivska L.</b> <i>On the problem of product of inner radii symmetric non-overlapping domains</i>	<b>103</b>
<b>Yildirim S., Ayyildiz N.</b> <i>A Study on Rectifying Curves in Semi-Euclidean Spaces</i>	<b>104</b>
<b>Арсеньева О. Е., Кириченко В. Ф., Суровцева Е. В.</b> <i>Эрмитова геометрия почти контактного метрического многообразия</i>	<b>105</b>